

Cyclization of *ortho*-Halogenated *N*-Acylbenzylamines: a Formal Synthesis of ($\pm$)-Cherylline¹

By SATINDER V. KESSAR,* PARAMJIT SINGH, RAVINDER CHAWLA, and PAWAN KUMAR
(Department of Chemistry, Panjab University, Chandigarh, India)

Summary *ortho*-Halogenated *N*-alkyl-*N*-acylbenzylamines can be cyclised to dihydroisoquinolones by reaction with KNH₂ in liquid NH₃ or lithium di-isopropylamide in tetrahydrofuran under photolytic and thermal conditions; the procedure has been employed to synthesize ($\pm$)-cherylline.

WE have investigated the cyclization of α -carbanions derived from *ortho*-halogenated *N*-alkyl-*N*-acylbenzyl (**1a**) and phenethyl (**5**) amines as a route to heterocyclic systems, using the following conditions: (A) KNH₂ (8 mmol) in liq. NH₃ for 2 h; (B) KNH₂ (1 mmol) in liq. NH₃ under irradiation† for 8 min; (C) lithium di-isopropylamide (LDA, 4 mmol) in tetrahydrofuran (THF) under irradiation for 1.5 h; (D) LDA (4 mmol) in THF for 15 h at 25 °C. The results are presented in the Table.

It is clear that under typical aryne-generating conditions (A) cyclization is successful with the unsubstituted compound (**1a**) (R¹ = R² = H), but when alkoxy-substituents are present (R¹ = R² = MeO) amination predominates.² With such substrates the desired ring closure can be brought about by irradiation (B). Remarkably, when LDA was used as a base, the cyclization proceeded smoothly under photolytic (C) as well as thermal³ (D) conditions. Irradiation of (**1a**; R¹ = MeO, R² = PhCH₂O, R³ = *p*-MeOC₆H₄) in THF containing 4 mol. equiv. of LDA (C) gave the corresponding dihydroisoquinolone (**3**) in 48% yield, m.p. 163–165 °C; ¹H n.m.r. and mass spectral data were in conformity with the reported values⁴ and the m.p. was not depressed on admixture with an authentic sample. It could be converted into ($\pm$)-cherylline (**4**) by the known procedure.⁴

† Medium-pressure mercury lamp (100 W) in a Pyrex immersion-type vessel.




TABLE. Cyclization of *ortho*-halogenated *N*-acylbenzylamines (**1a**).

Compound				Reaction conditions	Major product ^a	% Yield ^b
R ¹	R ₂	R ³	X			
H	H	H	Cl	A	(3)	51
MeO	MeO	H	Cl	A	(1b)	29
MeO	PhCH ₂ O	<i>p</i> -PhCH ₂ OC ₆ H ₄	Br	A	(1b)	47
MeO	MeO	<i>p</i> -MeOC ₆ H ₄	Br	B	(3)	48
MeO	MeO	<i>p</i> -MeOC ₆ H ₄	Br	C	(3)	59
MeO	MeO	<i>p</i> -MeOC ₆ H ₄	Br	D	(3)	51
MeO	MeO	H	Br	C	(3)	45
MeO	PhCH ₂ O	<i>p</i> -MeOC ₆ H ₄	Br	C	(3)	48

^a All new products were characterised by elemental analyses and n.m.r. and mass spectral data. ^b Reactions were carried out on a 1 mmol scale and products isolated by column or thick-layer chromatography.

Surprisingly, in the phenethyl system ring closure through the nitrogen atom is preferred. Thus, treatment of the amide (**5**) with LDA under conditions (C) or (D) gave the base (**6**) (42%) which was characterised by air oxidation to *N*-methyl-5,6-dimethoxyindole; ¹H n.m.r. (CDCl₃)⁵ δ 3.80 (3H, s, >NMe), 4.0 and 4.05 (each 3H, s, OMe), 6.45 (1H, d, 3-H), 6.88 (1H, s, 7-H), 7.0 (1H, d, 2-H), and 7.2 (1H, s, 4-H); *m/e* 191.0948 (*M*⁺) (calc. 191.0946). Additional

studies on the synthetic scope and the mechanism of these reactions are in progress; depending on the reaction conditions and the substrate, both aryne and aryl radical intermediates (*S*_{RN}¹ type mechanism) seem to be involved.

We thank Professor D. A. Evans and Dr. W. E. Scott for authentic samples of (**3**) and cherylline respectively.

(Received, 14th July 1981; Com. 833.)

¹ Abstracted in part from M.Phil. dissertation of P. K., Panjab University, Chandigarh, India, 1979.

² A similar deleterious effect of 3,4-alkoxy-groups on a benzyne cyclization has been reported earlier; S. V. Kessar, D. Pal, and M. Singh, *Tetrahedron*, 1973, **29**, 177.

³ This is in contrast with the recent findings with *N*-acyl-*o*-chloroanilines; J. F. Wolfe, M. C. Sleevi, and R. R. Goehring, *J. Am. Chem. Soc.*, 1980, **102**, 3646.

⁴ For a recent synthesis of cherylline see D. J. Hart, P. A. Cain, and D. A. Evans, *J. Am. Chem. Soc.*, 1978, **100**, 1556.

⁵ Cf. R. A. Heacock, O. Hutzinger, B. D. Scott, J. W. Daly, and B. Witkop, *J. Am. Chem. Soc.*, 1963, **85**, 1825.