

Received April 5 1989 accepted July 6 1989

PRELIMINARY NOTE

---

A Facile Synthesis of  $\alpha$ -Trifluorodiazoo Compounds

GUOQIANG SHI and YUANYAO XU\*

Shanghai Institute of Organic Chemistry, Academia Sinica,  
345 Lingling Lu, Shanghai (China)

SUMMARY

A variety of  $\alpha$ -trifluorodiazoo compounds which can be used as building blocks for the construction of organic molecules bearing a trifluoromethyl group have been prepared in high yields simply by refluxing trifluoromethylketone 2,4,6-triisopropylbenzenesulphonylhydrazones with a methanolic solution of KOH. For characterization, their triphenylphosphazine derivatives were prepared.

Trifluoromethyl substitution imparts interesting properties to important organic materials such as bioactive compounds and ferroelectric liquid crystals [1, 2]. An attractive method for introducing a  $\text{CF}_3$  group into organic molecules is to use appropriate  $\text{CF}_3$  containing intermediates as synthetic building blocks [2], but the development of this methodology has been hampered by either the abnormal reactivity\* or the limited availability of trifluoromethylated building blocks. Therefore, the search for new versatile building blocks from easily accessible fluorinated precursors has been of considerable significance for solving the inherent synthetic problem.

---

\* For example, various  $\alpha$ -trifluorocarbanions of otherwise synthetic importance, enolates in particular, are marred by their proclivity towards defluorination [3].




As shown in Scheme 1, when the trisylhydrazones **1** were treated with 2 equivalents of KOH in methanol<sup>++</sup> under reflux, they rapidly decomposed within a few minutes to give the desired diazo compounds **2** in almost quantitative yield (assayed by <sup>19</sup>F NMR). The products obtained from alkaline work-up were essentially free from impurities as indicated by spectroscopic

\* Only very few  $\alpha$ -trifluorodiazoo compounds are known and their preparations resulted in low yields and impure products [5].

There has been a report that  $\alpha$ -trifluorodiazoo compounds could react normally via carbene intermediate [6].

+ Diazo compounds have been postulated as intermediates in the Bamford-Stevens reaction of aliphatic or alicyclic ketone hydrazones, but the only isolable products were usually compounds derived from their decompositions [4].

Although the Bamford-Stevens reaction can be performed under various conditions (see ref. 4), we have found the present condition to be the most suitable and convenient.



TABLE

Trifluorodiazoo compounds prepared from trisylhydrazones

Hydrazones <sup>a</sup> 1	Reaction <sup>b</sup> time(min)	Products <sup>d</sup> 2	Phosphazines <sup>f</sup>	
			Yield <sup>g</sup> , %	m p, °C
R=		R=		
CH <sub>3</sub> 1a	15 <sup>c</sup>	CH <sub>3</sub> 2a	73	135-136 <sup>h</sup>
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>3</sub> 1b	10	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>3</sub> 2b	81	41-42
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>5</sub> 1c	10	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>5</sub> 2c	70	46-48
PhCH <sub>2</sub> 1d	10	PhCH <sub>2</sub> 2d	78	94-95
Ph 1e	5	Ph 2e	76	152-154
c-C <sub>6</sub> H <sub>11</sub> 1f	10	c-C <sub>6</sub> H <sub>11</sub> 2f	80	163-164
PhCH=CH 1g	5	PhCH=CH <sup>e</sup> 2g	82	160-161

<sup>a</sup> Prepared by condensation of trifluoromethylketones [11] with an equal molar amount of trisylhydrazide in methanol containing a few drops of conc hydrochloric acid Yield, 75-93%

<sup>b</sup> Trisylhydrazone (2mmol) and KOH(ca 4mmol) in methanol (5ml) heated at refluxing temperature for a given time

<sup>c</sup> Reaction performed in a sealed tube at 60°C

<sup>d</sup> All products except 2g which was unstable under the reaction condition have been fully characterized spectroscopically (IR, <sup>19</sup>F NMR, <sup>1</sup>H NMR and MS) and the elemental composition of their phosphazine derivatives was determined by combustion analysis

<sup>e</sup> Spontaneously cyclized to form 4, m p 126-127°C

<sup>f</sup> For the preparation of phosphazines, see text


<sup>g</sup> Yields based on trisylhydrazones

<sup>h</sup> Lit [5a], 127-128 5°C

examination ( $^1\text{H}$  NMR,  $^{19}\text{F}$  NMR and IR) and could be readily trapped by triphenylphosphine to afford the corresponding phosphazines [9] in high yield. The results are summarized in the Table.

All diazo compounds listed in the Table generally have reasonable thermal stability with the doubly stabilized one, **2e**, being the most stable. Thus further purification for most of the diazo compounds, if necessary, is possible by short-path distillation under slightly reduced pressure using an aspirator. Like most other  $\alpha, \beta$ -unsaturated diazo compounds [10], compound **2g** lost its red color to form a pyrazole derivative **4** concomitant with the decomposition of the corresponding trisylhydrazone, however, it could be trapped *in situ* by  $\text{Ph}_3\text{P}$  to afford the corresponding phosphazine.

Interestingly, when our method was extended to the preparation of a polyfluorodiazo compound **6** from heptafluoropropyl butyl ketone trisylhydrazone **5**, an  $\alpha$ -defluorinated diazo compound **7** was obtained exclusively. This unanticipated result revealed a general feature of compounds such as **6**, 1e, the propensity towards defluorination, which can be rationalized by the process depicted in the following equation.



In summary, we have developed a convenient preparative method for a variety of  $\alpha$ -trifluorodiazo compounds. In view of many transformations of synthetic importance that can be carried out with diazo compounds, *inter alia*, those catalyzed by transition metals [12], the ready access to these  $\alpha$ -trifluorodiazo compounds may open a promising area for the synthesis of trifluoromethylated organic molecules. Work along this line is in progress.

In a typical experiment, a mixture of trifluoroacetophenone trisylhydrazone (2 mmol) and KOH (ca 4 mmol) in methanol (5 ml) was heated under reflux for 5 min. Water (10 ml) was added and then three extractions with ether were carried out. The extract was washed several times with saturated  $\text{NaHCO}_3$  solution and dried over  $\text{Na}_2\text{SO}_4$ . Removal of the solvent in a rotary evaporator gave a fairly pure orange product **2e**, IR(neat) 2090 ( $\text{C}=\text{N}=\text{N}$ ), 1140 ( $\text{CF}_3$ )  $\text{cm}^{-1}$ ,  $^{19}\text{F}$  NMR ( $\text{CCl}_4$ ,  $\delta$  TFA) -16.6 (downfield, s),  $^1\text{H}$  NMR ( $\text{CCl}_4$ ,  $\delta$ ) 6.97-7.40 (m), MS,  $m/z$  (relative intensity) 186 (18), 159 (100), 139 (22), 77 (11), 69 (20). For the preparation of phosphazine, a slight excess of  $\text{Ph}_3\text{P}$  was directly added to the ether extract. After the orange color had faded completely (ca 30 min), the solvent was evaporated and the residue was recrystallized from petroleum ether to give colorless crystals in 76% yield, mp 152-154°C.  $\text{C}_{26}\text{H}_{20}\text{F}_3\text{N}_2\text{P}$  calcd C 69.62, H 4.46, N 6.25%, found C 69.46, H 4.41, N 6.24%.

Financial support from the National Natural Science Foundation of China and from Academia Sinica is gratefully acknowledged.

- 1 (a) R. Filler and Y. Kobayashi, 'Biomedical Aspects of Fluorine Chemistry', Kodansha and Elsevier Biomedical, Tokyo (1983). (b) C. Walsh, *Tetrahedron*, **38** (1982) 871.
- 2 T. Fuchigami, *J. Syn. Org. Chem. Jpn.*, **42** (1984) 775 and reference cited therein.
- 3 T. Fuchigami and Y. Nakagawa, *J. Org. Chem.*, **52** (1987) 5276.
- 4 R. H. Shapiro, *Org. React.*, **23** (1976) 405.
- 5 (a) R. A. Shepard and D. L. Sciaraffa, *J. Org. Chem.*, **31** (1966) 964, (b) R. A. Shepard and S. E. Wentworth, *ibid.*, **32** (1967) 3197.
- 6 A. Ya. Aizikovitch, I. T. Bazyl, *Zh. Org. Khim.*, **23** (1987) 133.
- 7 A. R. Chamberlin and F. T. Bond, *J. Org. Chem.*, **43** (1978) 154.
- 8 D. M. Gale, N. J. Middleton and C. G. Krespan, *J. Am. Chem. Soc.*, **88** (1966) 3617.

- 9 G Singh and H Zimmer, Organometall Chem Rev , 2 (1967) 297
- 10 J L Brewbarker and H Hart, J Am Chem Soc , 91 (1969) 711
- 11 For their preparation, see A sykes, J C Fatlow and C R Thomas, J Chem Soc , (1956) 835
- 12 M P Doyle, Chem Rev , 86 (1986) 919