

TRIHYDROCARBYLTIN COMPLEXES OF TRANSITION METAL CYANIDES

R. USON, J. FORNIES, M.A. USON and E. LALINDE

Department of Inorganic Chemistry, University of Zaragoza (Spain)

(Received July 17th, 1979)

Summary

Cationic single-bridged trinuclear cyano complexes are obtained by treating trihydrocarbyltin perchlorate ($\text{SnR}_3\text{O}_2\text{ClO}_2$) (R = alkyl or aryl) with neutral transition metal monocyanides, whereas treatment with the corresponding neutral or anionic transition metal polycyanides gives neutral single-bridged polymeric cyano compounds. In every case the tin atoms are pentacoordinated with the three R groups in the equatorial and two CN groups (or, respectively one CN and one OClO_3 group) in the axial position of a trigonal bipyramid.

Introduction

The synthesis of covalent perchlorato derivatives from the corresponding halo complexes, according to eq. 1 and the subsequent substitution of the perchlorato



group, (which has a low coordinating capacity) by neutral (eq. 2) or anionic ligands (eq. 3) has frequently been used by us [1] and by others [2] for the



preparation of neutral and anionic organometallic complexes of gold [1a–1c], nickel [2a–2c], palladium [1d–1g, 2d], platinum [1h–1j] and rhodium [1k–1l].

If reaction 1 is carried out in a non-donor solvent under anhydrous conditions the intermediate perchlorato complex can sometimes be isolated, but is frequently unstable. Even so, after removal of the precipitated silver halide, the filtrate can in every case be used for processes 2 and 3.

Mainly because of the solubility of L_nMX it is, however, sometimes necessary to work in another type of solvent. In these cases it seems appropriate to employ

acetone, which, though it displaces the perchlorato group giving a (sometimes isolable) cationic perchlorate (eq. 4), can afterwards readily be displaced by




reactions of type 2 or 3.

We consider below the reaction of solutions of perchloratotrialkyltin or perchloratotriaryltin compounds with several cyano complexes of transition metals. These behave as a ligand L' (eq. 2) since their cyano group still has a potentially N-donor atom which causes the displacement of the perchlorato group or the acetone. The process allows the preparation of polymeric (finite or infinite) complexes with the cyano group bridging the tin and the transition metal atoms.

Some of these results were the subject of a preliminary communication [3].


Results and discussion

The precursor solutions used in the present study were prepared by treating the organochlorocomplexes SnR_3Cl (R being Me, Bu or Ph) with benzene or acetone solutions of $AgClO_4$, and removal of the $AgCl$. As reported [4] for the methyl complex $[SnMe_3O_2ClO_2]_x$ in the solid state the pentacoordinated tin atoms are bridged by the perchlorato ligands.




This bridging system is cleaved in methanol to give the pentacoordinated cationic perchlorate $[SnR_3S_2]ClO_4$ (S = solvent). Thus, whilst our benzene solutions are likely to contain $[Sn(O_2ClO_2)R_3]$, the acetone solutions are assumed to contain $[SnR_3(Me_2CO)_2]ClO_4$, but the reaction products with cyano-transition metal complexes are in each case the same.

(a) Reactions with monocyanides



The reaction (1/2) of benzene or acetone solutions of perchlorato trihydrocarbyltin $\text{Sn}(\text{O}_2\text{ClO}_2)\text{R}_3$ with cyanopentafluorophenylbis(triphenylphosphine)palladium(II) $[\text{Pd}(\text{CN})(\text{C}_6\text{F}_5)(\text{PPh}_3)_2]$ gives cationic trinuclear complexes (see eq. 5). The complexes I and III are precipitated instantaneously whatever the used molar ratio of the reagents used but the butyl complex (II) can only be obtained by use of a 2/1 mole ratio as in eq. 5. A 1/1 mole ratio leads to the




precipitation of the neutral complex IV, (see eq. 6), which contains a monodentate (C_{3v}) perchlorato group (see below).


The tendency of tin to adopt a coordination number of 5 means that the reaction of $\text{Sn}(\text{O}_2\text{ClO}_2)\text{R}_3$ with di- or poly-cyano-transition metal complexes leads to polymeric end products. Thus the reaction with neutral polycyano complexes yields cationic polymers, whereas anionic polycyanides give neutral polymers.

(b) Reactions with di- and poly-cyanides

The reaction with *trans*-dicyanobis(triphenylphosphine)palladium(II), according to eq. 7 gives rise to the immediate precipitation of the cationic



polynuclear complex V. The reactions with other cationic dicyano complexes, such as $(\text{Bu}_4\text{N})[\text{Ag}(\text{CN})_2]$ or $\text{K}[\text{Au}(\text{CN})_2]$ (which are very little soluble in benzene require the use of acetone ($\text{Me}_2\text{CO} = \text{S}$) as reaction medium) also give neutral complexes (eq. 8).



The complexes separate out upon mixing the two solutions. The simultaneously

precipitated QClO_4 can be removed by washing with water. No reaction could be observed for $\text{M} = \text{Ag}$ and $\text{R} = \text{Bu}$.

Complex V is decomposed by an acetone solution of NaBPh_4 finally giving $\text{trans-Pd}(\text{CN})_2(\text{PPh}_3)_2$. Treatment with tetra- and hexa-cyanometallates gives immediate precipitation of the corresponding neutral polymers.



(XI, $\text{M} = \text{Pd}$, $\text{R} = \text{Me}$;
 XII, $\text{M} = \text{Pd}$, $\text{R} = \text{Bu}$;
 XIII, $\text{M} = \text{Pd}$, $\text{R} = \text{Ph}$;
 XIV, $\text{M} = \text{Pt}$, $\text{R} = \text{Me}$;
 XV, $\text{M} = \text{Pt}$, $\text{R} = \text{Bu}$;
 XVI, $\text{M} = \text{Pt}$, $\text{R} = \text{Ph}$)

In the light of the square planar coordination of Pd or Pt , these polymers can be expected to have an infinite twodimensional structure (Fig. 1).

When $\text{K}_3[\text{Fe}(\text{CN})_6]$ is used no reaction occurs for $\text{R} = \text{Me}$, whereas when $\text{R} = \text{Bu}$ or Ph it gives neutral polymeric complexes (eq. 10), which are assumed to



have a three-dimensional structure.

(c) Solubility and conductivity

As expected for macromolecular substances, all the polymeric complexes (V–XVIII) are insoluble in water and organic solvents. The bi- and tri-nuclear complexes (I–IV) are soluble in chloroform, dichloromethane and nitromethane, and complexes II–IV are also soluble in acetone. All are insoluble in n-hexane




Fig. 1. Proposed structure of the $\{[\text{R}_3\text{Sn}]_2[(\mu\text{-CN})_4\text{M}]\}_x$ complexes. ($\text{M} = \text{Pd}$ or Pt ; $\text{R} = \text{Me}$, Bu , Ph).

and benzene. The conductivities of the soluble complexes show [5] that they behave as 1/1 electrolytes, in accordance with the proposed formulae.

(d) IR spectra

The discussion above requires that the reaction products contain only bridging CN groups while the starting materials contain only terminal CN groups. In accordance with this, the $\nu(\text{CN})$ stretching vibration of the resulting complexes is always shifted towards higher energy, and only a single band is observed [6]. Table 1 shows the values of $\Delta\nu$ ($\Delta\nu = \nu(\text{CN})_{\text{bridge}} - \nu(\text{CN})_{\text{terminal}}$) for all the new complexes.

All the pentafluorophenyl derivatives exhibit absorption at $\sim 1500\text{s}$, 1050s , 950s and 800s cm^{-1} due to the C_6F_5 ligand [7] along with those of the two mutually *trans*- PPh_3 groups [8] at 525s , 515s and 502s cm^{-1} (complexes I–IV).

All the methyl derivatives show a strong absorption at $560\text{--}555\text{ cm}^{-1}$ assignable to the $\nu_{\text{as}}(\text{Sn—C})$ stretching mode, and only complex XIV also shows a very weak band at 505 cm^{-1} assignable to $\nu_{\text{asym}}(\text{Sn—C})$. This is in good agreement with the proposed trigonal bipyramidal structure, with the three methyl groups in equatorial position and the two cyano groups at axial sites. This agrees with the structure proposed for other trimethyl derivatives of SnR_3X , with X being a bridging ligand (X = F [9]; Cl, Br [10]; ClO_4 [4,11]; BF_4 [12], AsF_6 , SbF_6 [13]; RCOO [14] or alkyl- and aryl-imidazole [15]).

The corresponding absorptions for the butyl derivatives located in the $650\text{--}600\text{ cm}^{-1}$ region, which are always weak [16] are in our complexes masked by bands due to the PPh_3 , ClO_4 or CN group. Our phenyl derivatives show two medium bands at $280\text{--}270$ and $240\text{--}230\text{ cm}^{-1}$, assignable to symmetric and asymmetric $\nu(\text{Sn—C})$ [17], showing that the Sn—C bonds are not coplanar.

Finally, all the ionic complexes (I, II, III and V) show absorptions at 1100s and 630m cm^{-1} assignable, respectively, to ν_3 or ν_4 vibrations of the perchlorato group (T_d). In complex IV the bands expected at 1100 and 630 cm^{-1} are split, and are located, respectively, at 1140 and 1040 cm^{-1} and at 615 and 625 cm^{-1} , corresponding to the ν_1 , ν_4 and ν_3 , ν_5 vibrations of the OClO_3 group (C_{3v}). This complex shows another band (A_1) at 915 cm^{-1} assignable to $\nu(\text{Cl—O})$, which would be IR active for C_{3v} symmetry, and this confirms the formulation proposed for this compound [18].

Experimental

C, H and N analyses were carried out with a Perkin–Elmer 240 microanalyzer. Melting points were measured on a Buchi apparatus (model Dr. Tottoli) and are uncorrected. Conductivities were determined in approximately $5 \times 10^{-4}\text{ M}$ solutions with a Philips PW 9501/01 conductimeter. IR spectra were recorded (over the range $4000\text{--}200\text{ cm}^{-1}$) on a Perkin–Elmer 599 spectrophotometer using Nujol mulls between polyethylene plates.

$\text{Pd}(\text{CN})(\text{C}_6\text{F}_5)(\text{PPh}_3)_2$ was prepared as described elsewhere [19]. $\text{Pd}(\text{CN})_2\text{--}(\text{PPh}_3)_2$ was obtained by treating a solution of $\text{PdI}_2(\text{PPh}_3)_2$ (9.01 g, 10 mmol) in 200 ml of acetone with freshly prepared AgCN (2.60 g, 20 mmol) for 60 min at room temperature with exclusion of light. The AgI was filtered off and

TABLE 1

ANALYTICAL AND OTHER DATA FOR POLYNUCLEAR COMPLEXES

Complex		Found (calcd.) %			Λ_M (ohm ⁻¹ cm ² mol ⁻¹)	$\nu(\text{CN})$ complex (cm ⁻¹)	$\Delta\nu$ (cm ⁻¹)	M.p. (°C)
		C	H	N				
I	$\{\text{Me}_3\text{Sn}[(\mu\text{-NC})_2[\text{Pd}(\text{C}_6\text{F}_5)(\text{PPh}_3)_2]_2]\text{ClO}_4$	54.75 (55.93)	3.76 (3.64)	1.45 (1.47)	115 ^b	2177	47	154(dec.)
II	$\{\text{Bu}_3\text{Sn}[(\mu\text{-NC})_2[\text{Pd}(\text{C}_6\text{F}_5)(\text{PPh}_3)_2]_2]\text{ClO}_4$	57.62 (57.77)	4.31 (4.30)	1.41 (1.37)	112 ^b	2155	25	195(dec.)
III	$\{\text{Ph}_3\text{Sn}[(\mu\text{-NC})_2[\text{Pd}(\text{C}_6\text{F}_5)(\text{PPh}_3)_2]_2]\text{ClO}_4$	58.59 (58.55)	3.60 (3.60)	1.35 (1.34)	82.9 ^a	2155	25	179(dec.)
IV	$[\text{Bu}_3(\text{O}_3\text{ClO})\text{Sn}[(\mu\text{-NC})\text{Pd}(\text{C}_6\text{F}_5)(\text{PPh}_3)_2]$	54.46 (54.44)	4.85 (4.73)	1.19 (1.15)	130 ^b	2155	25	203(dec.)
V	$\{\text{Ph}_3\text{Sn}[(\mu\text{-NC})\text{Pd}(\text{PPh}_3)_2](\mu\text{-CN})\}_x(\text{ClO}_4)_x$	59.11 (59.39)	4.04 (4.01)	2.59 (2.47)	^c	2150	30	192(dec.)
VI	$\{\text{Me}_3\text{Sn}[(\mu\text{-NC})\text{Ag}](\mu\text{-CN})\}_x$	17.91 (18.55)	2.59 (2.80)	8.58 (8.65)		2168	26	263
VII	$\{\text{Ph}_3\text{Sn}[(\mu\text{-NC})\text{Ag}](\mu\text{-CN})\}_x$	47.02 (47.11)	2.92 (2.96)	5.13 (5.49)		2160	20	>300
VIII	$\{\text{Me}_3\text{Sn}[(\mu\text{-NC})\text{Au}](\mu\text{-CN})\}_x$	14.55 (14.55)	2.14 (2.20)	6.74 (6.79)		2171	26	215(dec.)
IX	$\{\text{Bu}_3\text{Sn}[(\mu\text{-NC})\text{Au}](\mu\text{-CN})\}_x$	31.38 (31.19)	5.03 (5.05)	5.09 (5.20)		2170	25	265(dec.)
X	$\{\text{Ph}_3\text{Sn}[(\mu\text{-NC})\text{Au}](\mu\text{-CN})\}_x$	40.59 (40.10)	2.65 (2.52)	4.82 (4.68)		2170	25	220(dec.)
XI	$\{\text{Me}_3\text{Sn}[(\mu\text{-NC})_4\text{Pd}]\}_x$	22.36 (22.32)	3.25 (3.27)	10.24 (10.41)		2165	29	243(dec.)
XII	$\{\text{Bu}_3\text{Sn}[(\mu\text{-NC})_4\text{Pd}]\}_x$	42.30 (42.54)	6.69 (6.89)	6.96 (7.09)		2165	29	287(dec.)
XIII	$\{\text{Ph}_3\text{Sn}[(\mu\text{-NC})_4\text{Pd}]\}_x$	52.75 (52.77)	2.85 (3.32)	5.77 (6.15)		2165	29	210(dec.)
XIV	$\{\text{Me}_3\text{Sn}[(\mu\text{-NC})_4\text{Pt}]\}_x$	19.74 (19.26)	2.98 (2.89)	9.25 (8.94)		2165	32	>300
XV	$\{\text{Bu}_3\text{Sn}[(\mu\text{-NC})_4\text{Pt}]\}_x$	38.78 (38.25)	6.27 (6.19)	6.05 (6.37)		2160	27	>300
XVI	$\{\text{Ph}_3\text{Sn}[(\mu\text{-NC})_4\text{Pt}]\}_x$	48.62 (48.08)	3.21 (3.03)	4.76 (5.61)		2160	27	>300
XVII	$\{\text{Bu}_3\text{Sn}[(\mu\text{-NC})_6\text{Fe}]\}_x$	45.61 (46.22)	7.07 (7.55)	7.78 (7.77)		2146	28	182(dec.)
XVIII	$\{\text{Ph}_3\text{Sn}[(\mu\text{-NC})_6\text{Fe}]\}_x$	56.46 (57.11)	3.59 (3.59)	7.16 (6.66)		2140	22	163(dec.)

^a In nitromethane. ^b In acetone. ^c Insoluble.

the filtrate evaporated to dryness. The residue was recrystallized from dichloromethane/ethanol.

$K[Au(CN)_2]$, $(BuN_4)[Ag(CN)_2]$, $K_2[Pd(CN)_4]$ and $K_2[Pt(CN)_4]$ were prepared by standard procedures [20,21]. $K_3[Fe(CN)_6]$ was acquired from commercial sources.

(i) *Preparation of the organotin perchlorate complexes $[Sn(O_2ClO_2)R_3]$ ($R = Me, Bu, Ph$)*

Depending on their subsequent use these complexes were prepared either in benzene or acetone by treating stoichiometric amounts of the organotin complex SnR_3Cl with $AgClO_4$ for 30 min at room temperature. After removal of the $AgCl$ the filtrate was used for further reactions without additional treatment.

(ii) $\{[R_3Sn](\mu-CN)_2[Pd(C_6F_5)(PPh_3)_2]_2\}ClO_4$ (I, $R = Me$; II, $R = Bu$, III, $R = Ph$)

Addition of $Sn(O_2ClO_2)R_3$ (0.25 mmol) in 40 ml of benzene to a stirred solution of $Pd(CN)(C_6F_5)(PPh_3)_2$ (0.4120 g, 0.5 mmol) in 40 ml of benzene immediately gave a white microcrystalline precipitate which was filtered off and repeatedly washed with hexane (yields: Complex I 84%, complex II 75%, complex III (61%).

(iii) $\{[Bu_3(O_3ClO)Sn](\mu-NC)[Pd(C_6F_5)(PPh_3)_2]\}$ (IV)

Complex IV was similarly prepared but using a different molar ratio: $Pd(CN)(C_6F_5)(PPh_3)_2$ (0.4120 g, 0.5 mmol) in 20 ml of benzene and $Sn(O_2ClO_2)Bu_3$ (0.5 mmol) in 40 ml of benzene (84% yield).

(iv) $\{[Ph_3Sn](\mu-CN)[Pd(PPh_3)_2](\mu-CN)]_x(ClO_4)_x$ (V)

To a solution of $Sn(O_2ClO_2)Ph_3$ (0.5 mmol) in 25 ml of benzene was added $Pd(CN)_2(PPh_3)_2$ (0.3415 g, 0.5 mmol) in 40 ml of dichloromethane. The microcrystalline precipitate, which was difficult to filter off, was separated by centrifugation and washed with hexane (70% yield).

(v) $\{[R_3Sn](\mu-NC)[M(\mu-CN)]_x$ (VI, $M = Ag$, $R = Me$; VII, $M = Ag$, $R = Ph$; VIII, $M = Au$, $R = Me$; IX, $M = Au$, $R = Bu$; X, $M = Au$, $R = Ph$)

To a solution of $Q[M(CN)_2]$ ($M = Ag$, $Q = Bu_4N$; $M = Au$, $Q = K$) (1 mmol) in 25 ml of acetone was added an equimolecular solution of $Sn(O_2ClO_2)R_3$ in 25 ml of acetone. The immediately precipitated white (or in the case of VIII yellowish) solid was filtered off and repeatedly washed with water/acetone (yields: complex VI, 63%; VII, 76%; VIII, 66%; IX, 64%; X, 81%).

(vi) $\{[R_3Sn]_2[(\mu-NC)_4]_x$ (XI, $M = Pd$, $R = Me$; XII, $M = Pd$, $R = Bu$; XIII, $M = Pd$, $R = Ph$; XIV, $M = Pt$, $R = Me$; XV, $M = Pt$, $R = Bu$; XVI, $M = Pt$, $R = Ph$)

To a solution of $K_2[M(CN)_4]$ (1 mmol) in 10 ml of H_2O and 15 ml of acetone was added a solution of $Sn(O_2ClO_2)R_3$ (2 mmol) in acetone (1/2 mole ratio). Stirring was continued for 30–60 min. The white precipitate was filtered off and washed with benzene or acetone and then with hexane (yields: complex XI, 81%; XII, 88%; XIII, 85%; XIV, 91%; XV, 81%; XVI, 93%).

(vii) $\{[R_3Sn]_3[(\mu-NC)_6Fe]\}_x$ (XVII, $R = Bu$; XVIII, $R = Ph$)

The reaction was carried out as before by treating 1 mmol of the iron complex with 3 mmol of the organotin perchlorato complex. The resulting compounds were washed with acetone (yields; XVII, 95%; XVIII, 96%).

References

- (a) R. Usón, P. Royo, A. Laguna and J. Carcía, *Rev. Acad. Ciencias Zaragoza*, **28** (1973) 67; (b) R. Usón, P. Royo and A. Laguna, *Syn. Inorg. Metal-org. Chem.*, **3** (1973) 237; (c) R. Usón, A. Laguna and J.L. Sanjoaquin, *J. Organometal. Chem.*, **80** (1974) 147; (d) R. Usón, P. Royo and J. Forniés, *Syn. React. Inorg. Metal-Org. Chem.*, **4** (1974) 157; (e) R. Usón, J. Forniés and S. Gonzalo, *J. Organometal. Chem.*, **104** (1976) 253; (f) R. Usón, J. Forniés and F. Martínez, *J. Organometal. Chem.*, **112** (1976) 105; (g) R. Usón, J. Forniés, G. Gonzalo, F. Martínez and R. Navarro, *Rev. Acad. Ciencias Zaragoza*, **32** (1977) 85; (h) R. Usón, P. Royo and J. Gimeno, *J. Organometal. Chem.*, **72** (1974) 299; (i) R. Usón, J. Forniés, P. Espinet and M.P. García, *Rev. Acad. Ciencias Zaragoza*, **32** (1977) 85; (j) R. Usón, J. Forniés, P. Espinet and M.P. García, *Rev. Acad. Ciencias Zaragoza*, **32** (1977) 193; (k) R. Usón, L.A. Oro, C. Claver and M.A. Garraza, *J. Organometal. Chem.*, **105** (1976) 365; (l) R. Usón, P. Lahuerta, D. Carmona, L.A. Oro and K. Hildenbrand, *J. Organometal. Chem.*, **157** (1978) 63;
- (a) M. Wada and K. Oguro, *Inorg. Chem.*, **15** (1976) 2346; (b) M. Wada and F. Shimohigashi, *Inorg. Chem.*, **15** (1976) 954; (c) T. Majima and H. Kurosawa, *J. Organometal. Chem.*, **134**, (1977) C45; (d) J.M. Coronas, G. Muller and J. Sales, *Syn. React. Inorg. Metal-Org. Chem.*, **6** (1976) 217.
- R. Usón, J. Forniés, P. Espinet, R. Navarro and M.A. Usón, *Inorg. Chim. Acta*, **33** (1979) L103.
- H.C. Clark and R.J. O'Brien, *Inorg. Chem.*, **2** (1963) 740.
- W.J. Geary, *Coord. Chem. Rev.*, **7** (1971) 81.
- K. Nakamoto, *Infrared and Raman Spectra of Inorganic and Coordination Compounds*, Wiley Interscience, New York, 3rd ed., 1978, p. 266.
- D.A. Long and D. Steele, *Spectrochim. Acta*, **19** (1961) 1955.
- S.H. Mastin, *Inorg. Chem.*, **13** (1974) 1003.
- (a) H.C. Clark, R.J. O'Brien and J. Trotter, *Proc. Chem. Soc.*, (1963) 85; (b) H.C. Clark, R.J. O'Brien and J. Trotter, *J. Chem. Soc.*, (1964) 2332.
- H. Kriegsmann and S. Pischtschan, *Z. Anorg. Allg. Chem.*, **308** (1961) 212.
- R. Okawara, B.J. Hathaway and D.E. Webster, *Proc. Chem. Soc.*, (1963) 13.
- B.J. Hathaway and D.E. Webster, *Proc. Chem. Soc.*, (1963) 14.
- H.C. Clark and R.J. O'Brien, *Inorg. Chem.*, **2** (1963) 1020.
- (a) R. Okawara, D.E. Webster and E.G. Rochow, *J. Amer. Chem. Soc.*, **82** (1960) 3278; (b) R. Okawara and M. Ohara, *J. Organometal. Chem.*, **1** (1964) 360.
- R. Gassend, J.C. Maire and J.C. Pommier, *J. Organometal. Chem.*, **132** (1977) 69.
- J. Mendelshon, A. Marchand and J. Valade, *J. Organometal. Chem.*, **6** (1966) 25.
- R.C. Poller, *Spectrochim. Acta*, **22** (1966) 935.
- B.J. Hathaway and A.E. Underhill, *Chem. Soc.*, (1961) 3091.
- R. Usón, P. Royo and J. Forniés, *Rev. Acad. Ciencias, Zaragoza*, **28** (1973) 349.
- J.H. Bigelow, *Inorganic Synth.*, **2** (1946) 245.
- G. Brauer, *Química Inorgánica Preparativa*, Ed. Reverté, Barcelona, 1958, p. 630, 949.