This article was downloaded by: [University of Wyoming Libraries] On: 01 October 2013, At: 22:05 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



### Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/lsyc20

# Facile One-Pot Conversion of Aldehydes into Amides

Samuel T. Chill <sup>a</sup> & Robert C. Mebane <sup>a</sup>

<sup>a</sup> Department of Chemistry, University of Tennessee at Chattanooga, Chattanooga, Tennessee, USA Published online: 08 Jun 2010.

To cite this article: Samuel T. Chill & Robert C. Mebane (2010) Facile One-Pot Conversion of Aldehydes into Amides, Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry, 40:13, 2014-2017, DOI: <u>10.1080/00397910903219443</u>

To link to this article: <u>http://dx.doi.org/10.1080/00397910903219443</u>

### PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <a href="http://www.tandfonline.com/page/terms-and-conditions">http://www.tandfonline.com/page/terms-and-conditions</a>



Synthetic Communications<sup>®</sup>, 40: 2014–2017, 2010 Copyright © Taylor & Francis Group, LLC ISSN: 0039-7911 print/1532-2432 online DOI: 10.1080/00397910903219443

## FACILE ONE-POT CONVERSION OF ALDEHYDES INTO AMIDES

Samuel T. Chill and Robert C. Mebane

Department of Chemistry, University of Tennessee at Chattanooga, Chattanooga, Tennessee, USA

A facile one-pot synthesis of amides from aldehydes has been developed. This tandem process involves the formation of a nitrile intermediate obtained from the reaction of an aldehyde with hydroxylamine hydrochloride in dimethylsulfoxide at  $100^{\circ}$ C and the subsequent treatment of the nitrile with basic hydrogen peroxide. The resulting amide products were produced in good yields (67–95%) and purity (>95%).

Keywords: Aldehyde; amide; hydrogen peroxide; nitrile; tandem reaction

Nitriles are versatile synthetic intermediates because they can be transformed into a variety of other functional groups.<sup>[1,2]</sup> One such transformation is the selective hydrolysis of nitriles into amides.<sup>[3]</sup> In part because of their unique electronic and structural properties,<sup>[4]</sup> amides are particularly important functional groups and are found in a number of natural products, polymers, and pharmaceuticals. Thus, methods for preparing amides are highly desirable.

Recently,<sup>[5]</sup> we described a convenient one-pot synthesis of nitriles from aldehydes using hydroxylamine hydrochloride in dimethylsulfoxide (DMSO) at 100°C. We have now extended this synthetic methodology to the one-pot conversion of aldehydes into amides and report our findings herein.

The experimental procedure for the one-pot conversion of aldehydes to amides is simple and straightforward and affords amides in good isolated yields (Table 1). As an illustrative example, octanal (2.0 mmol) was added to a solution of hydroxylamine hydrochloride (3.8 mmol) in DMSO (4 mL), and the resulting reaction solution was stirred and heated for 30 min in a sand bath maintained at 100°C. Upon removal of the sand bath, but while still hot, aqueous sodium hydroxide (1 mL of 5.68 M, 5.68 mmol) was slowly added to the reaction mixture over a 1-min period with stirring, followed by the slow and careful addition of 30% hydrogen peroxide (2.5 mL) over a 5-min period. After stirring for 1 min, the reaction mixture was extracted with ether ( $4 \times 20$  mL), and the combined ether layers were washed with water ( $4 \times 20$  mL) and dried (MgSO<sub>4</sub>). Removal of the ether

Received April 13, 2009.

Address correspondence to Robert C. Mebane, Department of Chemistry, #2252, University of Tennessee at Chattanooga, 615 McCallie Ave., Chattanooga, TN 37403-2598, USA. E-mail: robert-mebane@utc.edu

### **ONE-POT CONVERSION OF ALDEHYDES INTO AMIDES**





Table 1. Facile one-pot synthesis of amides from aldehydes

(Continued)

Entry	Aldehyde	Amide	Yield (%) <sup>a</sup>	Time
12	H trans-3-phenyl-2- propenal	NH <sub>2</sub> trans-3-phenyl-2- propenamide	84	30
13	H I-naphthaldehyde	O NH <sub>2</sub> I-naphthamide	70	60

Table 1. Continued

<sup>a</sup>Isolated yields.

<sup>b</sup>Time for complete conversion to amide.

solvent by rotary evaporation and high vacuum gave octanamide as a white solid (0.231 g, 80% yield). The <sup>1</sup>H and <sup>13</sup>C NMR spectra and the mass spectra of the isolated octanamide were identical to literature spectra.<sup>[6]</sup>

Although no mechanistic studies were performed, it seems reasonable that this tandem conversion of an aldehyde to an amide involves two steps. First, the formation of an aldoxime followed by its dehydration gives a nitrile, and second, the nitrile is selectively hydrolyzed to an amide by the basic hydrogen peroxide, as fully described by Wiberg.<sup>[7]</sup>

As seen in Table 1, the reaction is general for the preparation of both aliphatic and aromatic amides. The amides prepared in this study were confirmed by comparison of <sup>1</sup>H and <sup>13</sup>C NMR spectra and mass spectra with authentic samples. Furthermore, the <sup>1</sup>H and <sup>13</sup>C NMR spectra of the amide products indicated that the product purities were greater than 95%.

In conclusion, we have demonstrated that aldehydes can be readily converted into amides by heating (100°C) the aldehydes in DMSO containing hydroxylamine hydrochloride, followed by selective hydrolysis of the subsequent nitrile with basic hydrogen peroxide. This one-pot, tandem process does not involve expensive or hazardous materials and should offer an attractive alternative to recently described procedures for converting aldehydes into amides, including the TerpyRu(PPh<sub>3</sub>)Cl<sub>2</sub>catalyzed rearrangement of aldoximes to amides,<sup>[8]</sup> the metal-free oxidative amination of aromatic aldehydes with *tert*-butyl hydroperoxide,<sup>[9]</sup> the treatment of aldehydes with iodine in aqueous ammonia followed by hydrogen peroxide,<sup>[10,11]</sup> the rhodium-catalyzed oxidative amination of aldehydes with N-methylmorpholine N-oxide,<sup>[12]</sup> and the treatment of aldehydes with hydroxylamine hydrochloride/ wet alumina/methanesulfonyl chloride.<sup>[13]</sup>

### ACKNOWLEDGMENTS

The authors gratefully acknowledge financial support from the Research Corporation, the University of Chattanooga Foundation, and the Grote Chemistry Fund.

### REFERENCES

- Friedrich, K.; Waaensfesl, K. The Chemistry of the Cyano Group; Wiley-Interscience: New York, 1970.
- Fatiadi, A. J. Preparation and Synthetic Applications of Cyano Compounds; Wiley: New York, 1983.
- 3. March, J. Advanced Organic Chemistry; McGraw-Hill: New York, 1977; p. 809.
- Bruice, P. Y. Organic Chemistry, 5th ed.; Pearson Prentice Hall: Upper Saddle River, NJ, 2007; p. 811.
- Chill, S. T.; Mebane, R. C. A facile one-pot conversion of aldehydes into nitriles. Synth. Commun. 2009, 39, 3601–3606.
- SDBSWeb (National Institute of Advanced Industrial Science and Technology). Available at http://riodb01.ibase.aist.go.jp/sdbs/.
- Wiberg, K. B. The mechanism of hydrogen peroxide reactions, I: The conversion of benzonitrile to benzamide. J. Am. Chem. Soc. 1953, 75, 3961–3964.
- Gnanamgari, D.; Crabtree, R. H. Terpyridine ruthenium-catalyzed one-pot conversion of aldehydes into amides. *Organomet. Chem.* 2009, 28, 922–924.
- Ekove-Kovi, K.; Wolf, C. Metal-free one-pot oxidative amination of aldehydes to amides. Org. Lett. 2007, 9, 3429–3432.
- Dunlap, N. K.; Drake, J.; Ward, A.; Salyard, T. L. J.; Martin, L. Concise synthesis of enantiomers of 4-aminobutane-1,2,3-triol. J. Org. Chem. 2008, 73, 2928–2930.
- Jiun-Jie, S.; Jim-Min, F. Direct conversion of aldehydes to amides, tetrazoles, and triazines in aqueous media by one-pot tandem reactions. J. Org. Chem. 2003, 68, 1158–1160.
- 12. Tillack, A.; Rudloff, I.; Beller, M. Catalytic amination of aldehydes to amides. *Eur. J. Org. Chem.* **2001**, 523–528.
- Sharghi, H.; Sarvari, M. H. A direct synthesis of nitriles and amides from aldehydes using dry or wet alumina in solvent-free conditions. *Tetrahedron* 2002, 58, 10323–10328.