Unexpected P-Si or P-C Bond Cleavage in the Reaction of $Li_2[(C_5Me_4)SiMe_2PR]$ (R = Cyclohexyl, 2,4,6-Me₃C₆H₂) and Li $[(C_5H_4)CMe_2PHR]$ (R = Ph, ^tBu) with ZrCl₄ or [TiCl₃(thf)₃]: Formation and Molecular Structure of the ansa-Metallocenes [$\{(\eta - C_5Me_4)_2SiMe_2\}$ ZrCl₂] and $[\{(\eta - C_5 H_4)_2 CMe_2\} MCl_2] (M = Ti, Zr)^{\dagger}$

Thomas Koch, Steffen Blaurock, Fernando B. Somoza, Jr., Andreas Voigt, Reinhard Kirmse, and Evamarie Hey-Hawkins*

Institut für Anorganische Chemie der Universität, D-04103 Leipzig, Germany

Received September 28, 1999

The (phosphinosilyl)cyclopentadiene 1-SiMe₂PHMes-2,3,4,5-Me₄C₅H, prepared from 1-SiMe₂- $Cl-2,3,4,5-Me_4C_5H$ and LiPHMes (Mes = 2,4,6-Me_3C_6H_2), reacts with MeLi to yield Li₂[(C₅- Me_4)SiMe₂PMes] (2). The (phosphinomethyl)cyclopentadienide Li[(C₅H₄)CMe₂PH^tBu] (4) is obtained by the reaction of LiPH^tBu with 6,6-dimethylfulvene. Li₂[(C_5Me_4)SiMe₂PR] (R = cyclohexyl (Cy; 1), Mes (2)) reacts in situ with $ZrCl_4$ at -30 °C to give [{(η -C₅Me₄)₂SiMe₂}- $ZrCl_2$] (5). The formation of (PCy)₄ and (PHMes)₂, respectively, in these reactions was shown by ³¹P NMR spectroscopy. In the reaction of Li[$(C_5H_4)CMe_2PHR$] (R = Ph (3), ^tBu (4)) with $ZrCl_4$ or $[TiCl_3(thf)_3]$, $[{(\eta - C_5H_4)CMe_2PHR}_2ZrCl_2]$ (R = Ph (6), ^tBu (7)) and $[{(\eta - C_5H_4)CMe_2-}$ PHR₂TiCl] (R = Ph (9), ^tBu (10)) are formed. 6, 7, 9, and 10 slowly decompose on recrystallization from thf, Et₂O, or toluene with formation of $[\{(\eta - C_5H_4)_2CMe_2\}MCl_2]$ (M = Zr (8), Ti (11)). The formation of $(PR)_n$ (R = Ph, n = 4-6; R = ^tBu, n = 4) and PH₂R was shown by ³¹P NMR spectroscopy. **2**, **4**, **5–8**, and **11** were characterized by NMR spectroscopy (¹H, ¹³C, ³¹P) and the paramagnetic Ti(III) complexes 9 and 10 by EPR spectroscopy and mass spectrometry. 5, $\hat{\mathbf{8}}$, and $1\hat{\mathbf{1}}$ were also characterized by crystal structure determinations.

Introduction

After the discovery of α -olefin polymerization by Ziegler,¹ a vast number of metallocene and monocyclopentadienyl derivatives with different steric and electronic properties was prepared and employed as catalysts in polymerization reactions.

ansa-Metallocene dichlorides of Ti, Zr, and Hf (especially chiral ones; A in Figure 1) are important catalysts for the methylalumoxane (MAO)-promoted isotactic polymerization of olefins. Therefore, intensive research is being conducted toward synthesis of improved catalysts.2-5

references therein.
(3) Kaminsky, W.; Rabe, O.; Schauwienold, A.-M.; Schupfner, G. U.;
Hanss, J.; Kopf, J. J. Organomet. Chem. 1995, 497, 181.
(4) Chacon, S. T.; Coughlin, E. B.; Henling, L. M.; Bercaw, J. E. J.
Organomet. Chem. 1995, 497, 171.
(5) (a) Kaminsky, W.; Külper, K.; Brintzinger, H. H.; Wild, F. R. W.
P. Angew. Chem. 1985, 97, 507; Angew. Chem., Int. Ed. Engl. 1985, 24, 507. (b) Brintzinger, H. H.; Fischer, D.; Mülhaupt, R.; Rieger, B.;
Waymouth, R. M. Angew. Chem. 1995, 107, 1255; Angew. Chem., Int. Ed. Engl. 1995, 34, 1143. (c) Fink, G.; Mülhaupt, R.; Brintzinger, H.
Ziegler Catalysts-Recent Scientific Innovations and Technological Improvements, Springer-Verlag: Berlin, Heidelberg, New York, 1995.



Figure 1. Selected examples of group 4 cyclopentadienyl complexes.

Recently, monocyclopentadienyl derivatives in which the cyclopentadienyl ring has a heteroatom-functionalized side chain (B in Figure 1), were prepared and employed in the polymerization of olefins and copolymerization of ethylene with propylene, 1-hexene, 1-octene or styrene.⁶ These studies were also extended to phospholyl ligands (C in Figure 1).⁷

 $^{^\}dagger$ Parts of this work were presented as a poster at the XVIII International Conference on Organometallic Chemistry, Munich, Germany, Aug 16-22, 1998; Abstract No. A223.

⁽¹⁾ Ziegler, K.; Holzkamp, E.; Breil, H.; Martin, H. Angew. Chem. 1955, 67, 541.

⁽²⁾ Schneider, N.; Huttenloch, M. E.; Stehling, U.; Kirsten, R.; Schaper, F.; Brintzinger, H. H. Organometallics 1997, 16, 3413 and references therein.

^{(6) (}a) Shapiro, P. J.; Bunel, E.; Schaefer, W. P.; Bercaw, J. E. Organometallics **1990**, *9*, 867. (b) Shapiro, P. J.; Cotter, W. D.; Schaefer, W. P.; Labinger, J. A.; Bercaw, J. E. J. Am. Chem. Soc. **1994**, *116*, 4623. (c) Flores, J. C.; Chien, J. C. W.; Rausch, M. D. Organometallics **1994**, *13*, 4140. (d) Alt, H. G.; Föttinger, K.; Milius, W. J. Organomet. Chem. **1999**, *572*, 21. (e) Galan-Fereres, M.; Koch, T.; Hey-Hawkins, E.; Eisen, M. J. Organomet. Chem. **1999**, *580*, 145. (f) McKnight, A. L.; Waymouth, R. M. *Chem. Rev.* **1998**, *380*, 145. (1) McKnight, A. L.; Waymouth, R. M. *Chem. Rev.* **1998**, *98*, 2587. (g) Canich, J. A. M. Eur. Pat. 420 436, 1991. (h) Schmidt, G. F.; Timmers, F. J.; Knight, G. W.; Lai, S.-Y.; Nickias, P. N.; Rosen, R. K.; Stevens, J. C.; Wilson, D. R. Eur. Pat. 416 815, 1991.

While numerous O- and N-functionalized cyclopentadienyl ligands are known, the number of cyclopentadienes with a P-functionalized alkyl or silyl side chain is still small, and we have recently reported on the first compounds in which the P atom has a reactive phosphorus-substituent bond (e.g., P-H).8,9 Such compounds should be useful precursors for dianionic bifunctional ligands in transition-metal chemistry. In addition, the presence of a P-containing group is advantageous for reactivity studies by ³¹P NMR spectroscopy.¹⁰

We now report unexpected P-Si or P-C bond cleavage in the reactions of $Li_2[(C_5Me_4)SiMe_2PR]$ (R = cyclohexyl (Cy; 1),⁸ 2,4,6-Me₃C₆H₂ (Mes, 2)) with ZrCl₄ and of $Li[(C_5H_4)CMe_2PHR]$ (R = Ph (3),⁸ ^tBu (4)) with ZrCl₄ or [TiCl₃(thf)₃], which give the *ansa*-metallocenes $[\{(\eta - C_5Me_4)_2SiMe_2\}ZrCl_2]$ (5) and $[\{(\eta - C_5H_4)_2CMe_2\}$ - MCl_2 (M = Zr (8), Ti (11)). 5, 8, and 11 were characterized by spectroscopy and by crystal structure determination. While the ansa-metallocene 5 has been reported before,¹¹ its structure was unknown up to now.

Results and Discussion

Synthesis. The general synthetic method for introduction of a cyclopentadienyl ligand into Zr or Ti complexes is salt elimination. We therefore prepared the (phosphanidosilyl)cyclopentadienides Li₂[(C₅Me₄)SiMe₂-PR] (R = Cy (1),⁸ Mes (2)) from 1-SiMe₂PHR-2,3,4,5- Me_4C_5H and MeLi at -50 °C in thf. The neutral (phosphinosilyl)cyclopentadienes were obtained from (dimethylchlorosilyl)tetramethylcyclopentadiene and LiPHR at -80 °C followed by distillation. The (phosphinomethyl)cyclopentadienides Li[(C₅H₄)CMe₂PHR] $(R = Ph (3),^{8} ^{t}Bu (4))$ were obtained by the reaction of LiPHR with 6,6-dimethylfulvene in Et₂O.

After treating **1** and **2** with ZrCl₄ (ratio 1:1), we were, however, unable to isolate the expected cyclopentadienyl derivatives $[{(\eta - C_5 Me_4)SiMe_2 PR}MCl_2];$ instead, the ansa-metallocene [{ $(\eta$ -C₅Me₄)₂SiMe₂}ZrCl₂] (5) was obtained after recrystallization from thf in good yield. The formation of $(PCy)_4^{12,13}$ or $(PHMes)_2^{14}$ in this reaction (eq 1) was shown by ³¹P NMR spectroscopy. The Si compound which must have been formed could not be characterized by ²⁹Si NMR spectroscopy. These results are contrary to what was originally described by Dow Chemical Co. in a patent application. There, the reaction between the related dilithio salt Li₂[(C₅Me₄)SiMe₂PPh] and TiCl₄ or ZrCl₄ was reported to give the expected products $[{(\eta - C_5Me_4)SiMe_2PPh}MCl_2]$, but no analytical or spectroscopic data were reported to support this assumption.^{6h} 5 was previously prepared from [ZrCl₄-



 $(thf)_2$ and the dimetalated ligand in low yield (15%);¹¹ this is a general preparative method for ansa-metallocenes of Ti, Zr, and Hf.^{2,3,4,11}

In the reaction of **3** or **4** with $ZrCl_4$ (Scheme 1) or [TiCl₃(thf)₃] (Scheme 2), the formation of [$\{(\eta - C_5H_4)CMe_2 PHR_{2}ZrCl_{2}$ (R = Ph (6), ^tBu (7)) as pale yellow solids and of $[{(\eta - C_5H_4)CMe_2PHR}_2TiCl]$ (R = Ph (9), ^tBu (10)) as dark-green solids was observed. However, these compounds decompose slowly on recrystallization from thf, Et₂O, or toluene with formation of $[{(\eta - C_5H_4)_2}]$ CMe_2 }MCl₂] (M = Zr (8), Ti (11)). The Ti compound that is presumably formed in addition to 11 could not be isolated. The decomposition is not influenced by light. The formation of cyclooligophosphanes $(PR)_n^{13}$ and PH₂R in this reaction was shown by ³¹P NMR spectroscopy.

For 6, the decomposition to 8 was studied in more detail. 6 was suspended in d_8 -toluene (in which it is only slightly soluble), and d₈-thf was added until a clear solution had formed. The decomposition was studied by ¹H NMR spectroscopy. After 3 days at room temperature, the ratio of complexes 6 and 8 was 5:1, while at 50 °C, the ratio was 7:1 after 4 h. Complete formation of 8 from 6 had occurred after 8 days at room temperature. As the decomposition is spontaneous in methanol, 6 was dissolved in methanol, and the decomposition products were analyzed by GC-MS. Besides products which stem from a reaction with methanol, signals for phenylphosphine (m/z 110, high intensity), PhP=CMe₂ (m/z 150, medium intensity), and $(PPh)_4$ (m/z 429, m/z 429)medium intensity) were observed. While this experiment shows that neither propene nor polypropylene is formed, the mechanism of the formation of 5, 8, and 11 is still speculative.

The Ti(III) complexes 9 and 10 are highly airsensitive. Brief exposure of toluene solutions to minute traces of oxygen apparently also triggers the quantitative formation of 11.

A stable zirconocene derivative with [(diarylphosphino)isopropyl]cyclopentadienyl ligands, namely, [$\{(\eta C_5H_4$)CMe₂PR₂ $_2$ ZrCl₂] (R = *p*-tolyl, Ph), has been reported.15

The complexes 6, 7, 9, and 10 were characterized by IR and ³¹P NMR spectroscopy (the signals are only slightly broadened for 9 and 10), 6 and 7 were also

^{(7) (}a) Brown, S. J.; Gao, X.; Harrison, D. G.; Koch, L.; Spence, R. E. V. H.; Yap, G. P. A. Organometallics 1998, 17, 5445. (b) Mathey, F. Chem. Rev. 1998, 88, 429. (c) Janiak, C.; Lange, K. C. H.; Versteeg, U.; Lentz, D.; Budzelaar, P. Chem. Ber. 1996, 129, 1517. (d) Janiak, C.; Versteeg, U.; Lange, K. C. H.; Weimann, R.; Hahn, E. J. Organomet. Chem. **1995**, 501, 219.

⁽⁸⁾ Koch, T.; Hey-Hawkins, E. Polyhedron 1999, 18, 2113 and references therein.

⁽⁹⁾ Slone, C. S.; Weinberger, D. A.; Mirkin, C. A. Prog. Inorg. Chem. 1999. 48. 233.

⁽¹⁰⁾ Berger, S.; Braun, S.; Kalinowski, H. D. NMR Spectroscopy, Thieme Verlag: Stuttgart, Germany, 1993; Vol. 3 (³¹P NMR Spectroscopy)

⁽¹¹⁾ Jutzi, P.; Dickbreder, R. Chem. Ber. 1986, 119, 1750.

⁽¹²⁾ Bart, J. C. J. Acta Crystallogr. 1969, 25B, 762.

 ⁽¹³⁾ Baudler, M.; Glinka, K. Chem. Rev. 1993, 93, 1623.
 (14) Kurz, S.; Oesen, H.; Sieler, J.; Hey-Hawkins, E. Phosphorus,

Sulfur Silicon Relat. Elem. 1996, 117, 189.

⁽¹⁵⁾ Bosch, B. E.; Erker, G.; Fröhlich, R.; Meyer, O. Organometallics 1997. 16. 5449.





studied by ¹H NMR spectroscopy. The paramagnetic complexes **9** and **10** were also characterized by mass spectrometry and studied by EPR spectroscopy (vide infra). **6**, **7**, **9**, and **10** exhibit a doublet in the ³¹P NMR



Figure 2. Experimental (a) and simulated (b) X-band EPR spectra of **10** in toluene at 295 K. The low-field part is additionally increased in intensity by a factor of 5.



Figure 3. Experimental (a) and simulated (b) X-band EPR spectra of **10** in toluene at 130 K.

spectrum (in C₆D₆), which is shifted to low field relative to **3** and **4** (**6**, 17.1 ppm (${}^{1}J_{P-H} = 206.0 \text{ Hz}$); **7**, 42.3 ppm (${}^{1}J_{P-H} = 193.8 \text{ Hz}$); **9**, 18.4 ppm (d, ${}^{1}J_{P-H} = 212.0 \text{ Hz}$); **10**, 38.6 ppm (d, ${}^{1}J_{P-H} = 195.0 \text{ Hz}$); **3**, -1.20 ppm (d of two q, ${}^{1}J_{P-H} = 207.5 \text{ Hz}$, ${}^{3}J_{P-H} = 11.4 \text{ Hz}$ and ${}^{3}J_{P-H} =$ 15.8 Hz);⁸ **4**, -26.0 ppm (d of two dec, ${}^{1}J_{P-H} = 189.9 \text{ Hz}$, ${}^{3}J_{P-H} = 11.0 \text{ Hz}$).

For **6** and **7**, molecular ion peaks were observed in the mass spectra, confirming their composition.

EPR Spectra of $[{(\eta - C_5H_4)CMe_2PHR}_2TiCl]$ (R = Ph (9), ^tBu (10)). For both Ti(III) complexes 9 and 10 intense EPR signals were observed. The room-temperature EPR spectrum of **10** is shown in Figure 2 together with its simulation. It consists of an intense doublet due to the interaction of the unpaired electron with one ³¹P nucleus (nuclear spin I = 1/2). Each of the doublet lines is symmetrically flanked by a hyperfine sextet resulting from the ⁴⁷Ti nucleus ($I = \frac{5}{2}$, natural abundance 7.4%) and a hyperfine octet caused by the 49 Ti nucleus (I = $^{7/_{2}}$, natural abundance 5.4%). Because of the similarity of the nuclear properties of the two Ti isotopes,¹⁶ for each "31P hyperfine line" an octet with the intensity ratio 0.7:1.9:1.9:1.9:1.9:1.9:0.7 was expected and observed. However, the intense "31P doublet" prevents the observation of the "inner" lines (Figure 2) of the ^{47,49}Ti hyperfine pattern.

Figure 3 shows the EPR spectrum of **10** in a frozen solution (T = 130 K) together with its simulation. The spectrum is rhombic-symmetric, like that recorded for

⁽¹⁶⁾ BRUKER Almanac; Bruker Analytic GmbH: Rheinstetten/ Karlsruhe, Germany, 1999.

Table 1. EPR Parameters^a for $[{(\eta - C_5H_4)CMe_2PHPh}_2TiCl]$ (9)^b and $[{(\eta - C_5H_4)CMe_2PH'Bu}_2TiCl] (10)^c$

tensor component	9	10	tensor component	9	10
g_1	2.006	2.007	A_1^{P}	14.5	14.8
g_2	1.990	1.992	$A_2^{\rm P}$	15.2	15.4
g_3	1.975	1.975	$A_3^{\tilde{P}}$	18.4	18.9
$g_{\mathrm{av}}{}^d$	1.990	1.991	$A_{av}^{P} d$	16.0	16.4
g_0	1.989	1.990	a_0^{P}	16.3	16.9
a_0^{Ti}	9.8	9.6	solvent	thf	toluene

^{*a*} Hyperfine coupling constants are given in 10^{-4} cm⁻¹. ^{*b*} $g_i \pm 0.004$; $A_i^{\text{Ti},\text{P}} \pm 1.0$. ^{*c*} $g_i \pm 0.002$; $A_i^{\text{Ti},\text{P}} \pm 0.5$. ^{*d*} $g_{av} = (g_1 + g_2 + g_3)/3$; $A_{av}^{\text{P}} = (A_1^{\text{P}} + A_2^{\text{P}} + A_3^{\text{P}})/3$.

9. All spectra can be described with the spin Hamiltonian (2), with $S = \frac{1}{2}$; all other symbols have their usual meaning.

$$\mathbf{H}_{sp} = \mathbf{g} \cdot \boldsymbol{\mu}_{B} \cdot \mathbf{S} \cdot \mathbf{B}_{0} + \mathbf{S} \cdot \mathbf{A}^{P} \cdot \mathbf{I}^{P} + \mathbf{S} \cdot \mathbf{A}^{Ti} \cdot \mathbf{I}^{Ti}$$
(2)

In Table 1 the principal values obtained for the g tensor and the ³¹P and ^{47,49}Ti hyperfine structure (=hfs) tensors \mathbf{A}^{P} and \mathbf{A}^{Ti} are listed. Anisotropic ^{47,49}Ti hfs couplings were observed in the frozen-solution spectra, but unfortunately they could not be analyzed in detail because (i) the line widths of the frozen-solution spectra are 2.5 times larger than those observed in liquid solution, (ii) the intensity of the Ti satellite lines is low (<2% of the major signals), and (iii) the small **g** tensor anisotropy results in excessive line overlap, which prevents an unambiguous analysis of the observed ^{47,49}Ti satellites. In addition, ^{47,49}Ti hfs coupling parameters so far observed¹⁷ are small and lie only in the range of $(10-20) \times 10^{-4}$ cm⁻¹. In our case (compound 10) a maximum Ti hfs coupling constant of about 18 imes 10^{-4} cm⁻¹ can be estimated, which we ascribe to A_z^{Ti} .

The hfs parameters can be used to estimate the spin density on the nuclei and, therefore, allow conclusions to be made about the nature of the M-L bonds, in particular the hybridization of the orbitals.^{18,19} For this, the complete parameters of the hfs tensors are needed. The spin densities $(c_{s,p,d}^X)^2$ can then be obtained by using eqs 3a,b, where s, p, and d represent atomic

$$(c_{\rm s}^{\rm X})^2 = a_{\rm av,exptl}^{\rm X}/a_{\rm av,theor}^{\rm X} \tag{3a}$$

$$(c_{\rm p,d}^{\rm X})^2 = b_{\rm exptl}^{\rm X}/b_{\rm theor}^{\rm X}$$
 (3b)

orbitals, X is the nucleus under discussion, a_{av}^X is the isotropic hyperfine coupling constant, and b^X is the dipolar part of the hyperfine tensors. For the ³¹P nucleus the complete tensor has been obtained. Using the values for the parameters $a_{av,theor}^{X}$ and b_{theor}^{X} calculated by Morton and Preston¹⁹ and eqs 3a,b gave the following spin densities for the ³¹P 3s and 3p orbitals: $(c_s^P)^2 =$ 0.004 and $(c_p^P)^2 = 0.010$. The latter values can be used to estimate the 3s/3p hybridization of the P atom (eq

4), where $\alpha_{\rm P}^2$ is the overall spin density on the P atom and amounts to $\alpha_{\rm P}^2 = 1.4\%$.

$$(c_{\rm s}^{\rm P})^2 = \alpha_{\rm P}^2 (1 - n^2) \qquad (c_{\rm p}^{\rm P})^2 = \alpha_{\rm P}^2 n^2$$
 (4)

For the degree of hybridization n^2 a value of 0.71 was obtained, which is close to that expected for a distorted tetrahedrally coordinated P atom. This corresponds to the structure proposed for the Ti(III) complex under study.

To our knowledge, up to now, ³¹P hfs interactions were only observed for the Ti(III) complex [Cp₂TiMe(PPh₃)].²⁰ The observed isotropic ³¹P hfs coupling constant of $a_0^P = 17.6 \times 10^{-4} \text{ cm}^{-1}$ is close to those obtained by us. Unfortunately, no anisotropic ³¹P hfs data were reported,²⁰ so that no conclusions can be drawn about the overall spin density and the degree of hybridization n^2 on the P atom.

Similar properties were observed by Xin et al. for $[(Cp_2Ti)_2(\mu-H)(\mu-PCy_2)]$.²¹ This Ti(III) complex exhibits no EPR signal at 253 K; this indicates the presence of diamagnetic species. At room temperature, the EPR spectrum shows an intense singlet ($g_0 = 1.9808$, $a_0^{\text{Ti}} =$ 9.43 G). No ³¹P hfs interaction was observed.

Because of the lack of a complete parameter set for the ^{47,49}Ti hyperfine tensor **A**^{Ti}, only a rough estimate of the spin density on the metal atom can be made. From the isotropic Ti hfs coupling and by using eq 3a and $a_{av,theor}^{Ti} = 260.8 \times 10^{-4} \text{ cm}^{-1}$,¹⁹ a Ti 4s contribution of 3.7% can be calculated. To estimate the Ti 3d contribution, we assume the observed maximum Ti hfs coupling constant of about 18×10^{-4} cm⁻¹ as being the axial component A_z^{Ti} of the Ti hfs tensor. Then, from $A_z^{\text{Ti}} - a_0^{\text{Ti}} = 2b$, a *b* value of 4.2×10^{-4} cm⁻¹ was obtained. With $b_{\text{theor}}^X = 7 \times 10^{-4}$ cm⁻¹, ¹⁹ eq 3 yields a Ti 3d contribution of about 60%. Therefore, roughly 65% of the unpaired spin density is located on the metal. Since the spin density determined for the coordinating P atom is small, the residual spin density is expected to be located on the Cl and Cp ligands.

Molecular Structures of $[{(\eta - C_5Me_4)_2SiMe_2}]$ - $ZrCl_2$] (5) and [{(η -C₅H₄)₂CMe₂}MCl₂] (M = Zr (8), **Ti** (11)). Compound 5 was obtained as pale green crystals from thf and crystallizes in the monoclinic space group C2/c (Figure 4). Complex 8 was obtained as yellow crystals on recrystallizing 6 or 7 from thf after several days. Accordingly, 11 is obtained on recrystallizing 9 or 10 from thf after several days as dark red crystals. 8 (Figure 5) and **11** (Figure 6) are isotypic and isostructural and crystallize in the monoclinic space group $P2_1/m$.

In **5**, the Zr atom and the Si atom of the SiMe₂ bridge are located on a crystallographic C_2 axis which generates the atoms "a". In 8 and 11, the metal atom (Zr or Ti) as well as Cl(1), Cl(2), and the C atoms of the CMe_2 bridge (C(1), C(2), C(3)) are located on a crystallographic mirror plane, by which the atoms "a" are generated. A comparison of 5, 8, and 11 with related ansa-metallocenes is given in Tables 2 and 3, respectively. The Zr-Cl and Si–C bond lengths of **5** are in the range observed for related complexes (Table 2). The Zr-CEN distance

⁽¹⁷⁾ Bajgur, C. S.; Tikkanen, W. R.; Petersen, J. L. Inorg. Chem. 1985, 24, 2539.

⁽¹⁸⁾ Kirmse, R.; Stach, J. *ESR-Spektroskopie-Anwendungen in der Chemie*, Akademie-Verlag: Berlin, 1985; pp 86–95.
(19) Morton, J. R.; Preston, K. F. *J. Magn. Reson.* 1978, *30*, 577.

⁽²⁰⁾ Klei, E.; Teuben, J. H. *J. Organomet. Chem.* **1980**, *188*, 97. (21) Xin, S.; Woo, H. G.; Harrod, J. F.; Samuel, E.; Lebuis, A. J. Am. Chem. Soc. 1997, 119, 5307.



Figure 4. Molecular structure of $[\{(\eta-C_5Me_4)_2SiMe_2\}ZrCl_2]$ (5) showing the atom-numbering scheme employed (ORTEP plot, 50% probability, SHELXTL PLUS; XP³²). Hydrogen atoms are omitted for clarity.



Figure 5. Molecular structure of $[\{(\eta-C_5H_4)_2CMe_2\}ZrCl_2]$ (**8**) showing the atom-numbering scheme employed (ORTEP plot, 50% probability, SHELXTL PLUS; XP³²). Hydrogen atoms are omitted for clarity.

is about 0.1 Å larger. As no data for the $CMe_2-C(Cp)$ bond lengths are given in the literature, the bond lengths of C(1)-C(4) and C(1)-C(4A) of 1.532(3) Å in **8** and **11** cannot be compared with known values (Table 3). Of the reported complexes, **8** has the smallest CEN-Zr-CEN bite angle (116.7°).

Experimental Section

All experiments were carried out under purified dry argon. Solvents were dried and freshly distilled under argon. The NMR spectra were recorded at 25 °C in C_6D_6 with an AVANCE DRX 400 spectrometer (Bruker), ¹H NMR: internal standard,



Figure 6. Molecular structure of $[\{(\eta-C_5H_4)_2CMe_2\}TiCl_2]$ (**11**) showing the atom-numbering scheme employed (ORTEP plot, 50% probability, SHELXTL PLUS; XP³²). Hydrogen atoms are omitted for clarity.

solvent (benzene); external standard, TMS. ¹³C NMR: external standard, TMS; internal standard, solvent. ³¹P NMR: external standard, 85% H₃PO₄. The IR spectra were recorded on a FT-IR spectrometer Perkin-Elmer System 2000 in the range 350–4000 cm⁻¹. GC-MS: Hewlett-Packard HP 6890 GC, HP 5973 MSD (mass selective detector). EPR spectra were recorded in the X-band ($\nu \approx 9.5$ GHz) with a Bruker ESP 300E spectrometer in the temperature range 295 K $\geq T \geq 130$ K by using about 10⁻³ M solutions of **9** (thf) and **10** (toluene). The EPR parameters were obtained by using the computer program Win-EPR-SimFonia.²⁸ The melting points were determined in sealed capillaries under argon and are uncorrected. Li₂[(C₅-Me₄)SiMe₂PCy] **(1**).⁸ Li[(C₅H₄)CMe₂PHPh] **(3**).⁸ [TiCl₃(thf)₃].²⁹ and 1-SiMe₂Cl-2,3,4,5-Me₄C₅H³⁰ were prepared according to the literature procedures. ZrCl₄ is commercially available.

Synthesis of 1-SiMe₂PHMes-2,3,4,5-Me₄C₅H (2a). LiPH-Mes (3.50 g, 22.2 mmol) was suspended in a mixture of *n*-hexane (50 mL) and Et₂O (5 mL) and cooled to -80 °C. (Dimethylchlorosilyl)tetramethylcyclopentadiene (4.83 g, 22.5 mmol) was added, and the reaction mixture was warmed to ambient temperature. The mixture was stirred for 3 days, after which the yellow color of the lithium phosphanide had disappeared completely. The mixture was filtered, the solvent evaporated in vacuo, and the resulting yellow oil distilled under reduced pressure (100–115 °C, 2·10⁻³ Torr), giving 2.2 g (30%) of **2a** as a yellow oil. ¹H NMR (C₆D₆): δ 6.76 (s, 2H, *m*-H in Mes), 3.30 (d, 1H, PH, ¹J_{P-H} = 211.4 Hz), 2.86 (s, 1H, C₅HMe₄), 2.33 (s, 6H, *o*-CH₃ in Mes), 2.10 (s, 3H, *p*-CH₃ in

- (24) Razavi, A.; Atwood, J. L. *J. Organomet. Chem.* 1995, 497, 105.
 (25) Willoughby, C. A.; Davis, W. M.; Buchwald, S. L. *J. Organomet. Chem.* 1995, 497, 11.
- (26) Smith, J. A.; von Seyerl, J.; Huttner, G.; Brintzinger, H. H. J. Organomet. Chem. **1979**, *173*, 175.
- (27) Nifant'ev, I. E.; Churakov, A. V.; Urazowski, I. F.; Mkoyan, Sh. G.; Atovmyan, L. O. *J. Organomet. Chem.* **1992**, *435*, 37.
- (28) Weber, R. T. WIN-EPR-SimFonia, Version 1.2; EPR Division, Bruker Instruments Inc., 1995.
 - (29) Manzer, L. F. Inorg. Synth. 1982, 21, 135.
 - (30) Plenio, H. Chem. Ber. 1991, 124, 2185.

⁽²²⁾ Herrmann, W. A.; Rohrmann, J.; Herdtweck, E.; Spaleck, W.; Winter, A. Angew. Chem. **1989**, *101*, 1536; Angew. Chem., Int. Ed. Engl. **1989**, *28*, 1511.

⁽²³⁾ Spaleck, W.; Antberg, M.; Rohrmann, J.; Winter, A.; Bachmann, B.; Kiprof, P.; Behm, J.; Herrmann, W. A. Angew. Chem. 1992, 104, 1371; Angew. Chem., Int. Ed. Engl. 1992, 31, 1345.
(24) Razavi, A.; Atwood, J. L. J. Organomet. Chem. 1995, 497, 105.

 Table 2. Comparison of Selected Bond Lengths (Å) and Angles (deg) of 5 with Those of Related

 Silylene-Bridged Zirconocene Dichloride Complexes

					CEN-Zr-	C(Cp)-Si-	
compd	Zr-C	Si-C(Cp)	Zr-CEN	Cl-Zr-Cl	CEN^{a}	C(Cp)	ref
rac-[{(2-methyltetrahydrobenz[e]indenyl) ₂ - SiMe ₂ }ZrCl ₂]	2.4100(7)	1.876(3)	2.238	97.35(4)	128.1	95.6(2)	2
<i>rac</i> -[({bis(4,5,6,7-tetrahydroindenyl)}SiMe ₂)ZrCl ₂]	2.444, 2.440	b	2.226, 2.225	97.3	126.3	b	3
[({bis(2,4,7-trimethylindenyl)}SiMe ₂)ZrCl ₂]	2.414, 2.414	b	2.236, 2.236	97.2	129.5	b	3
rac -[{(2-SiMe ₃ -4-Bu ^t -C ₅ H ₂) ₂ SiMe ₂ }ZrCl ₂]	2.435(1), 2.422(1)	1.872(3), 1.879(3)	2.233, 2.233	97.6(3)	126.7	94.0(1)	4
$[\{(\eta-C_5H_4)_2SiMe_2\}ZrCl_2]$	2.435(1)	1.866(4)	2.197	97.98(4)	125.4	93.2(2)	17
rac-[({bis(indenyl)}SiMe ₂)ZrCl ₂]	2.431(1)	1.871(1)	2.293	98.76(1)	119.0	94.57(6)	22
$\label{eq:rac-left} \begin{array}{l} rac\mbox{-}[(\{bis(2\mbox{-}methylindenyl)SiMe_2\})ZrCl_2] \\ [\{(\eta\mbox{-}C_5Me_4)_2SiMe_2\}ZrCl_2] \mbox{(5)} \end{array}$	2.419 2.4334(7)	b 1.887(2)	2.244 2.329	99.06 99.28(3)	128.1 128.6	94.3(1) 95.7(1)	23 this work

^{*a*} CEN = center of the cyclopentadienyl ring. ^{*b*} No value given.

 Table 3. Comparison of Selected Bond Lengths (Å) and Angles (deg) of 8 and 11 with Those of Related

 Methylene-Bridged Titanocene and Zirconocene Dichloride (and Dimethyl) Complexes

compd	M-Cl	C _{br} -C(Cp)	M-CEN	Cl-M-Cl	CEN-M- CEN ^a	C(Cp)-C- C(Cp)	ref
[Me ₂ C{(3-Bu ^t -C ₅ H ₃)-9-fluorenyl}ZrCl ₂]	2.420, 2.420	b	2.184, 2.245	96.2	118.5	b	3
$[Me_2C{(3-Me-C_5H_3)-9-fluorenyl}ZrCl_2]$	2.425(3), 2.403(3)	b	2.16, 2.26	99.1	118.6	98.4	24
$[cyclo-C_6H_{10}(\eta-C_5H_4)(4,5,6,7-tetrahydro-indenvib))$ T;Cl	2.346(2), 2.338(2)	b	2.07, 2.08	97.6(1)	121.8	96.3(4)	25
$[\{(\eta-C_5H_4)_2CH_2\}$ TiCl ₂]	2.33(1), 2.34(1)	1.53(5)	2.05, 206	97.2(3)	121	99(2)	26
$[\{(\eta - C_5H_4)_2CMe_2\}TiMe_2]$		1.465(6)	b		120.9	b	27
$[\{(\eta - C_5H_4)_2CMe_2\}ZrMe_2]$		1.441(6)	b		115.9	b	27
$[{(\eta - C_5H_4)_2CMe_2}ZrCl_2]$ (8)	2.4321(8), 2.4406(8)	1.532(3)	2.213	100.25(3)	116.7	99.5(2)	this work
$[{(\eta - C_5H_4)_2CMe_2}TiCl_2]$ (11)	2.333(1), 2.352(1)	1.532(3)	2.193	98.04(4)	121.6	96.6(2)	this work

^{*a*} CEN = center of the cyclopentadienyl ring. ^{*b*} No value given.

Mes), 1.97 (s, 6H, CH₃ in C₅Me₄), 1.84 (s, 6H, CH₃ in C₅Me₄), 0.17 (d, 6H, SiMe₂, ${}^{3}J_{P-H} = 5.0$ Hz). ${}^{31}P$ NMR (C₆D₆): δ –168.6 (d, ${}^{1}J_{P-H} = 212.0$ Hz). EI-MS: m/z 328 (25) [M⁺], 179 (100) [(CH₃)₄C₅Si(CH₃)₂⁺], 151 (85) [PHMes⁺], and fragmentation thereof. Anal. Calcd for C₂₀H₃₁SiP (328.51): C, 72.7; H, 9.4; P, 9.4. Found: C, 67.3 (lower due to silicon carbide formation); H, 9.0; P, 9.2.

Synthesis of Li₂[(C₅Me₄)SiMe₂PMes] (2). 2a (2.2 g, 6.7 mmol) was dissolved in thf (50 mL), and at -50 °C an MeLi solution (8.9 mL of a 1.5 M solution in Et₂O) was added slowly. The mixture was then brought to ambient temperature and refluxed for 5 h. The compound was used without further purification. ³¹P NMR (thf/ C₆D₆): δ –181.2 br, s.

Synthesis of Li[(C₅H₄)CMe₂PH^tBu] (4). 6,6-Dimethylfulvene (1.34 g, 12.6 mmol) was added to a suspension of LiPH^tBu (1.21 g, 12.6 mmol) in Et₂O (30 mL), and the reaction mixture was stirred for 12 h. The resulting solid was isolated by filtration and washed twice with *n*-hexane to yield **4** as a pale yellow powder (1.5 g, 58%). Mp: 120 °C dec. ¹H NMR (thf- d_8): δ 6.06 (s, 2H, CH in C₅H₄), 6.00 (s, 2H, CH in C₅H₄), 3.57 (d, 1H, P-H, ${}^{1}J_{P-H} = 199.2$ Hz), 1.82 (d, 3H, C(CH₃)₂, ${}^{3}J_{P-H} =$ 14.2 Hz), 1.80 (d, 3H, C(CH₃)₂, ${}^{3}J_{P-H} = 14.3$ Hz), 1.16 (d, 9H, $C(CH_3)_3$, ${}^3J_{P-H} = 10.6$ Hz). ${}^{13}C$ NMR (thf- d_8): δ 132.23 (d, *ipso-*C in C_5H_4 , ${}^2J_{P-C} = 8.5$ Hz), 103.39 (C3, C4 in C_5H_4), 102.46 (d, C2, C5 in C₅H₄, ${}^{3}J_{P-C} = 3.3$ Hz), 39.53 (d, C-*C*H₃, ${}^{2}J_{P-C} =$ 10.3 Hz), 35.33 (d, $C(CH_3)_2$, ${}^1J_{P-C} = 15.6$ Hz), 34.08 (d, $C(CH_3)_3$, ${}^{1}J_{P-C} = 15.1$ Hz), 33.52 (d, C-*C*H₃, ${}^{2}J_{P-C} = 11.7$ Hz), 31.89 (d, $C(CH_3)_3$, ${}^2J_{P-C} = 12.7$ Hz). ${}^{31}P$ NMR (thf, C_6D_6): δ 26.0 (d of two dec, ${}^{1}J_{P-H} = 189.9$ Hz, ${}^{3}J_{P-H} = 11.0$ Hz). ⁷Li NMR (thf*d*₈): δ –4.6, s. IR (KBr): ν (cm⁻¹): 2282 m (ν (PH)), 1606 br, m (Cp, v(C=C)). Anal. Calcd for C₁₂H₂₀PLi (202.20): C, 71.3; H, 10.0; P, 15.3. Found: C, 70.9; H, 9.8; P, 15.9. Recrystallization from Et₂O gave colorless crystals.

Synthesis of ansa-{Dimethylsilylbis(tetramethylcyclopentadienyl)}zirconium Dichloride (5). (a) From ZrCl₄ and Li₂[(C₅Me₄)SiMe₂PCy] (1). A 0.76 g (3.3 mmol) amount of ZrCl₄ was suspended in 30 mL of toluene, and a solution of 1.00 g (3.3 mmol) of Li₂[(C₅Me₄)SiMe₂PCy] (1) in 25 mL of thf was added dropwise at -30 °C. The solution turned dark red immediately. The reaction mixture was stirred at room tem-

perature for 12 h. Then the solvent was removed in vacuo and the residue washed with 20 mL of pentane. The resulting yellow powder was recrystallized from thf, yielding 0.72 g of **5** (82% relative to **1**) as pale green crystals. The product $(PCy)_{4}^{13}$ (³¹P NMR (C₆D₆): δ –68.4 s) was detected by ¹H and ³¹P NMR spectroscopy.

(b) From ZrCl₄ and Li₂[(C_5Me_4)SiMe₂PMes] (2). Alternatively, **5** was obtained from ZrCl₄ (1.00 g, 4.4 mmol) in 30 mL of toluene and Li₂[(C_5Me_4)SiMe₂PMes] (2; 1.50 g, 4.4 mmol) in 25 mL of thf at -30 °C. Yield: 1.41 g (70% relative to 2). The product (PHMes)₂¹⁴ (³¹P NMR (C_6D_6): δ -111.1, *meso* isomer; -118.5, *d* and *l* isomer) was detected by ¹H and ³¹P NMR spectroscopy.

¹H and ¹³C NMR (in CDCl₃) are in agreement with the literature values.¹¹ In C_6D_6 , the signals are shifted slightly: ¹H NMR δ 2.07 (s, 12H, C_5Me_4), 1.79 (s, 12H, C_5Me_4), 0.61 (s, 6H, Si Me_2).

Synthesis of ansa-{Dimethylmethanylbis(cyclopentadienyl)}zirconium Dichloride (8). (a) From ZrCl4 and Li-[(C₅H₄)CMe₂PHPh] (3). A 1.30 g (5.6 mmol) amount of ZrCl₄ was suspended in 50 mL of toluene, and a solution of 1.24 g (5.6 mmol) of Li[(C₅H₄)CMe₂PHPh] (3) in 20 mL of thf was added dropwise at -50 °C. The solution turned dark red immediately. The reaction mixture was stirred at room temperature for 12 h. Then the solvent was removed in vacuo and the residue washed with 50 mL of pentane. The resulting pale yellow powder exhibits a doublet in the ³¹P NMR spectrum (C_6D_6) at 17.1 ppm $(^1J_{P-H} = 206.0 \text{ Hz})$, which is consistent with the presence of the complex $[{(\eta - C_5H_4)CMe_2PHPh}_2ZrCl_2]$ (6). On recrystallization from thf, 6 decomposed over several days with formation of 8. Yield: 1.17 g of 8 (61% relative to 3) as yellow crystals. The products $(PPh)_n^{13}$ (n = 4-6) and PH_2Ph were detected by ³¹P NMR spectroscopy (C₆D₆: δ -4.0 (m, (PPh)₅), -26.4 (s, (PPh)₆), -47.7 (s, (PPh)₄), -123.8 (t, PH₂-Ph, ${}^{1}J_{P-H} = 197$ Hz)).

(b) From ZrCl₄ and Li[(C_5H_4)CMe₂PH^tBu] (4). Alternatively, **8** was obtained from ZrCl₄ (1.10 g, 4.7 mmol) in 50 mL of toluene and Li[(C_5H_4)CMe₂PH^tBu] (4; 0.95 g, 4.7 mmol) in 20 mL of Et₂O at -50 °C. The solution turned dark red immediately. The reaction mixture was stirred at room tem-

perature for 24 h. Then the solvent was removed in vacuo and the residue washed with 25 mL of pentane. The resulting pale yellow powder exhibits a doublet in the ³¹P NMR spectrum (C₆D₆) at 42.3 ppm (¹J_{P-H} = 193.8 Hz), consistent with the presence of the complex [{(η -C₅H₄)CMe₂PH^tBu}₂ZrCl₂] (7). On recrystallization from thf, 7 slowly decomposed over several days with formation of **8**. Yield: 1.2 g of **8** (75% relative to **4**) as yellow crystals. The products (P^tBu)₄¹³ and PH₂^tBu were detected by ³¹P NMR spectroscopy (C₆D₆: δ –32.4 (s, ?), –59.0 (s, (P^tBu)₄), –80.1 (t, PH₂^tBu, ¹J_{P-H} = 174 Hz)).

[{(η-C₅H₄)**CMe₂PHPh**}**2Zr**Cl₂] (6). ¹H NMR (thf-*d*₈): δ 7.36 (m, 2H, Ph), 7.28 (m, 2H, Ph), 7.10–6.93 (m, 6H, Ph), 6.03 (m, 1H, C*H* in C₅H₄), 5.99 (t, 1H, C*H* in C₅H₄, ³*J*_{H-H} = 2.7 Hz), 5.95 (t, 1H, C*H* in C₅H₄, ³*J*_{H-H} = 2.7 Hz), 5.88 (t, 1H, C*H* in C₅H₄, ³*J*_{H-H} = 2.7 Hz), 5.85 (m, 1H, C*H* in C₅H₄, ³*J*_{H-H} = 3.5 Hz), 5.79 (m, 1H, C*H* in C₅H₄), 5.77 (m, 1H, C*H* in C₅H₄), 5.73 (m, 1H, C*H* in C₅H₄), 4.00 (d, 2H, P–H, ¹*J*_{P-H} = 207.3 Hz), 1.11 (d, 6H, CH₃, ³*J*_{H-H} = 9.8 Hz), 1.10 (d, 6H, CH₃, ³*J*_{H-H} = 9.8 Hz). ³¹P NMR (C₆D₆): δ 17.1 (d, ¹*J*_{P-H} = 206.0 Hz). IR (KBr): ν (cm⁻¹): 2292 m (ν(PH)).

[{ $(\eta$ -C₅H₄)**CMe₂PH'Bu**}₂**ZrCl**₂] (7). ¹H NMR (C₆D₆): δ 6.26 (t, 4H, CH in C₅H₄, ³J_{H-H} = 2.8 Hz), 6.01 (m, 4H, CH in C₅H₄), 3.57 (d, 2H, P-H, ¹J_{P-H} = 189.0 Hz), 1.75 (m, 6H, CH₃ in CMe₂) 1.71 (m, 6H, CH₃ in CMe₂), 0.86 (d, 18H, CH₃ in ^tBu, ³J_{P-H} = 11.4 Hz). ³¹P NMR (C₆D₆): δ 42.3 (d, ¹J_{P-H} = 193.8 Hz). IR (KBr): ν (cm⁻¹): 2379 m (ν (PH)).

[{ $(\eta$ -C₅H₄)₂CMe₂}ZrCl₂] (8). ¹H NMR (C₆D₆): δ 6.38 (t, 4H, CH in C₅H₄, ³J_{H-H} = 2.7 Hz), 5.15 (t, 4H, CH in C₅H₄, ³J_{H-H} = 2.7 Hz), 1.07 (s, 6H, CH₃). ¹³C NMR (C₆D₆): δ 123.23 (s, 4C, C₅H₄), 123.05 (s, 2C, *ipso*-C in C₅H₄), 107.17 (s, 4C, C₅H₄), 37.92 (s, 1C, C-CH₃), 24.10 (s, 2C, CH₃). EI-MS: 332 (100) [M⁺], 296 (25) [M⁺ - Cl], 281 (25) [M⁺ - Cl - Me]. Anal. Calcd for C₁₃H₁₄ZrCl₂ (332.38): C, 47.0; H, 4.2. Found: C, 46.7; H, 4.1.

Synthesis of ansa-{Dimethylmethanylbis(cyclopentadienyl) { titanium Dichloride (11). (a) From [TiCl₃(thf)₃] and Li[(C₅H₄)CMe₂PHPh] (3). A 1.25 g (3.4 mmol) amount of [TiCl₃(thf)₃] was dissolved in 25 mL of thf, and a solution of 0.75 g (3.4 mmol) of Li[(C₅H₄)CMe₂PHPh] (3) in 30 mL of thf was added at -20 °C. The solution turned dark green immediately. The reaction mixture was stirred at room temperature for 12 h. Then the solvent was removed in vacuo and the residue washed with 20 mL of hexane. The resulting green powder (complex 9) was recrystallized from thf. The EPR spectrum of this highly air-sensitive solution showed the presence of a paramagnetic Ti(III) species ([$\{(\eta - C_5H_4)CMe_2-$ PHPh₂TiCl] (9)) which exhibits Ti-P interaction. The ³¹P NMR spectrum (C₆D₆) showed a doublet at 18.4 ppm (${}^{1}J_{P-H} =$ 212.0 Hz). The dark green solution decomposes slowly over several days with formation of a red solution from which dark red crystals of 11 can be obtained. Yield: 0.54 g (55% relative to **3**). The products $(PPh)_n^{13}$ (n = 4-6) and PH_2Ph were detected by ³¹P NMR spectroscopy (C₆D₆: δ -4.0 (m, (PPh)₅), -26.4 (s, (PPh)₆), -47.7 (s, (PPh)₄), -123.8 (t, PH₂Ph, ${}^{1}J_{P-H} =$ 197 Hz)).

(b) From [TiCl₃(thf)₃] and Li](C₅H₄)CMe₂PH'Bu] (4). Alternatively, **11** was obtained from 1.00 g (2.7 mmol) of [TiCl₃-(thf)₃], dissolved in 25 mL of thf and a solution of 0.60 g (2.7 mmol) of Li](C₅H₄)CMe₂PH'Bu] (4) in 30 mL of thf at -20 °C. The solution turned dark green immediately. The reaction mixture was stirred at room temperature for 12 h. Then the solvent was removed in vacuo and the residue washed with 20 mL of hexane. The resulting green powder (complex **10**) was recrystallized from toluene. An EPR spectrum of this highly air-sensitive solution showed the presence of a paramagnetic Ti(III) species ([{ $(\eta$ -C₅H₄)CMe₂PH'Bu}₂TiCl] (**10**)) which exhibits Ti-P interaction. The dark green solution decomposes slowly with formation of a red solution, from which dark red crystals of **11** can be obtained. Yield: 0.48 g (62% relative to **4**). The products (P'Bu)₄¹³ and PH₂'Bu were detected by ³¹P

Table 4. Crystal Data and Structure RefinementDetails for 5, 8, and 11

	5	8	11
formula	C ₂₀ H ₃₀ Cl ₂ SiZr	C ₁₃ H ₁₄ Cl ₂ Zr	C ₁₃ H ₁₄ Cl ₂ Ti
$M_{\rm r}$	460.65	332.38	289.04
temp (K)	220(2)	213(2)	220(2)
cryst syst	monoclinic	monoclinic	monoclinic
space group	C2/c (No. 15)	$P2_1/m$ (No. 11)	$P2_1/m$ (No. 11)
a (Å)	15.454(3)	7.118(1)	7.102(2)
b (Å)	12.219(2)	8.831(1)	8.604(4)
<i>c</i> (Å)	11.135(2)	10.163(1)	9.967(3)
β (deg)	105.05(3)	90.49(1)	90.67(2)
$V(Å^3)$	2030.5(6)	638.84(3)	609.0(3)
Z	2	2	2
ρ_{calcd} (Mg m ⁻³)	1.507	1.790	1.576
F(000)	952	344	296
cryst size (mm)	$0.5\times0.4\times0.3$	$0.5\times0.4\times0.3$	$0.4\times0.2\times0.1$
abs coeff (mm^{-1})	0.864	1.251	1.108
$2\theta_{\rm max}$ (deg)	4.3 - 56.1	4.0 - 55.5	4.1 - 54.7
no. of rflns	10942	3565	2666
collected			
no. of indep rflns	2306	1466	1313
$R_{\rm int}$	0.0286	0.0264	0.0173
no. of params	170	112	112
$R(I \geq 2\sigma(I))$	0.0247	0.0227	0.0282
wR2 (all data)	0.0626	0.0609	0.0836
$(\Delta/\rho)_{\rm min}$ (e Å ⁻³)	-0.374	-0.396	-0.328
$(\Delta/ ho)_{ m max}$ (e Å ⁻³)	0.330	0.367	0.436

NMR spectroscopy (C₆D₆: δ -32.4 (s, ?), -59.0 (s, (P^tBu)₄), -80.1 (t, PH₂^tBu, ¹J_{P-H} = 174 Hz)).

[{(η-C₅H₄)CMe₂PHPh}₂TiCl] (9). ³¹P NMR (C₆D₆): δ 18.4 (d, ¹J_{P-H} = 212.0 Hz). EI-MS: *m*/*z* 513 (7) [M⁺], 297 (20) [M⁺ - C₅H₄Me₂PHPh], 260 (75) [M⁺ - 2 Ph], 216 (25) [C₅H₄-CMe₂PHPh⁺], and fragmentation thereof. IR (KBr): ν (cm⁻¹): 2287 m (ν (PH)), 1579 m (ν (C=C)), 1569 m (ν (C=C)). EPR: see Table 1.

[{ $(\eta$ -C₅H₄)CMe₂PH^tBu}₂TiCl] (10). ³¹P NMR (C₆D₆): δ 38.6 (d, ¹*J*_{P-H} = 195.0 Hz). EI-MS: *m/z* 473 (2) [M⁺], 385 (10) [M⁺ - P^tBu], 296 (12) [M⁺ - 2 P^tBu], 260 (12) [(C₅H₄)₂CMe₂-Ti⁺], and fragmentation thereof. IR (KBr): ν (cm⁻¹): 2282 m (ν (PH)), 1471 s (ν (C=C)), 1261 s (ν (C=C)). EPR: see Table 1.

[{ $(\eta$ -C₅H₄)₂CMe₂}TiCl₂] (11). ¹H NMR (CDCl₃): δ 6.90 (t, 4H, CH in C₅H₄, ³J_{H-H} = 2.5 Hz), 5.55 (t, 4H, CH in C₅H₄, ³J_{H-H} = 2.5 Hz), 1.74 (s, 6H, CH₃). ¹³C NMR (CDCl₃): δ 132.08 (s, 4C, C₅H₄), 116.42 (s, 2C, *ipso*-C in C₅H₄), 111.86 (s, 4C, C₅H₄), 37.77 (s, 1C, C-CH₃), 24.11 (s, 2C, C-CH₃). EI-MS: *m*/*z* 288 (28) [M⁺], 252 (30) [M⁺ - Cl], 237 (10) [M⁺ - Cl -Me]. Anal. Calcd for C₁₃H₁₄TiCl₂ (289.04): C, 54.0; H, 4.8. Found: C, 53.8; H, 4.7.

X-ray Crystal Structure Determination of 5, 8, and 11. Data (λ (Mo K α) = 0.710 73 Å) were collected with a Siemens CCD (SMART) diffractometer. All observed reflections were used for refinement (SAINT) of the unit cell parameters. Empirical absorption correction was carried out with SAD-ABS.³¹ The structures were solved by direct methods (SHELX-TL PLUS).³² Zr, Ti, Si, Cl, and C atoms were refined anisotropically; H atoms were located by difference maps and refined isotropically. Table 4 lists crystallographic details. Crystallographic data (excluding structure factors) for the structures reported in this paper have been deposited with the Cambridge Crystallographic Data Centre (5, CCDC-133344; 8, CCDC-133345; 11, CCDC-133346). Copies of the data can be obtained free of charge on application to The Director, CCDC, 12 Union Road, Cambridge, CB2 1EZ, U.K. (fax, int. code +(1223)336-033; e-mail, deposit@ccdc.cam.ac.uk).

⁽³¹⁾ Sheldrick, G. M. SADABS—A Program for Empirical Absorption Correction; University of Göttingen, Göttingen, Germany, 1998.

⁽³²⁾ SHELXTL PLUS, Siemens Analytical X-ray Instruments Inc., 1990: XS, Program for Crystal Structure Solution; XL, Program for Crystal Structure Determination; XP, Interactive Molecular Graphics.

Unexpected P-Si or P-C Bond Cleavage

Acknowledgment. We gratefully acknowledge financial support from the *Fonds der chemischen Industrie*, the *Alexander von Humboldt-Foundation* (F.B.S.), the *Landesgraduiertenförderung Sachsen* (S.B.) and the *German-Israeli Foundation* (Grant No. I-142/95).

Supporting Information Available: Tables giving X-ray crystal structure data for **5**, **8**, and **11**. This material is available free of charge via the Internet at http://pubs.acs.org.

OM990763Z