

AN EFFICIENT SYNTHESIS OF 5-ARYL(or ALKYL)AMINO-4-ETHOXYCARBONYL-2-METHYLTHIO-1,3-THIAZOLES FROM DIMETHYL N-(ETHOXYCARBONYLMETHYL)IMINODITHIOCARBONATE AND ISOTHIOCYANATES.

Carlos Alvarez-Ibarra*, Matilde Gil, Paloma Ortiz, and Maria L. Quiroga.

Departamento de Química Orgánica I. Facultad de Química. Universidad Complutense.


28040 Madrid. España.

Abstract- An efficient synthesis of perfunctionalized 1,3-thiazoles 2a-2f with good yields has been carried out by a cyclocondensation reaction between the α -metallated dimethyl N-(ethoxycarbonylmethyl)iminodithiocarbonate and aryl or alkylisothiocyanates.

In this paper we give account of an efficient synthesis of 5-aryl(or alkyl)amino-4-ethoxycarbonyl-2-methylthio-1,3-thiazoles 2a-2f. These perfunctionalized 1,3-thiazoles can be used as building blocks to obtain other heterocyclic structural complex systems by a convergent connection with specific synthons. Thus, the methylthio on C2 can be substituted by halogen, O-, N-, and C-nucleophiles by or without a previous oxidation of the sulfide.¹ This strategy can be used, for instance, to obtain the famotidine, an interesting antiulcer agent.² By other hand, the ethoxycarbonyl on C4 can be the starting point for a new heterocyclization³ to azadiaryl compounds that can be used as target molecules to obtain compounds of biological interest.⁴ Additionally, the presence of the amine substituent on C5 can make possible the fusion of other hetero- or homocycles of variable topology with 1,3-thiazole moiety.

We have previously reported that the dimethyl N-(ethoxycarbonylmethyl)iminodithiocarbonate (EMIC), 1, is an useful synthetic equivalent of the $\text{EtO}_2\text{C}-\text{C}_4-\text{N}=\text{C}_2-\text{SMe}$ synthon that is a structural unit present into 1,3-thiazoles whose substituents on C5 position come from an unsaturated electrophile.⁵ Thus, 5-alkylthio-4-ethoxycarbonyl-2-methylthio-1,3-thiazoles have been obtained by a cyclocondensation reaction of EMIC with carbon disulfide, and later alkylation of intermediate 1,3-thiazole-5-thiolate with alkyl halides.^{5,6}

We describe now an efficient synthesis of 5-aryl(or alkyl)amino-4-ethoxycarbonyl-2-methylthio-1,3-thiazoles, 2a-2f, by a cyclocondensation reaction between the α -metallated EMIC and aryl and alkyl isothiocyanates (Scheme 1).



R: a, Ph; b, *p*-FC₆H₄; c, *p*-MeC₆H₄; d, *p*-MeOC₆H₄; e, Et; f, *n*-Bu

-Scheme 1-

Different base/solvent systems, base/EMIC proportions, and reaction conditions for metallation of 1, and condensation of α -metallated EMIC with phenyl isothiocyanate were tested. The results have been gathered in Table 1.

Table 1. Observed yields for 4-ethoxycarbonyl-2-methylthio-5-phenylamino-1,3-thiazole, 2a, from 1 and phenyl isothiocyanate.

Run	Base/solvent	Base/EMIC/PhNCS	reaction conditions		yield(%) ^a
			metallation	condensation	
1	KBu ^t O/THF	1.1/1/1	20°C/0.5 h	20°C/0.5 h	36
2	KBu ^t O/THF	1.1/1/1	-78°C/0.5 h	-78°C/0.5 h 20°C/2 h	49
3	KBu ^t O/THF	1.4/1/1	-78°C/0.5 h	-78°C/0.5 h 20°C/2 h	90
4	NaH/DMSO	1.5/1/1		20°C/2 h	b)
5	NaH/DME	1.5/1/1	0°C/0.5 h	20°C/0.5 h	c)
6	NaOH ^d	---/1/1		20°C/2 h	19

^aIsolated product. Except in the indicated cases none other product was detected by TLC except 1.

^bBis(4-ethoxycarbonyl-2-methylthio-1-phenyl imidazolyl)disulfane was obtained (10%) by acidification to pH 5 of the reaction crude after of the extraction with diethyl ether at basic pH with negative result.⁷ ^cMethyl N-phenyldithiocarbamate was obtained with a yield of 45%. ^dThe following conditions were used: CH₂Cl₂/NaOH-H₂O (4%)/*n*-Bu₄N⁺, Br⁻.

In all cases, except runs 4 and 5, the 1,3-thiazole 2a was the unique isolated product. The best result in 2a was obtained with the KBu^tO/THF system (Table 1, run 3). From run 4 a solid with a melting point of 200°C was isolated and identified as bis[5-(4-ethoxycarbonyl-2-methylthio-1-phenyl)imidazolyl]disulfane by according with ir, ¹H- and ¹³C-nmr data.⁷ A related result has been described in the literature⁸ for the cyclocondensation reaction of N-tosylmethylisocyanide and isothiocyanates with the NaH/DMSO system.

From the results gathered in Table 1, the synthesis of 5-aryl(or alkyl)amino-4-ethoxycarbonyl-2-methylthio-1,3-thiazoles, 2a-2f, was verified with an 40% excess of KBu^tO, and π -deficient and π -excess aryl isothiocyanates, and alkyl isothiocyanates. The results have been collected in Table 2.

All new compounds have been fully characterized from their ir, ¹H-nmr, ¹³C-nmr, and ms data together a satisfactory analytical data. The ir, ¹H-, ¹³C-nmr, and ms key data have been gathered in Table 3.

The ir data support the presence of an amine and conjugated carbonyl groups. The single band at 3220-3420 cm⁻¹ can be assigned to a NH group,⁹ and the band at 1660-1730 cm⁻¹ to a carbonyl group of a conjugated ester with an unsaturated system (Ref. 9, p. 177).

Table 2. Observed yields by the synthesis of 5-aryl(or alkyl)amino-4-ethoxycarbonyl-2-methylthio-1,3-thiazoles 2a-2f.

1,3-thiazol	R	Yield(%) ^a	Mp(°C)
2a	Ph	90	79-80 ^b
2b	<i>p</i> -FC ₆ H ₄	65	54-55 ^c
2c	<i>p</i> -MeC ₆ H ₄	73	64-65 ^b
2d	<i>p</i> -MeOC ₆ H ₄	55	57-58 ^b
2e	Et	59	liquid ^d
2f	<i>n</i> -Bu	77	liquid ^d

^aYield in purified product. ^bRecrystallized from methanol. ^cRecrystallized from hexane. ^dPurified by silica gel flash chromatography.

Table 3. Ir, ¹H-Nmr, ¹³C-Nmr, and Ms Key data of 1,3-thiazoles 2a-2f.

Compound	Ir (cm ⁻¹)		¹ H-Nmr							Ms	
	NH(s)	CO(s)	NH	SCH ₃	C2	C4	C5	CO	SCH ₃	M ⁺ (100%)	MeS-C≡S ⁺ (%)
2a	3400	1660	9.80 ^a	2.65	164.72 ^b	140.64	122.90	164.76	17.56	294	20
2b	3280	1690	9.58 ^a	2.62	156.89	146.20	122.80	164.49	17.59	312	44
2c	3220	1730	9.60 ^a	2.62	156.44	145.50	122.15	164.35	17.32	308	13
2d	3300	1700	9.47 ^a	2.59	158.17	133.84	121.54	164.34	17.37	324	25
2e	3310	1710	7.40 ^c	2.56	161.83	142.50	117.87	162.96	16.51	246	15
2f	3420	1700	7.27 ^c	2.52	162.73	143.02	118.33	163.56	17.08	274	12

^aBroadened singlet. ^bInterchangeable assignments. ^cBroadened triplet.


The ¹H-nmr data prove the presence of an ethyl ester, a methylthio, and a secondary amine groups. The signals observed at 2.52-2.65 ppm support the presence of a methylthio group bounded to a sp² carbon,¹⁰ and the signals at 9.47-9.80 ppm and 7.27-7.40 ppm an aromatic (2a-2d) or aliphatic (2e-2f) secondary amine groups (Ref. 10, p. 118).

The ¹³C-nmr data support the proposed structure to 2a-2f compounds. The assignment of signals to C2 and C4 carbons is proposed by comparison with the chemical shifts reported by us for C2 and C4 carbons of 5-aryl(or alkylthio)-4-ethoxycarbonyl-2-methylthio-1,3-thiazoles (C2: 161.73-163.74 ppm; C4: 137.74-141.46 ppm).⁵

The ms data are according to the proposed structure to 2a-2f 1,3-thiazoles. In all cases the base peak has been the molecular ion, and a common ion m/z-91 has been detected. This peak can be reasonably assigned to MeS-C≡S⁺ ion (Ref. 1, p. 244), and its formation precludes any other relative disposition of the sulfur heteroatoms in the molecular ion.

Thus, the synthesis of 1,3-thiazoles by this method has shown to be highly selective because four

different products can be originated taking into account the regioselectivities of condensation and cyclization steps⁶ (Scheme 2).



- Scheme 2 -

Although some related syntheses have been previously described -cyclocondensation reactions of nitrile ylides with monothioesters,¹¹ α-metallated isocyanides with carbon disulfide,⁸ monothio,¹² and dithioesters;¹³ EMIC with carbon disulfide,^{5,6} and rearrangement of mesoionic 1,3-oxazoles to mesoionic 1,3-thiazoles¹⁴ - none of these methods allows to obtain the perfunctionalized 1,3-thiazoles described in this paper.

EXPERIMENTAL

Melting points were determined in a Büchi 520 apparatus in capillary tubes and are uncorrected. The IR spectra were recorded on a Perkin Elmer 781 spectrometer. The ¹H- and ¹³C-nmr spectra were recorded on a Varian FT 80A spectrometer (79.542 MHz for ¹H and 20.00 MHz for ¹³C). Solutions in CDCl₃ (13% and 25% w/v, respectively) at 303°K were used. Chemical shifts are quoted in δ values using TMS as internal reference. Mass spectra were recorded on a Varian Matt 711 spectrometer by electron impact technique (70 eV). All elemental analyses were satisfactory. Silica gel 60 F254 plates (Merck) were used for TLC, and silica gel 273-400 mesh (Merck) was used for conventional flash column chromatographies.

Synthesis of **13** was accomplished from ethyl ester glycine hydrochloride (Aldrich Chemie), carbon disulfide, and methyl iodide following the procedure described previously by us.⁵ α -Metallated **1** was condensed with phenyl isothiocyanate in different base/solvent systems and/or reaction conditions (Table 1). The isolation of products was performed following a general procedure with special features gathered in Table 1.

General procedure. To a stirred solution of K^tBuO (0.785 g, 7 mmol) in 50 ml dry THF (at -78°C under nitrogen) was added dropwise a solution of **1** (1 g, 5 mmol) in 4 ml dry THF. After 0.5 h. at -78°C , a solution of 5 mmol of isothiocyanate in 4 ml dry THF was slowly added. After 30 min. at -78°C , the mixture was allowed to stand at room temperature for 2 h., and was then quenched with water, and extracted with diethyl ether (4x30 ml). The combined ethereal extracts were dried on MgSO_4 (12 h.), and evaporated. The product was isolated by precipitation with pentane and recrystallized (**2a-2d**), or from silica gel flash chromatography **2e**: *n*-hexane/ethyl acetate: 90/10 v/v; **2f**: *n*-hexane/ethyl acetate: 95/5 v/v.

4-Ethoxycarbonyl-2-methylthio-5-phenylamino-1,3-thiazol, 2a. Ir (KBr pellet) 3400, 2910-2820, 1660, 1620, 1590, 1570, 1420, 1255, 1205, 790 cm^{-1} . $^1\text{H-Nmr}$, $\delta(\text{ppm})$: 1.42 (t, 3H, CH_3CH_2 , $J=7.0$ Hz), 2.65 (s, 3H, SCH_3), 4.41 (q, 2H, CH_2CH_3 , $J=7.0$ Hz), 7.05-7.53 (m, 5H, Ph), 9.80 (bs, 1H, NH). $^{13}\text{C-Nmr}$, $\delta(\text{ppm})$: 14.56 (CH_3CH_2), 17.56 (SCH_3), 60.87 (CH_2CH_3), 118.61 (*p*-C), 122.90 (C5), 123.81 (*o*-C), 129.69 (*m*-C), 140.64 (C4), 155.79 (*ipso*-C), 164.72 (C2), 164.76 (CO_2Et). Ms (%): 296 (9.4), 295 (14.3), 294 (100.0), 248 (37.1), 247 (3.7), 215 (77.1), 214 (19.4), 91 (19.7), 77 (15.1).

4-Ethoxycarbonyl-5-*p*-fluorophenylamino-2-methylthio-1,3-thiazole, 2b. Ir (KBr pellet): 3280, 3100, 2990, 1690, 1670, 1600, 1580, 1470, 1230, 1200, 1190, 860, 780 cm^{-1} . $^1\text{H-Nmr}$, $\delta(\text{ppm})$: 1.43 (t, 3H, CH_3CH_2 , $J=7.0$ Hz), 2.62 (s, 3H, SCH_3), 4.41 (q, 2H, CH_2 , $J=7.0$ Hz), 6.81-7.34 (m, 4H, Ar), 9.58 (bs, 1H, NH). $^{13}\text{C-Nmr}$, $\delta(\text{ppm})$: 14.53 (CH_3CH_2), 17.59 (SCH_3), 60.95 (CH_2CH_3), 116.48 (d,*m*-C, $^2J(^{13}\text{C},^{19}\text{F})=22.6$ Hz), 120.28 (d,*o*-C, $^3J(^{13}\text{C},^{19}\text{F})=11.8$ Hz), 122.80 (C5), 137.06 (d, *ipso*-C, $^4J(^{13}\text{C},^{19}\text{F})=2.6$ Hz), 146.20 (C4), 156.89 (C2), 159.38 (d,*p*-C, $^1J(^{13}\text{C},^{19}\text{F})=244.4$ Hz), 164.69 (CO_2Et). Ms (%): 314 (10.3), 313 (19.1), 312 (100.0), 266 (45.6), 265 (4.1), 233 (69.1), 232 (45.6), 95 (19.1), 91 (44.1).

4-Ethoxycarbonyl-5-*p*-methylphenylamino-2-methylthio-1,3-thiazole, 2c. Ir (KBr pellet): 3220, 3000, 2900, 1730, 1660, 1590, 1580, 1420, 1270, 1200, 820, 780 cm^{-1} . $^1\text{H-Nmr}$, $\delta(\text{ppm})$: 1.42 (t, 3H, CH_3CH_2 , $J=7.1$ Hz), 2.32 (s, 3H, $\text{CH}_3\text{-Ar}$), 2.62 (s, 3H, SCH_3), 4.41 (q, 2H, CH_2CH_3 , $J=7.1$ Hz), 7.10 (m, 4H, Ar), 9.60 (bs, 1H, NH). $^{13}\text{C-Nmr}$, $\delta(\text{ppm})$: 14.22 (CH_3CH_2), 17.32 (SCH_3), 20.46 ($\text{CH}_3\text{-Ar}$), 60.47 (CH_2CH_3), 118.38 (*o*-C), 122.15 (C5), 129.84 (*p*-C), 133.48 (*ipso*-C), 137.92 (*m*-C), 145.50 (C4), 156.44 (C2), 164.35 (CO_2Et). Ms (%): 310 (9.2), 309 (14.9), 308 (100.0), 262 (48.6), 261 (8.1), 229 (64.5), 228 (16.7), 91 (13.2).

4-Ethoxycarbonyl-5-*p*-methoxyphenylamino-2-methylthio-1,3-thiazole, 2d. Ir (KBr pellet): 3300, 3160, 3110, 2990, 1700, 1660, 1540, 1510, 1480, 1300, 1270, 770, 670 cm^{-1} . $^1\text{H-Nmr}$, $\delta(\text{ppm})$: 1.41 (t, 3H, CH_3CH_2 , $J=7.0$ Hz), 2.59 (s, 3H, SCH_3), 3.78 (s, 3H, OCH_3), 4.42 (q, 2H, CH_2CH_3 , $J=7.0$ Hz), 6.77-7.47 (m, 4H, Ar), 9.47 (bs, 1H, NH). $^{13}\text{C-Nmr}$, $\delta(\text{ppm})$: 14.26 (CH_3CH_2), 17.37 (SCH_3), 55.19 (OCH_3), 60.44 (CH_2CH_3), 113.92 (*ipso*-C), 114.54 (*m*-C), 121.14 (*o*-C), 121.54 (C5), 133.89 (C4), 156.48 (*p*-C), 158.17 (C2), 164.34 (CO_2Et). Ms (%): 326 (10.6), 325 (15.8), 324 (100.0), 278 (79.0), 277 (5.3), 245 (31.1), 244 (21.6), 91 (25.3).

4-Ethoxycarbonyl-5-ethylamino-2-methylthio-1,3-thiazole, 2e. Ir (film): 3310, 2980, 2860, 1710, 1660, 1550, 1390, 1220, 1150 cm^{-1} . $^1\text{H-Nmr}$, $\delta(\text{ppm})$: 0.93 (t, 3H, $\text{CH}_3\text{CH}_2\text{NH}$, $J=7.1$ Hz), 1.35 (t, 3H, $\text{CH}_3\text{CH}_2\text{OCO}$, $J=7.0$ Hz), 2.56 (s, 3H, SCH_3), 3.11 (qd, 2H, $\text{CH}_3\text{CH}_2\text{NH}$, $J=7.1$, 5.0 Hz), 4.27 (q, 2H, $\text{CH}_3\text{CH}_2\text{OCO}$, $J=7.0$ Hz), 7.40 (bt, 1H, NH, $J=5.0$ Hz). $^{13}\text{C-Nmr}$, $\delta(\text{ppm})$: 13.39 ($\text{CH}_3\text{CH}_2\text{OCO}$, $\text{CH}_3\text{CH}_2\text{NH}$), 16.51 (SCH_3), 42.88 ($\text{CH}_3\text{CH}_2\text{NH}$), 58.85 ($\text{CH}_3\text{CH}_2\text{OCO}$), 117.87 (C5), 142.50 (C4), 161.83 (C2), 162.96 (CO_2Et). Ms (%): 248 (8.3), 247 (9.2), 246 (100.0), 200 (20.2), 199 (3.5), 167 (25.1), 166 (22.1), 91 (15.0).

4-Ethoxycarbonyl-5-*n*-butylamino-2-methylthio-1,3-thiazole, 2f. Ir (film): 3420, 2900, 1700, 1660, 1550, 1460, 1420, 1200, 1150 cm^{-1} . $^1\text{H-Nmr}$, $\delta(\text{ppm})$: 0.93 (t, 3H, $\text{CH}_3(\text{CH}_2)_3$, $J=7.0$ Hz), 1.35 (m, 4H, $\text{CH}_3(\text{CH}_2)_2\text{CH}_2\text{NH}$), 1.42 (t, 3H, $\text{CH}_3\text{CH}_2\text{OCO}$, $J=7.1$ Hz), 2.52 (s, 3H, SCH_3), 3.10 (m, 2H, $\text{CH}_3(\text{CH}_2)_2\text{CH}_2\text{NH}$), 4.27 (q, 2H, $\text{CH}_3\text{CH}_2\text{OCO}$, $J=7.1$ Hz), 7.27 (bt, 1H, NH, $J=7.27$ Hz). $^{13}\text{C-Nmr}$, $\delta(\text{ppm})$: 12.87 ($\text{CH}_3(\text{CH}_2)_3$), 13.82 ($\text{CH}_3\text{CH}_2\text{OCO}$), 17.08 (SCH_3), 19.08 ($\text{CH}_3\text{CH}_2(\text{CH}_2)_2$), 30.51 ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}$), 48.18 ($\text{CH}_3(\text{CH}_2)_2\text{CH}_2\text{NH}$), 59.40 ($\text{CH}_3\text{CH}_2\text{OCO}$), 118.33 (C4), 143.02 (C5), 162.73 (C2), 163.73 (CO_2Et). Ms (%): 276 (8.7), 275 (10.1), 274 (100.0), 228 (35.3), 227 (4.1), 195 (40.1), 194 (8.3), 91 (12.0).

REFERENCES AND NOTES.

1. A. R. Katritzky and Ch. W. Rees, 'Comprehensive Heterocyclic Chemistry. The Structure, Reactions, Synthesis and Uses of Heterocyclic Compounds', Vol. 6, ed. by A.R. Katritzky, Pergamon Press, New York, 1982, pp. 273-293.
2. a) E. Monserrat Faba, ES. P. 542,806 (Chem. Abst., 1987, 106, 84592n); b) Lab. Fides, ES. P. 548,523 (Chem. Abst., 1987, 106, 84591m); c) F. López Calahorra, F. A. Geijo, J. M. Vallés, and J. Prats ES. P. 544,816 (Chem. Abst., 1987, 107, 77786v); d) P. Airno, ES. P. 544,597 (Chem. Abst., 1987, 107, 77787w); e) L. Salvador, ES. P. 548,521 (Chem. Abst., 1987, 107, 77788y); f) G. Richter and G. Rt. Vegyeszeti, BE. P. 905,409 (Chem. Abst., 1987, 107, 39795r).
3. L. A. Reiter, EP. P. 161,841 (Chem. Abst., 1986, 104, 168456t).
4. a) A. Bycroft and M. S. Gowland, J. Chem. Soc., Chem. Commun., 1978, 256; b) H. Sugne, J. Am. Chem. Soc., 1982, 104, 5537; c) C. M. Ireland, J. Org. Chem., 1982, 47, 1807; d) S. Halazy,

- Tetrahedron Lett., 1984, 1421.
5. C. Alvarez Ibarra, E. Dávila, A. Mateo, P. Ortiz, and M. L. Guiroga, Tetrahedron Lett., 1987, 6667.
 6. W. Dolling, P. Kindt, and M. Augustin, Z. Chem., 1987, 27, 212.
 7. Spectroscopic data of bis(5-(4-ethoxycarbonyl-2-methylthio-1-phenyl)imidazolyl)disulfane: Ir (CHCl₃): 1700, 1540, 1520, 1400, 1220 cm⁻¹. ¹H-Nmr, δ(ppm): 1.38 (t, 3H, CH₃CH₂, J=7.12 Hz), 2.66 (s, 3H, SCH₃), 4.29 (q, 2H, CH₂CH₃, J=7.12 Hz), 7.02-7.55 (m, 5H, Ph). ¹³C-Nmr, δ(ppm): 14.18 (CH₃CH₂), 14.49 (SCH₃), 60.68 (CH₂CH₃), 128.20, 128.53, 129.00, 129.56 (aromatic carbons), 133.98 (-CO-C=), 138.69 (-S-C=), 149.66 (-S-C=N), 161.33 (CO₂Et).
 8. S. P. J. van Nispen, J. H. Bregman, D. G. van Engen, A. M. van Leusen, M. Saikachi, J. Kitigawa and H. Sasaki, J. Royal Neth. chem. Soc., 1982, 101, 28.
 9. R. T. Conley, 'Infrared Spectroscopy', Allyn and Bacon Inc., Boston, 2nded., 1972, pp. 143-146.
 10. E. Pretsch, T. Clerc, J. Seibl, and W. Simon, 'Tablas para la elucidación estructural de compuestos orgánicos por métodos espectroscópicos', Alhambra, Madrid, 1980, p. 122.
 11. H. Bunge, R. Huisgen, R. Raab, and H.J. Sturman, Chem. Ber., 1972, 105, 1037.
 12. G. D. Hartman and L. M. Weisntock, Synthesis, 1976, 681.
 13. W. Schollkopf, P. H. Porsch, and E. Blume, Liebigs Ann. Chem., 1976, 2212.
 14. G. C. Barrett and R. Walker, Tetrahedron, 1976, 32, 583.

Received, 23rd April, 1988