Metal-Containing Oligonucleotides: Solid-Phase Synthesis and Luminescence Properties

Dennis J. Hurley and Yitzhak Tor*

Department of Chemistry and Biochemistry University of California, San Diego La Jolla, California 92093-0358

Received November 24, 1997

The incorporation of photo- and redox-active transition metal ions into oligonucleotides is a key design target for the study of energy and electron-transfer processes through DNA,¹ as well as the development of DNA hybridization probes and sensors.² Metal-containing oligonucleotides have been predominantly constructed via two major pathways: (a) the synthesis of a chelatorcontaining oligonucleotide followed by metal complexation³ and (b) the synthesis of an end-functionalized oligonucleotide to which a metal complex can be conjugated.^{1,4} These approaches are restricted primarily to modifications at the oligonucleotides' termini and/or require the exposure of oligonucleotides to reactive metal precursors.²⁻⁵ A direct method for the site-specific incorporation of metal complexes during solid-phase oligonucleotide synthesis has never been reported.

We now disclose a general methodology for the incorporation of polypyridine metal complexes into oligonucleotides using automated DNA synthesizers. We report the synthesis of novel Ru^{II}- and Os^{II}-containing nucleosides and their phosphoramidite derivatives. These building blocks are sequence-specifically incorporated into oligonucleotides in high yields using standard solid-phase phosphoramidite chemistry. The uniquely modified oligonucleotides form stable DNA duplexes and are useful probes for the study of energy-transfer processes in nucleic acids.

We have previously reported that functionalized tris-chelate complexes are excellent substrates for the powerful palladiummediated cross-coupling methodologies.⁶ This approach provides a convenient entry into metal-containing nucleosides and is key

Pfleiderer, W.; Müller, F. Helv. Chim. Acta 1991, 74, 1991-1999. (d) Telser, J.; Cruickshank, K. A.; Schanze, K. S.; Netzel, T. L. J. Am. Chem. Soc. **1989**, 111, 7221–7226. (e) Sammes, P. G.; Yahioglu, G. Nat. Prod. Rep. **1996**, 13, 1-28. (f) Ihara, T.; Nakayama, M.; Murata, M.; Nakano, K.; Maeda, M. Chem. Commun. 1997, 1609-1610.

(3) Bashkin, J. K.; Frolova, E. I.; Sampath, U. J. Am. Chem. Soc. 1994, 116, 5981-5982. Matsumura, K.; Endo, M.; Komiyama, M. J. Chem. Soc., Chem. Commun. 1994, 2019-2020. Dreyer, G. R.; Dervan, P. B. Proc. Natl. Acad. Sci. U.S.A. 1985, 82, 968-972. Chen, C.-H. B.; Sigman, D. S. J. Am. Chem. Soc. 1988, 110, 6570-6572.

(4) Magda, D.; Miller, R. A.; Sessler, J. L.; Iverson, B. L. J. Am. Chem. (c) Magda, D., Miller, R. A., Sessiel, J. L., Iverson, B. L. J. M., Chem. Soc. **1994**, *116*, 7439–7440. Hall, J.; Hüsken, D.; Pieles, U.; Moser, H. E.; Häner, R. Chem. Biol. **1994**, *1*, 185–190. Hall, J.; Hüsken, D.; Häner, R. Nucleic Acids Res. 1996, 24, 3522-3526.

(5) For recent approaches utilizing metal-containing phosphoramidites or H-phosphonates, see: ref 2b. Manchanda, R.; Dunham, S. U.; Lippard, S. J. J. Am. Chem. Soc. **1996**, 118, 5144–5145. Schliepe, J.; Berghoff, U.; Lippert, B.; Cech, D. Angew. Chem., Int. Ed. Engl. **1996**, 35, 646–648. Mucic, R. C.; Herrlien, M. K.; Mirkin, C. A.; Letsinger, R. L. Chem. Commun. **1996**, 557. 557. Magda, D.; Crofts, S.; Lin, A.; Miles, D.; Wright, M.; Sessler, J. L. J. Am. Chem. Soc. **1997**, 119, 2293–2294. For a photoaddition of Ru^{II} complex

I. M. Soc. 1997, 119, 1229, 14, Kirsch-De Mesmaeker, A.; Kelly, J. M. J. Am. Chem. Soc. 1997, 119, 11763–11768.
 (6) Tzalis, D.; Tor, Y. Chem. Commun. 1996, 1043–1044. Tzalis, D.; Tor, Y. J. Am. Chem. Soc. 1997, 119, 852–853. Connors, P. J., Jr.; Tzalis, D.; Dunnick, A. L.; Tor, Y. Inorg. Chem. In press.

Scheme 1. Synthesis of Phosphoramidites 6a and 6b^{*a,b*}



^a Reagents: (a) 4,4'-dimethoxytrityl chloride (DMT-Cl), DMAP, pyridine, Et₃N, 92% yield; (b) **3a** or **3b**, (Ph₃P)₂PdCl₂, CuI, DMF, Et₃N, sonication, 84% yield; (c) (*i*Pr₂N)₂POCH₂CH₂CN, (1*H*)-tetrazole, CH₃CN; 70-85% yield. ^b All metal-modified nucleosides were isolated as their PF6⁻ salts.⁸

to the successful preparation of the modified nucleosides and their phosphoramidites. Thus, palladium-catalyzed cross-coupling reactions between 5-ethynyldeoxyuridine⁷ (1) and [(bpy)₂Ru(3bromo-1,10-phenanthroline)]²⁺(PF₆⁻)₂ (**3a**) or [(bpy)₂Os(3-bromo-1,10-phenanthroline)]²⁺(PF₆⁻)₂ (bpy = bipyridyl) (**3b**) afford nucleosides 4a and 4b, respectively (Scheme 1).8,9 The mild conditions of this reaction allow us to apply it for the modification of 4,4'-dimethoxytrityl-protected nucleosides. Thus, 1 is first treated with 4,4'-dimethoxytrityl (DMT) chloride in the presence of 4-(dimethylamino)pyridine (DMAP) to provide the DMTprotected nucleoside 2, which is then cross-coupled to 3a or 3b to afford the protected metal-containing nucleosides 5a and 5b, respectively (Scheme 1). Phosphitylation of the protected nucleosides 5a and 5b using (2-cyanoethoxy)bis(diisopropylamino)phosphine in the presence of (1H)-tetrazole provides the corresponding metal-modified phosphoramidites 6a and 6b.¹⁰

Target 20-mer oligonucleotides incorporating one or two metalmodified 2'-deoxyuridine bases at various positions were synthesized on a 0.2 μ mol scale using an automated DNA synthesizer (Figure 1). When coupling times for the phosphoramidites 6a and 6b in 0.5 M (1H)-tetrazole were extended to 5 min, reaction efficiencies were greater than 90%.¹¹ Removal of the finished 20-mers from the solid support using concentrated ammonium hydroxide was followed by incubation at 55 °C for 8 h to afford

⁽¹⁾ Murphy, C. J.; Arkin, M. R.; Jenkins, Y.; Ghatlia, N. D.; Bossmann, S. H.; Turro, N. J.; Barton, J. K. *Science* **1993**, *262*, 1025–1029. Meade, T. S. H.; Turro, N. J.; Barton, J. K. Science 1993, 262, 1025–1029. Meade, T. J.; Kayyem, J. F. Angew. Chem., Int. Ed. Engl. 1995, 34, 352–354. Dandliker, P. J.; Holmlin, R. E.; Barton, J. K. Science 1997, 275, 1465–1468. Meggers, E.; Kusch, D.; Giese, B. Helv. Chim. Acta 1997, 80, 640–652. See also: Netzel, T. L. J. Chem. Educ. 1997, 74, 646–651. Diederichsen, U. Angew. Chem., Int. Ed. Engl. 1997, 36, 2317–2319.
(2) (a) Bannwarth, W.; Schmidt, D.; Stallard, R. L.; Hornung, C.; Knorr, R.; Müller, F. Helv. Chem. Acta 1988, 71, 2085–2099. (b) Bannwarth, W.; Schmidt, D. Tetrahedron Lett. 1989, 30, 1513–1516. (c) Bannwarth, W.; Pleiderer W.: Müller F. Helv. Chim. Acta 1917, 74, 1940–1999. (d) Telser

⁽⁷⁾ Robins, M. J.; Barr, P. J. J. Org. Chem. 1983, 48, 1854-1862.

 ⁽⁸⁾ See the Supporting Information for experimental details.
 (9) Polypyridine complexes of Ru^{II} and Os^{II} were selected due to their chemical stability and favorable redox and photophysical characteristics. See: Sauvage, J.-P.; Collin, J.-P.; Chambron, J.-C.; Guillerez, S.; Coudret, C.; Balzani, V.; Barigelletti, F.; De Cola, L.; Flamigni, L. *Chem. Rev.* **1994**, 94, 993-1019. Balzani, V.; Juris, A.; Venturi, M.; Campagna, S.; Serroni, S. Chem. Rev. 1996, 96, 759-833.

⁽¹⁰⁾ All compounds were characterized by ¹H NMR, ESI-MS, UV-vis, IR and square-wave and cyclic voltammetry. See the Supporting Information.

⁽¹¹⁾ To control the amount of reagents and reaction time, the coupling of the modified bases was performed manually (ref 8). The decreased coupling efficiencies relative to standard phosphoramidites are likely due to the steric bulk of the appended metal complex and have been reported with other bulky phosphoramidites. See: Kobertz, W. R.; Essigmann, J. M. J. Am. Chem. Soc. 1997, 119, 5960–5961.

7	5'	ТCG	GCG	CGA	ΑΤΤ	CGC	GTG	СС	3'
8	5 '	$\mathrm{T}\mathrm{C}\mathrm{G}$	GCG	CGA	A U T	CGC	GTG	СС	3 '
9	5 '	ТCG	GCG	CGA	а ў т	CGC	GTG	СС	3'
10	5 '	T C G	GCG	CGA	ΑΤΤ	CGC	GTG	СС	3'
11	5 '	ТCG	GCG	CGA	А й т	CGC	$\mathbf{G} \overset{\mathtt{Ru}}{\boldsymbol{U}} \mathbf{G}$	CC	3'
12	3 '	AGC	CGC	GСТ	ΤΑΑ	GCG	CAC	$\mathbf{G}\mathbf{G}$	5'
13	3'	AGC	CGC	$\operatorname{GC} \overset{\mathbf{os}}{\boldsymbol{U}}$	ΤΑΑ	GCG	CAC	GG	5'
14	3'	AGC	CGC	GCT	Τ Τ Α	GCG	CAC	GG	5'

Figure 1. Sequences of oligonucleotides synthesized. The Ru^{II}- and Os^{II}- containing deoxyuridine nucleosides (**4a** and **4b**, respectively) are shown in bold.



Figure 2. Thermal denaturation curves for control duplex $7 \cdot 12$ (\bullet), Os^{II}containing duplex $9 \cdot 12$ (\triangle), and a single-mismatch duplex $7 \cdot 14$ (\Box) determined in 0.01 M sodium phosphate buffer pH 7, 0.1 M NaCl.⁸

the deprotected oligomers 8-11 and 13 that were purified by gel electrophoresis.⁸ Analytical denaturing polyacrylamide gel electrophoresis confirmed the purity of the modified oligonucleotides, and enzymatic digestion followed by HPLC analysis verified the presence of the intact metal-containing nucleosides.⁸

The presence of the a metal-containing nucleoside has a relatively small effect on duplex stability as determined by thermal denaturation curves (Figure 2). The melting temperature (T_m) of the unmodified duplex derived from oligonucleotide 7 and its complementary sequence 12 is 78 °C. When the metal-containing nucleoside is located at the 5'-end, as in duplex 10·12, the T_m is essentially the same. Duplexes 8·12 and 9·12, in which the metal-containing nucleoside is in the middle of the duplex, are slightly less stable with a T_m at 75 °C. Yet, this destabilization is far smaller than the effect of a single mismatch on duplex stability as demonstrated for duplex 7·14 containing a T–T "pair" at the same position ($T_m = 69$ °C, Figure 2).

Steady-state emission profiles of iso-absorptive oligonucleotide solutions in degassed phosphate buffer are shown in Figure 3. The Ru^{II}-containing duplex **8**•12 shows a typical metal-centered emission at 630 nm upon excitation of the visible metal-to-ligand charge-transfer (MLCT) band at 456 nm. Upon hybridization of the Ru^{II}-containing oligonucleotide **8** to a complementary Os^{II}-containing oligonucleotide **13**, a substantial drop in the emission intensity (ca. 70-85%) is observed. This suggests an "intraduplex" quenching of the excited Ru^{II} center by the proximal Os^{II}



Figure 3. Steady-state emission spectra of modified oligonucleotides in degassed 0.01 M sodium phosphate buffer pH 7.0, 0.1 M NaCl. Top: duplex 8·12 (solid line), a 1:1 mixture of 8 and 9 (---), and a duplex containing proximal Ru and Os 8·13 (dashed line). Bottom: duplex 10·12 (solid line) and duplex 10·13 (dashed line).

center.^{9,12} Intermolecular quenching can be excluded since a 1:1 mixture of the noncomplementary oligonucleotides **8** and **9** shows essentially the same emission intensity as duplex **8·12** (Figure 3a, top). This behavior is distance-dependent as demonstrated by comparing duplex **8·13** to duplex **10·13**. In this case, where the Os^{II} center is more remote, only 40% quenching of the Ru^{II}-based emission is observed (Figure 3b, bottom). To the best of our knowledge, this is the first example of energy-transfer processes in DNA oligonucleotides that are sequence-specifically modified with polypyridine metal complexes.

The data presented here establish a novel and powerful approach for the site-specific incorporation of polypyridine—metal complexes into synthetic oligonucleotides using automated phosphoramidite chemistry. The versatile phosphoramidite synthesis and the compatibility with existing automated DNA synthesizers provides enormous flexibility for rapid construction of oligonucleotide—metal conjugates. The presence of photo- and redoxactive metal centers in these oligonucleotides makes them extremely important probes for the study of photophysical processes in nucleic acids.

Acknowledgment. We thank Clinical Micro Sensors, Inc., and the UC Biotechnology STAR Project for generous support of this research and Professor David R. Kearns for the use of his temperature-controlled UV spectrophotometer.

Supporting Information Available: Synthetic procedures and analytical data for all new derivatives as well as procedures and data for oligonucleotide synthesis, purification, digestion, melting, and fluorescence studies (13 pages). See any current masthead page for ordering information and Web access instructions.

JA9739998

⁽¹²⁾ De Cola, L.; Barigelletti, F.; Balzani, V.; Belser, P.; Von Zelewsky, A.; Seel, C.; Frank, M.; Vögtle, F. in *Supramolecular Chemistry*; Balzani, V., De Cola, L., Eds, Kluwer: Dordrecht, The Netherlands, 1992; pp 157–180.