

Dalton Communications


Synthesis and Crystal Structure of $[\{\text{Au}_2(\text{C}_6\text{H}_4\text{S}_2-1,2)(\text{PEt}_3)\}_2]$ Masami Nakamoto,^a Annette Schier^b and Hubert Schmidbaur^{*,b}^a Osaka Municipal Technical Research Institute, 6-50, 1-Chome, Morinomiya, Joto-ku, Osaka 536, Japan^b Anorganisch-chemisches Institut der Technischen Universität München, Lichtenbergstrasse 4, D-8046 Garching, Germany

Treatment of chloro(triethylphosphine)gold(I) with benzene-1,2-dithiol in the presence of a tertiary amine base affords $[\{\text{Au}_2(\text{C}_6\text{H}_4\text{S}_2-1,2)(\text{PEt}_3)\}_2]$ (60.4% Au by weight) which has a compact, centrosymmetric structure containing a rhomboid Au_4 unit supported by the sulfur and phosphorus ligands.

Gold(I) thiolate complexes have received continued interest in recent years owing to their widespread use in medicine,^{1,2} electrical conductor and semiconductor technology,^{3,4} and in the glass and ceramics industry,^{5,6} e.g., a (triethylphosphine)-(thioglucose)gold(I) complex is one of the most successful drugs in the therapy of arthritis, and sulfur-rich liquid-gold pastes, largely based on natural products, are the traditional gold sources for the regioselective gilding of materials.


In the course of our studies on polynuclear gold(I) compounds with main-group element co-ordination centres^{7,8} we have now obtained a novel tetranuclear complex with some interesting properties: (i) it is readily obtained in pure form; (ii) it is stable to air and water; (iii) it is soluble in polar organic solvents; (iv) it has an exceedingly high gold content (60.4% by weight), since half of the auxiliary phosphine ligand provided in the synthesis is not retained owing to the formation of a compact, thermally stable cluster structure.

Treatment of chloro(triethylphosphine)gold(I) with benzene-1,2-dithiol in the presence of triethylamine as a base in tetrahydrofuran at ambient temperature affords a light yellow crystalline product (m.p. 198 °C with decomposition) of stoichiometry $\text{Au}_2(\text{C}_6\text{H}_4\text{S}_2)(\text{PEt}_3)$ as determined by elemental analysis.[†] The bis(phosphine) complex $[\text{Au}_2(\text{C}_6\text{H}_4\text{S}_2-1,2)(\text{PEt}_3)_2]$ suggested by the stoichiometry of the reagents and a green compound identified as a mixed-valent salt $[\text{Au}(\text{PEt}_3)_2][\text{Au}(\text{C}_6\text{H}_4\text{S}_2-1,2)]^9$ (probably originating from partially oxidized dithiole) are by-products of the reaction [equation (1)]. Solutions of complex **1** in chloroform show a singlet



resonance in the $^{31}\text{P}\{-^1\text{H}\}$ NMR spectrum at δ 32.5. The ^1H NMR spectrum shows the pattern of an $\text{A}_3\text{B}_2\text{X}$ spin system for the PEt_3 groups [δ 1.01 and 1.43, $J(\text{HH})$ 7.6, $J(\text{PH})$ 18.5 and 9.6

[†] To a solution of $[\text{AuCl}(\text{PEt}_3)]$ (0.65 g, 1.87 mmol) in tetrahydrofuran (10 cm³) a mixture of $\text{C}_6\text{H}_4(\text{SH})_2-1,2$ (0.13 g, 0.91 mmol) and NEt_3 (0.19 g, 1.87 mmol) in the same solvent (10 cm³) was added dropwise at room temperature with stirring to give a faintly green solution and a white precipitate. The solvent was evaporated after 1 h, the residue was treated with water (20 cm³) to remove $\text{NHEt}_3^+\text{Cl}^-$, collected by filtration and washed with water (2 \times 10 cm³) and diethyl ether (2 cm³) to give 0.55 g of crude product. Crystallization from chloroform-hexane (1:1) afforded 0.21 g of pure material (35% yield). The by-products can be isolated from the mother-liquor⁹ (Found: C, 22.05; H, 2.95; Au, 60.40; P, 4.75; S, 9.85. $\text{C}_{12}\text{H}_{10}\text{Au}_2\text{PS}_2$ requires C, 22.10; H, 2.95; Au, 60.40; P, 4.75; S, 9.85%).



Hz, respectively] and of an $\text{ABB}'\text{A}'$ spin system for the dithiolate group (δ 7.03 and 7.99). In the $^{13}\text{C}\{-^1\text{H}\}$ NMR spectrum, the carbon resonances appear at δ 9.0 (Me), 19.1 [CH_2 , $J(\text{PC})$ 31.8 Hz], 125.3, 135.3 and 144.4 (C_6H_4). These data are distinct from those of the by-products (above).⁹

According to fast atom bombardment mass spectrometry data (*m*-nitrobenzyl alcohol, m/z 1304), complex **1** forms a dimer the structure of which has been determined by single-crystal X-ray diffraction.[‡]

Crystals of **1** (from chloroform-*n*-hexane) are monoclinic,

[‡] Crystal data. $\text{C}_{24}\text{H}_{38}\text{Au}_4\text{P}_2\text{S}_4$, $M_r = 1304.64$, colourless crystals (0.25 \times 0.30 \times 0.35 mm), monoclinic, space group $P2_1/n$ (no. 14), $a = 9.866(2)$, $b = 13.274(1)$, $c = 12.776(2)$ Å, $\beta = 103.46(2)^\circ$, $U = 1627.2$ Å³, $Z = 2$, $D_c = 2.662$ g cm⁻³, $F(000) = 1184$. Enraf-Nonius CAD4 diffractometer, graphite-monochromated Mo-K α radiation ($\lambda = 0.71069$ Å), $\mu(\text{Mo-K}\alpha) = 182.81$ cm⁻¹, $T = 22^\circ\text{C}$, θ - 2θ scan mode. During data collection three standard reflections were periodically measured as a general check of crystal and instrument stability. No significant change was observed. Reduced-cell calculations (DELOS, LAPAGE)^{10,11} did not indicate any higher symmetry. Intensity data were corrected for absorption effects (DIFABS).¹² The structure was solved by direct methods and refined by full-matrix least-squares calculations (SHELXS 86,¹³ SHELX 76¹⁴). Of 4248 measured reflections, 3292 were unique, and 2412 were considered 'observed' [$F_o \geq 4.0\sigma(F_o)$] and used for refinement. Thermal motion was treated anisotropically for all non-hydrogen atoms except for carbon atoms C(22) and C(32) of two of the ethyl groups, which were disordered in split positions with site-occupancy factors of 0.5:0.5 and 0.6:0.4 for C(22):C(222) and C(32):C(322), respectively. Hydrogen atoms of ethyl groups were neglected, ring hydrogen atoms were placed in calculated fixed positions with isotropic displacement parameters ($U_{\text{iso}} = 0.05$ Å²). Number of refined parameters 152. The functions minimized were $R = \Sigma(|F_o| - |F_c|)/|F_o|$, $R' = [\Sigma w(|F_o| - |F_c|)^2 / \Sigma w F_o^2]^{1/2}$, $w = 1/\sigma^2(F_o)$. Final R and R' values are 0.061 and 0.075, respectively. Residual electron density is in the range +3.72 to -4.42 e Å⁻³ (near the gold atoms). Atomic coordinates, thermal parameters and bond lengths and angles have been deposited at the Cambridge Crystallographic Data Centre. See Instructions for Authors, *J. Chem. Soc., Dalton Trans.*, 1993, Issue 1, pp. xxiii-xxviii.




Fig. 1 Molecular structure of $[\{Au_2(C_6H_4S_2-1,2)(PEt_3)_2\}_2]$ with atomic numbering (the molecule has a centre of inversion). Selected bond lengths (Å) and angles (°): Au(1)···Au(1a) 3.114(1), Au(1)···Au(2) 3.052(1), Au(1)···Au(2a) 3.129(1), Au(1)–S(1) 2.311(5), Au(1)–S(2) 2.294(4), Au(2)–S(1) 2.407(5), Au(2)–S(2a) 2.625(4), Au(2)–P(1) 2.224(6); Au(1)–S(1)–Au(2) 80.6(1), Au(1)–S(2)–Au(2a) 78.7(1), Au(1)···Au(2)···Au(1a) 60.5(1), Au(2)···Au(1)···Au(2a) 119.5(1), S(1)–Au(1)–S(2) 173.3(1), S(1)–Au(2)–P(1) 149.1(2), S(2a)–Au(2)–P(1) 126.5(2), S(1)–Au(2)–S(2a) 84.4(1)

space group $P2_1/n$, with two molecules in the unit cell. There is no solvent contained in the crystals. The structure of the complex is shown in Fig. 1. The molecule has a crystallographic centre of inversion which relates the two chelating benzene-1,2-dithiolato ligands and the two triethylphosphine units. The four gold atoms are arranged in the form of a parallelogram with Au···Au edges of 3.052(1) and 3.129(1) Å, and a short transannular Au···Au distance of 3.114(1) Å. The gold atoms at the wide-angle vertices of the parallelogram (119.5°) are two-co-ordinate with S–Au–S angles of 173.3° , while those at the acute-angle vertices (60.5°) are three-co-ordinate in a quasi-planar environment of one phosphorus and two sulfur atoms. Each of the bidentate dithiolate ligands is bridging three gold atoms, while the triethylphosphines are monodentate. There is a slight disorder of the ethyl groups of these units in the crystal, which, however, could be accounted for by a suitable computational model.

With $[AuCl(PPh_3)]$ instead of $[AuCl(PEt_3)]$ as the starting material, no PPh_3 is lost under similar reaction conditions to those used above,⁹ and a complex $[Au_2(C_6H_4S_2-1,2)(PPh_3)_2]$ is obtained as the sole product. Studies with a variety of other ligands have therefore been initiated in order to investigate the influence of electronic and/or steric effects on the stoichiometry of the reaction. (*Note added at proof.* For a methyl homologue, derived from toluene-3,4-dithiol, see ref. 15).

Acknowledgements

This work was supported by the Deutsche Forschungsgemeinschaft (Leibniz-Programm), the Fonds der Chemischen Industrie, and, through the donation of chemicals, by Degussa AG and Heraeus GmbH. We are indebted to Mr. J. Riede for collecting the X-ray data set. M. N. is grateful to the Yamada Foundation, Tokyo, for a fellowship.

References

- 1 K. C. Dash and H. Schmidbaur, in *Metal Ions in Biological Systems*, ed. H. Sigel, Marcel Dekker, New York, 1982, vol. 14, p. 179.
- 2 B. M. Sutton, *Gold Bull.*, 1986, **19**, 15; A. A. Isab, A. L. Hormann, D. T. Hill, D. E. Griswold, M. J. Dimartino and C. F. Shaw III, *Inorg. Chem.*, 1989, **28**, 1321.
- 3 W. S. Rapson, *Gold Usage*, Academic Press, London, 1978.
- 4 R. J. Puddephatt, *The Chemistry of Gold*, Elsevier, Lausanne, 1978.
- 5 L. B. Hunt, *Gold Bull.*, 1979, **12**, 116.
- 6 *Im Zeichen von Sonne und Mond*, ed. Degussa AG, Frankfurt, 1993.
- 7 H. Schmidbaur, *Gold Bull.*, 1990, **23**, 11.
- 8 H. Schmidbaur, *Interdisc. Sci. Rev.*, 1992, **17**, 213.
- 9 M. Nakamoto, W. Hiller and H. Schmidbaur, *Chem. Ber.*, 1993, **126**, 605; M. Nakamoto, H. Koijman, M. Paul, W. Hiller and H. Schmidbaur, *Z. Anorg. Allg. Chem.*, in the press and refs. therein.
- 10 H. Zimmermann and H. Burzlaff, *Z. Kristallogr.*, 1985, **170**, 241.
- 11 Y. LePage, *J. Appl. Crystallogr.*, 1982, **15**, 255.
- 12 N. Walker and D. Stuart, *Acta Crystallogr., Sect. A*, 1983, **39**, 158.
- 13 G. M. Sheldrick, SHELXS 86, in *Crystallographic Computing 3*, eds. G. M. Sheldrick, C. Krüger and R. Goddard, Oxford University Press, Oxford, 1985.
- 14 G. M. Sheldrick, SHELX 76, Program for Crystal Structure Determination, University of Cambridge, 1976.
- 15 R. M. Davila, A. Elduque, T. Grant, R. J. Staples and J. P. Fackler, jun., *Inorg. Chem.*, in the press.

Received 8th February 1993; Communication 3/00760J