

Preliminary Note

3-Trifluoromethylthiophens from hexafluoroacetone

Klaus Burger*, Brigitte Helmreich and Oliver Jendrewski

Institut für Organische Chemie, Technische Universität München, Lichtenbergstr. 4, D-85747 Garching (Germany)

(Received December 30, 1992; accepted March 25, 1993)

Abstract


A one-pot procedure for the synthesis of 2-fluoro-3-trifluoromethyl-substituted thiophens **6** from 1-aryl-4,4-difluoro-3-trifluoromethylbut-3-en-1-one (**3**) and phosphorus pentasulfide is described. The fluorine atom at C-2 of compounds **6** undergoes nucleophilic displacement reactions.

The development of synthetic methodology for the regioselective introduction of short-chain perfluoroalkyl groups into organic molecules is of current interest [1, 2]. From the biomedical viewpoint, fluoro substitution often confers unique properties on a molecule, e.g. in terms of increasing lipophilicity, which in turn changes *in vivo* absorption and transport rates. Hence, biological activity is often enhanced [3].

Although numerous synthetic routes to thiophens have been reported [4], only very few routes to fluorine- and/or perfluoroalkyl-substituted thiophens have been described [5]. Here we report on a versatile new route to 3-trifluoromethylthiophens starting from hexafluoroacetone (**1**).

α,β -Unsaturated ketones **2** can easily be prepared from hexafluoroacetone (**1**) via procedures reported in the literature [6]. Heterodienes **2** can be transformed into the β,γ -unsaturated ketones **3** by [4 + 1] cycloaddition of tin(II) chloride followed by thermally-induced fluoride elimination [7]. Treatment of **3** with phosphorus pentasulfide at 120–140 °C results in an oxygen/sulfur exchange reaction (**3** → **5**). The higher nucleophilicity of sulfur compared with that of oxygen facilitates an intramolecular 1,5-cyclization. Elimination of HF from **5** results in aromatization to give the 2-fluoro-3-trifluoromethylthiophens **6**.

*Author to whom correspondence should be addressed.



$\text{R}^1 =$ (a) phenyl, (b) 4-methoxyphenyl, (c) 4-chlorophenyl, (d) 4-fluorophenyl, (e) 2-fluorophenyl

Scheme 1.

Compounds **6** were characterized by ^1H , ^{13}C and ^{19}F NMR data as well as by mass spectrometry and elemental analyses.

General procedure for the preparation of compounds 6 – Compound **3** (10 mmol) and 2.22 g (10 mmol) phosphorus pentasulfide were heated for 2–6 h at 120–140 °C. After cooling to room temperature, 20 ml of hexane was added to the reaction mixture which was then stirred for an additional 2 h. The reaction mixture was filtered, concentrated under reduced pressure and purified via column chromatography (silica gel 0.2–0.063 mm; eluent, hexane).

Compound 6a: Yield, 1.35 g, 55%, oil. Analysis: Calc. for $\text{C}_{11}\text{H}_6\text{F}_4\text{S}$: C, 53.66; H, 2.46%. Found: C, 53.52; H, 2.52%. ^{19}F NMR (Bruker AC 250, 235.3 MHz, CDCl_3) δ : –45.99 (dq, $J_{\text{FH}} = 3$ Hz, $J_{\text{FF}} = 12$ Hz, 1F, thiophen-F); 19.30 (d, $J_{\text{FF}} = 12$ Hz, 3F, CF_3) ppm. ^1H NMR (Bruker AM 360, 360.1 MHz, CDCl_3) δ : 6.98 (d, $J_{\text{HF}} = 3$ Hz, 1H, thiophen-H); 7.36 (m, 3H, phenyl-H); 7.44 (m, 2H, phenyl-H) ppm. ^{13}C NMR (Bruker 360, 90.6 MHz, CDCl_3) δ : 113.11 (dq, $J_{\text{CF}} = 4$ Hz, $J_{\text{CF}} = 36$ Hz, thiophen-C3); 116.40 (m, thiophen-C4); 120.94 (dq, $J_{\text{CF}} = 3$ Hz, $J_{\text{CF}} = 270$ Hz, CF_3); 125.52 (d, $J_{\text{CF}} = 1$ Hz); 128.53, 129.16, 132.42 (phenyl-C); 132.65 (d, $J_{\text{CF}} = 2$ Hz, thiophen-C5);

163.39 (dq, $J_{\text{CF}} = 301$ Hz, $J_{\text{CF}} = 3$ Hz, thiophen-C2) ppm. MS (m/z): 246 $[\text{M}]^+$; 227 $[\text{M} - \text{F}]^+$; 226 $[\text{M} - \text{HF}]^+$; 207 $[\text{M} - \text{HF}]^+$; 182 $[\text{M} - \text{CS}]^+$; 133 $[\text{M} - \text{CF}_3 - \text{CS}]^+$. IR (film) (cm^{-1}): 1600; 1520; 1490; 1450; 1410.

Because of the electron-withdrawing properties of the trifluoromethyl group, the fluorine atom at C-2 of compounds **6** is susceptible to nucleophilic displacement reactions with a broad range of nucleophiles [7]. Hence, the reaction sequence $1 \rightarrow 3 \rightarrow 6 \rightarrow 7$ offers a convenient method for the synthesis of 3-trifluoromethylthiophens with a variety of substitution patterns at C-2 and C-5. In particular, the possibility of introducing various side-chains with additional functional groups into ring position 2 in the final step of the reaction sequence ($6 \rightarrow 7$), in order to enhance and/or modify biological activity [8], makes this strategy versatile and valuable from a preparative viewpoint.

A typical example of the nucleophilic substitution process involves the reaction of **6a** with sodium 2-propoxide. The alkoxide was generated *in situ* from 0.08 ml (1 mmol) 2-propanol and 0.02 g (1 mmol) sodium in 10 ml dioxane. Compound **6a** (0.25 g, 1 mmol) was added and the mixture stirred at room temperature for 1 h. The mixture was then washed with 10 ml water and extracted three times with 10 ml ether. The residue obtained on evaporation of the solvent was purified by column chromatography [eluent, hexane/trichloromethane (1:1)].

Compound **7a** ($\text{R}^1 = \text{C}_6\text{H}_5$, $\text{Nuc} = \text{OCH}(\text{CH}_3)_2$): Yield, 0.27 g, 93%, m.p. 57 °C. Analysis: Calc. for $\text{C}_{14}\text{H}_{13}\text{F}_3\text{OS}$: C, 58.73; H, 4.58%. Found: C, 58.81; H, 4.58%. ^{19}F NMR (Bruker AC 250, 235.3 MHz, CDCl_3) δ : 19.73 (s, 3F, CF_3) ppm. ^1H NMR (Bruker WP 200, 200.1 MHz, CDCl_3) δ : 1.43 (d, $J_{\text{HH}} = 6$ Hz, 6H, $\text{CH}(\text{CH}_3)_2$); 4.44 (sept., $J_{\text{HH}} = 6$ Hz, 1H, $\text{OCH}(\text{CH}_3)_2$); 7.05 (s, 1H, thiophen-H); 7.32 (m, 3H, phenyl-H); 7.47 (m, 2H, phenyl-H) ppm. ^{13}C NMR (Bruker AC 250, 50.3 MHz, CDCl_3) δ : 21.97 ($\text{CH}(\text{CH}_3)_2$); 80.59 ($\text{OCH}(\text{CH}_3)_2$); 113.79 (q, $J_{\text{CF}} = 34$ Hz, thiophen-C3); 117.95 (q, $J_{\text{CF}} = 3$ Hz, thiophen-C4); 122.13 (q, $J_{\text{CF}} = 270$ Hz, CF_3); 125.04, 127.49, 128.96 (phenyl-C); 130.51 (thiophen-C5); 133.52

(phenyl-C); 163.36 (q, $J_{\text{CF}} = 3$ Hz, thiophen-C2) ppm. MS (m/z): 286 $[\text{M}]^+$; 244 $[\text{M} - \text{C}_3\text{H}_6]^+$; 243 $[\text{M} - \text{C}_3\text{H}_7]^+$; 225 $[\text{M} - \text{F}]^+$; 224 $[\text{M} - \text{F}]^+$; 196 $[\text{M} - \text{CO}]^+$; 121 $[\text{C}_6\text{H}_5\text{CS}]^+$; 77 $[\text{C}_6\text{H}_5]^+$; 43 $[\text{C}_3\text{H}_7]^+$; 42 $[\text{C}_3\text{H}_6]^+$. IR (KBr) (cm^{-1}): 1590; 1525; 1500; 1460; 1420; 1400.


The scope of the concept of transforming bis(trifluoromethyl)-substituted hetero-1,3-dienes into partially fluorinated five-membered heteroaromatic systems, and the synthetic potential of compounds **6**, will be described elsewhere.

Acknowledgement

We wish to thank Deutsche Forschungsgemeinschaft for financial support and Hoechst AG, Frankfurt, for a generous supply of chemicals.

References

- 1 D.J. Burton and Z.Y. Yang, *Tetrahedron*, **48** (1992) 189, and literature cited therein.
- 2 M.A. McClinton and D.A. McClinton, *Tetrahedron*, **48** (1992) 6555, and literature cited therein.
- 3 R. Filler and Y. Kobayashi, *Biomedical Aspects of Fluorine Chemistry*, Kodansha, Tokyo, 1982; Rhône Poulenc Conference: *Synthesis of Aromatic and Heteroaromatic Compounds Substituted by a Limited Number of Fluorine Atoms or Short-Chain Fluorinated Groups*, Lyon, September 1986; Abstracts: *L'Actualité Chim.*, (1987) 135; J.T. Welch, *Selective Fluorination in Organic and Bioorganic Chemistry*, ACS Symp. Ser. No. 456, Am. Chem. Soc., Washington, DC, 1991, and literature cited therein.
- 4 S. Gronowitz, in A. Weissberger and E.C. Taylor (eds.), *The Chemistry of Heterocyclic Compounds, Thiophene and its Derivatives*, Wiley, New York, 1985, Part I, pp. 1–125.
- 5 M.W. Briscoe, R.D. Chambers, S.J. Mullins, T. Nakamura and F.T. Drakesmith, *J. Chem. Soc., Chem. Commun.*, (1990) 1127; R.D. Chambers, C.G.P. Jones, M.J. Silvester and D.B. Speight, *J. Fluorine Chem.*, **25** (1984) 47; C.F. Krespan and B.C. McKusick, *J. Am. Chem. Soc.*, **83** (1961) 3438; J. Burdon, J.G. Campbell, I.W. Parsons and J.C. Tatlow, *J. Chem. Soc. C*, (1971) 352.
- 6 T. Ishihara, H. Shinjo, Y. Inoue and T. Ando, *J. Fluorine Chem.*, **22** (1983) 1; K. Burger and B. Helmreich, *J. Prakt. Chem.*, **334** (1992) 219.
- 7 For similar nucleophilic displacement reactions, see K. Burger and B. Helmreich, *J. Chem. Soc., Chem. Commun.*, (1992) 348; K. Burger, K. Geith and D. Hübl, *Synthesis*, (1988) 194, 199.
- 8 D. Hübl, M. Ganzer, F. Arndt and R. Rees, *Ger. Offen. DE 3 614 229* (1988); [*Chem. Abs.*, **109** (1988) 124415c]; K. Burger, E. Höss and K. Geith, *Synthesis*, (1990) 360, and literature cited therein.



$\text{Nuc}^- = \text{R}^2\text{O}^-, \text{R}^2\text{S}^-, \text{CN}^-, \text{H}^-, \text{Ph}^-$ $\text{NucH} = \text{R}^2_2\text{NH}$

Scheme 2.