

# The Juliá-Colonna Type Asymmetric Epoxidation Reaction Catalyzed by Soluble Oligo-L-leucines Containing an $\alpha$ -Aminoisobutyric Acid Residue: Importance of Helical Structure of the Catalyst on Asymmetric Induction

Ryukichi Takagi, Akiko Shiraki, Toshiki Manabe, Satoshi Kojima, and Katsuo Ohkata\*

Department of Chemistry, Graduate School of Science, Hiroshima University, 1-3-1 Kagamiyama, Higashi-Hiroshima 739-8526


(Received January 5, 2000; CL-000008)

Length-defined organic solvent soluble oligo-L-leucines containing an Aib residue were prepared by stepwise elongation and fragment condensation methods, and were used as catalysts in the Juliá-Colonna asymmetric epoxidation reaction. The yield and enantioselectivity rose by increasing the number of amino acid units in the catalyst. The enantioselectivity was very sensitive to the reaction solvent. The IR characteristic bands (the amide I region) in  $\text{CH}_2\text{Cl}_2$  indicated the soluble catalysts to be of helical structure in solution.

Epoxides are a widely used group of compounds in organic synthesis.<sup>1</sup> For this reason several approaches for enantioselective epoxidation have been developed.<sup>2</sup> Juliá-Colonna asymmetric epoxidation which utilizes poly-L-leucine as catalyst has been established as a highly selective method of epoxidizing enones<sup>2d,3</sup> and has found application in natural product synthesis.<sup>4</sup> Our interest in the chemistry of small ring compounds<sup>5</sup> has attracted us to this very intriguing reaction, of which mechanistically little has been uncovered. The primary problem concerning mechanistic examinations has been the insolubility of the catalyst in ordinary solvents. One advance has been the introduction of cross-linked aminomethylpolystyrene support.<sup>6</sup> This paved the way for preparation of polymer-bound length-defined catalysts, from which it was deduced that the chirality of the *N*-terminus dictates the stereochemistry of the product and that a certain number of consecutive amino acids of the same chirality at this end are required for reasonable levels of asymmetric induction.<sup>7</sup> Still here, however, the state of the catalysts has hampered spectroscopic examinations, and thus has complicated speculations about the mechanism of the asymmetric induction.


In their early work, Juliá and Colonna had already suggested that  $\alpha$ -helical structure might be important for asymmetric induction in the epoxidation reaction under tri- and biphasic conditions,<sup>8</sup> unfortunately they could not produce evidence to back it up. Our observation of IR spectra of length-defined oligo-L-leucines in the solid state agreed with their suggestion.<sup>9</sup> Based on the report that the presence of even only a single  $\alpha$ -aminoisobutyric acid (Aib) residue placed in the mid-section of oligo-L-amino acid chains promotes helix formation and greatly improves the solubility of the oligomers towards organic solvents,<sup>10</sup> we prepared several length-defined oligo-L-leucines containing an Aib residue, and have carried out the first examination of the Juliá-Colonna asymmetric epoxidation reaction with soluble catalysts. Herein we disclose our results.

The protected oligo-L-leucines ( $\text{Boc-L-Leu}_m\text{-Aib-L-Leu}_n\text{-OBzl}$ ) were prepared by the fragment condensation of  $\text{Boc-L-Leu}_m\text{-OH}$  and  $\text{TFA-H-Aib-L-Leu}_n\text{-OBzl}$  as shown in Scheme 2 by modifying reported procedures.<sup>10</sup> The oligo-L-leucines were



**Scheme 1.** Juliá-Colonna asymmetric epoxidation using soluble oligo-L-leucine catalysts.

fully characterized by MS (MALDI-TOF),  $^1\text{H}$  NMR and IR spectra. All the  $\text{Boc-L-Leu}_m\text{-Aib-L-Leu}_n\text{-OBzl}$  peptides prepared here were soluble in a variety of organic solvents.



**Scheme 2.** Preparation of  $\text{Boc-L-Leu}_m\text{-Aib-L-Leu}_n\text{-OBzl}$  and Free  $\text{NH}_2$  catalysts: NMM = *N*-methylmorpholine; HOBT = hydroxybenzotriazole; WSCI = 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride; TFA = trifluoroacetic acid.

Epoxidation reactions of chalcone with the urea-hydrogen peroxide adduct and DBU in the presence of soluble catalysts ( $\text{Boc-L-Leu}_m\text{-Aib-L-Leu}_n\text{-OBzl}$  and  $\text{TFA-H-L-Leu}_m\text{-Aib-L-Leu}_n\text{-OBzl}$  oligomers) were investigated at room temperature in organic solvents.<sup>11</sup> Table 1 shows the stereoselectivity in the epoxidation catalyzed by  $\text{Boc-L-Leu}_m\text{-Aib-L-Leu}_n\text{-OBzl}$  ( $m = 4, 6$  and  $n = 4, 6$ ) in THF and some *N*-deprotected oligomers having a free  $\text{NH}_2$  group. Both  $\text{Boc-L-Leu}_m\text{-Aib-L-Leu}_n\text{-OBzl}$  and  $\text{TFA-H-L-Leu}_m\text{-Aib-L-Leu}_n\text{-OBzl}$  oligomers were effective as catalysts from the viewpoint of both chemical yield and stereoselectivity. The general trend was that the longer catalyst gave better results, a tendency similarly observed in the epoxidation reaction under tri- or biphasic conditions.<sup>7,8,9</sup> Contrary to previous observations under heterogeneous tri- and biphasic conditions where protection of the *N*-terminus as amides<sup>8</sup> and carbamates<sup>9</sup> lead to significant deterioration in reactivity, our protected catalysts furnished comparable results with corresponding unprotected catalysts. Coupled with the fact that  $\text{NMe}_2$  (in the place  $\text{NH}_2$ ) is also compatible,<sup>7,8</sup> neither the hydrogen-bonding ability of the hydrogen atom attached to nitrogen nor the basicity of the nitrogen atom are involved in the stereo-determining step, and thus the role of the *N*-terminus seems to be minimal. The negative results with previous carbonyl protected catalysts<sup>8,9</sup> could be due to unfavorable catalyst conformations induced by the groups.

The solvent effect on the enantioselectivity is shown in Table 2. The stereoselectivity was very sensitive to the reaction solvent. The highest enantioselectivity was observed in THF. In  $\text{CHCl}_3$  solution, very low enantioselectivity was observed although the epoxide was obtained in comparable yield. Since

**Table 1.** Boc-L-Leu<sub>m</sub>-Aib-L-Leu<sub>n</sub>-OBzl and TFA·H-L-Leu<sub>m</sub>-Aib-L-Leu<sub>n</sub>-OBzl catalyzed epoxidation in THF

| Entry | Catalyst                                                          | yield /% | % ee <sup>a</sup> |
|-------|-------------------------------------------------------------------|----------|-------------------|
| 1     | Boc-L-Leu <sub>4</sub> -Aib-L-Leu <sub>4</sub> -OBzl <sup>b</sup> | 50       | 61                |
| 2     | TFA·H-L-Leu <sub>4</sub> -Aib-L-Leu <sub>4</sub> -OBzl            | 61       | 68                |
| 3     | Boc-L-Leu <sub>4</sub> -Aib-L-Leu <sub>6</sub> -OBzl              | 60       | 78                |
| 4     | TFA·H-L-Leu <sub>4</sub> -Aib-L-Leu <sub>6</sub> -OBzl            | 54       | 73                |
| 5     | Boc-L-Leu <sub>6</sub> -Aib-L-Leu <sub>4</sub> -OBzl              | 89       | 85                |
| 6     | Boc-L-Leu <sub>6</sub> -Aib-L-Leu <sub>6</sub> -OBzl              | 73       | 94                |

<sup>a</sup> Estimated by HPLC using a chiral OD column. <sup>b</sup> Known compound.**Table 2.** Solvent effect in Boc-L-Leu<sub>6</sub>-Aib-L-Leu<sub>6</sub>-OBzl catalyzed epoxidation

| Entry | Solvent                         | yield /% | % ee <sup>a</sup> |
|-------|---------------------------------|----------|-------------------|
| 1     | THF                             | 73       | 94                |
| 2     | Toluene                         | 47       | 84                |
| 3     | CH <sub>2</sub> Cl <sub>2</sub> | 41       | 76                |
| 4     | CHCl <sub>3</sub>               | 72       | 15                |

<sup>a</sup> Estimated by HPLC using a chiral OD column

IR measurements indicated the homogeneous catalyst to be helical even in CHCl<sub>3</sub>, this experimental result can be ascribed to the presence of a relatively fast background uncatalyzed epoxidation process.<sup>12</sup>

Examination of the IR spectra of Boc-L-Leu<sub>6</sub>-Aib-L-Leu<sub>6</sub>-OBzl, which exhibited the highest selectivity, showed a weak band at 3424 cm<sup>-1</sup> (w, free N-H stretching), and strong bands at 3324 (s, hydrogen bonding N-H stretching) and 1661 cm<sup>-1</sup> (s, C=O stretching) in CH<sub>2</sub>Cl<sub>2</sub> solution, which can be assigned to helical structure.<sup>13</sup>

We have demonstrated that soluble oligo-L-leucine catalysts that show a high degree of helical conformational structure give results comparable to those of insoluble catalysts. Therefore, we draw the conclusion that in general the segment of the catalysts in the Juliá-Colonna reaction involved in the asymmetric induction process assumes helical conformation. This improvement in solubility of the catalyst brings in a new dimension to the Juliá-Colonna reaction and should lead to further understanding of the reaction.

We are grateful to Professor Katsuyoshi Yoshizato and Mr. Dan Kristensen, Hiroshima University, for the use of their MALDI mass spectrometer, and Professor Kenichi Ohno of Hiroshima University for IR spectra measurements. NMR, MS, and CD spectra were measured at the Instrument Center for Chemical Analysis, Hiroshima University.

## References and Notes

- a) M. Bartok and K. L. Lang, "The Chemistry of Ethers, Crown Ethers, Hydroxyl Groups and Their Sulphur Analogs, Supplement E," ed by S. Patai, John Wiley, New York (1980) vol. 2, p. 609. b) A. S. Rao, S. K. Paknikar, and J. G. Kirtane, *Tetrahedron*, **39**, 2323 (1983). c) J. G. Smith, *Synthesis*, **1984**, 629. d) E. G. Lewars, in "Comprehensive Heterocyclic Chemistry," ed by A. R. Katritzky and C. W. Rees, Pergamon, Oxford (1984), vol. 7, p. 95. e) I. Erden, in "Comprehensive Heterocyclic Chemistry II," ed by A. Padwa, Elsevier, Oxford (1996), vol. 1A, p. 97.
- a) T. Katsuki and K. B. Sharpless, *J. Am. Chem. Soc.*, **102**, 5874 (1980). b) W. Zhang, J. L. Loebach, S. R. Wilson, and E. N. Jacobsen, *J. Am. Chem. Soc.*, **112**, 2801 (1990). c) R. Irie, K. Noda, Y. Ito, N. Matsumoto, and T. Katsuki, *Tetrahedron Lett.*, **31**, 7345 (1990). d) S. Juliá, J. Masana, and J. C. Vega, *Angew. Chem., Int. Ed. Engl.*, **19**, 929 (1980). e) D. Yang, Y.-C. Yip, M.-W. Tang, M.-K. Wong, J.-H. Zheng, and K.-K. Cheung, *J. Am. Chem. Soc.*, **118**, 491 (1996). f) Y. Tu, Z.-X. Wang, and Y. Shi, *J. Am. Chem. Soc.*, **118**, 9806 (1996). g) D. Enders, J. Zhu, and G. Raabe, *Angew. Chem., Int. Ed. Engl.*, **35**, 1725 (1996). h) M. Bougauchi, S. Watanabe, T. Arai, H. Sasaki, and M. Shibasaki, *J. Am. Chem. Soc.*, **119**, 2329 (1997). i) C. L. Elston, R. F. W. Jackson, S. J. F. MacDonald, and P. J. Murray, *Angew. Chem., Int. Ed. Engl.*, **36**, 410 (1997). j) B. Lygo and P. G. Wainwright, *Tetrahedron Lett.*, **39**, 1599 (1998). k) S. Watanabe, T. Arai, H. Sasaki, M. Bougauchi, and M. Shibasaki, *J. Org. Chem.*, **63**, 8090 (1998).
- For reviews on Juliá-Colonna epoxidation see: a) M. E. Lasterra-Sanchez and S. M. Roberts, *Curr. Org. Chem.*, **1**, 187 (1997). b) S. Ebrahim and M. Wills, *Tetrahedron: Asymmetry*, **8**, 3163 (1997). c) L. Pu, *Tetrahedron: Asymmetry*, **9**, 1457 (1998).
- a) J. R. Flisak, K. J. Gombatz, M. M. Holmes, A. A. Jamas, I. Lantos, W. L. Mendelson, V. J. Navack, J. J. Remich, and L. Snyder, *J. Org. Chem.*, **58**, 6247 (1993). b) R. J. J. Nel, P. S. van Heerden, H. van Rensburg, and D. Ferreira *Tetrahedron Lett.*, **39**, 5623 (1998). c) B. M. Adger, J. V. Barkley, S. Bergeron, M. W. Cappi, B. E. Flowerdew, M. P. Jackson, R. McCague, T. C. Nugent, and S. M. Roberts, *J. Chem. Soc., Perkin Trans. 1*, **1997**, 3501. d) M. W. Cappi, W.-P. Chen, R. W. Flood, Y.-W. Liao, S. M. Roberts, J. Skidmore, J. A. Smith, and N. M. Williamson, *Chem. Commun.*, **1998**, 1159. e) W.-p. Chen and S. M. Roberts, *J. Chem. Soc., Perkin Trans. 1*, **1999**, 1043.
- a) K. Ohkata, J. Kimura, Y. Shinohara, R. Takagi, and Y. Hiraga, *Chem. Commun.*, **1996**, 2411. b) R. Takagi, J. Kimura, Y. Shinohara, Y. Ohba, K. Takezono, Y. Hiraga, S. Kojima, and K. Ohkata, *J. Chem. Soc., Perkin Trans. 1*, **1998**, 689.
- S. Itsuno, M. Sakakura, and K. Ito, *J. Org. Chem.*, **53**, 6047 (1990).
- P. A. Bentley, M. W. Cappi, R. W. Flood, S. M. Robert, and J. A. Smith, *Tetrahedron Lett.*, **39**, 9297 (1998).
- a) S. Juliá, J. Gulixer, J. Masana, J. Rocas, S. Colonna, R. Annuziata, and H. Molinari, *J. Chem. Soc., Perkin Trans. 1*, **1982**, 1317. b) S. Colonna, H. Molinari, S. Banfi, S. Juliá, J. Masana, and A. Alvarez, *Tetrahedron*, **39**, 1635 (1983). c) S. Banfi, S. Colonna, H. Molinari, S. Juliá, and J. Guixer, *Tetrahedron*, **40**, 5207 (1984).
- Manuscript in preparation.
- M. Narita, K. Ishikawa, H. Sugawara, and M. Doi, *Bull. Chem. Soc. Jpn.*, **58**, 1731 (1985).
- General method: To a mixture of chalcone 25 mg (0.12 mmol), catalyst 0.030 mmol, urea-hydrogen peroxide adduct 12.5 mg (0.16 mmol) and 1 ml of THF, 0.1 ml of DBU (0.67 mmol) was added at 0 °C. The mixture was allowed to warm to room temperature and stirred for 24 h. The resulting mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The combined extracts were washed with aqueous Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>, water and brine, dried over MgSO<sub>4</sub> and evaporated. The crude product was purified by preparative TLC [silica gel, petroleum ether : ether = 10 : 1 (v/v)]. As the reaction progresses, the turbidity caused by the lowly soluble urea-hydrogen peroxide adduct decreases.
- Uncatalyzed reactions in THF and CHCl<sub>3</sub> solutions: By the same procedure except without catalyst, the racemic epoxide was furnished in 26 and 68% yield, respectively.
- a) T. Miyazawa and E. J. Blout, *J. Am. Chem. Soc.*, **83**, 712 (1961). b) D. F. Kennedy, M. Crisma, C. Toniolo, and D. Chapman, *Biochemistry*, **30**, 6541 (1991).
- The CD spectrum of a trifluoroethanol solution of Boc-L-Leu<sub>6</sub>-Aib-L-Leu<sub>6</sub>-OBzl in the absorption region of oligomer chromophores displayed a negative CD band at 203 nm along with a shoulder centered near 222 nm with an R = [Θ]<sub>222</sub>/[Θ]<sub>203</sub> ratio of 0.5. In addition, the peak at 190 nm was positive with a further negative maximum at 181 nm. These spectral characteristics are extremely similar to those of a reported oligopeptide with established right-handed 3<sub>10</sub>-helical conformation, thus implying the presence of this structure: a) M. Manning and R. W. Woody, *Biopolymers*, **31**, 569 (1991). b) C. Toniolo, A. Polese, F. Formaggio, M. Crisma, and J. Kamphuis, *J. Am. Chem. Soc.*, **118**, 2744 (1996).