Communications

Organocatalytic Cyclopropanation



Organic-Catalyst-Mediated Cyclopropanation Reaction**

Charles D. Papageorgiou, Steven V. Ley, and Matthew J. Gaunt*

Cyclopropane units are a fundamental class of functional group that are the focus of many organic synthesis programs and perform a key structural role in a range of bioactive natural and nonnatural molecules.^[1] The importance of cyclopropanes is reflected in the enormous effort that has been invested in their diastereo- and enantioselective synthesis.^[1,2]

Recently, a strong emphasis has been placed on the development of catalytic methods to generate cyclopropanes. While there have been significant advances in the development of catalytic versions of the Simmons–Smith and carbene transfer reactions, there are few reports of catalytic methods based on the reaction of ylides with electron-deficient alkenes.^[2] The use of ylides as cyclopropanation reagents dates back to work by Corey et al.^[3] Of the many developments that have since been reported, the contributions from the groups of Hanessian,^[4] Dai and Tang,^[5] and Aggarwal^[6] deserve special mention. In these processes, however, the ylide precursor is usually generated in a separate step and there is only one reported process that is catalytic in the ylide species.^[6,7]

We were interested in utilizing a nitrogen ylide to effect a cyclopropanation reaction. Surprisingly, there are relatively few examples that employ nitrogen-derived ylides.^[8,9] An ammonium ylide based reaction becomes an attractive target for a general diastereo- and enantioselective cyclopropanation process owing to the vast range of tertiary amines that are commercially available. In this communication we describe the development of a stoichiometric "one-pot" ammonium ylide based cyclopropanation process and our initial studies towards an enantioselective reaction. We also report the development of the first catalytic cyclopropanation reaction involving ammonium ylides where the catalyst is a tertiary amine.

We began our study by investigating the reactivity of preformed quaternary ammonium salts such as **1** as nitrogen ylide precursors. Treatment of salt **1** with base followed by

[*] Dr. M. J. Gaunt, C. D. Papageorgiou, Prof. Dr. S. V. Ley
University Chemical Laboratory
Lensfield Road, Cambridge, CB2 1EW (UK)
Fax: (+44) 1223-336-442
E-mail: mjg32@cam.ac.uk

[**] We gratefully acknowledge financial support from the EPSRC and Pfizer Global Research and Development (Sandwich, UK) for Industrial Case Award (to C.D.P.), the Novartis Research Fellowship (to S.V.L.), and the British Ramsay Memorial Trust and Magdalene College for Research Fellowship (to M.J.G.). We also thank Dr. A. J. Morrison for helpful advice and discussion.

Supporting information for this article is available on the WWW under http://www.angewandte.org or from the author.

828

Table 1: Optimization of reaction conditions.^[a]

	Ph The Ph				
	1	2a	3a		
Solvent	Base	<i>t</i> [h]	Yield ^[b]	d.r. (<i>trans:cis</i>) ^[c]	
MeCN	NaOH	6	96 %	> 95:5	
THF	NaOH	3.5	66%	15:1	
DCE	NaOH	19	73%	>95:5	
DMSO	NaOH	1.5	54%	>95:5	
$CH_2Cl_2^{[d]}$	NaOH	4.5	65%	> 95:5	
MeCN	КОН	4.5	68%	15:1	
MeCN	Na ₂ CO ₃	25	79 %	>95:5	
MeCN	DBU	16	87 %	>95:5	
MeCN	DABCO ^[e]	24	5%	> 95:5	
MeCN	Et₃N	24	10%	>95:5	

[a] Salt 1, acrylate 2a (1 equiv), base (2 equiv), solvent (0.25 m), 80 °C (oil bath temperature). [b] Yield of isolated product after chromatography. [c] Determined by ¹H NMR spectroscopy. [d] Reaction stirred at 45 °C. [e] 2.5 equiv of DABCO was used. DABCO = 1,4-diazobicyclo[2.2.2]octane, DCE = 1,2-dichloroethane, DBU = 1,8-diazabicyclo[5.4.0]undec-7ene.

tert-butyl acrylate (2a) formed cyclopropane 3a in varying yields depending on the conditions employed; the best results are highlighted in Table 1. Of crucial importance was the temperature, with little or no reaction observed at room temperature. However, at 80 °C the reaction produced the highest yield of *trans*-disubstituted cyclopropane. It is important to note that the corresponding reaction between phenacyl chloride (instead of salt 1) and acrylate 2a does not form the cyclopropane.^[10]

To increase the efficiency of this process we investigated the development of a stoichiometric "one-pot" cyclopropanation reaction, in which the ammonium salt and hence the ylide could be generated in situ from readily available starting materials. Accordingly, DABCO (5) was added to phenacyl chloride (4) in acetonitrile followed by the base and alkene 2a, and the reaction mixture was stirred at 80 °C. After 18 h, cyclopropane 3a was isolated in 79% yield after chromatography as a single diastereoisomer (Scheme 1).^[11] It is note-

$$\begin{array}{c} 0 \\ Ph \overset{O}{\underbrace{}} \\ 4 \end{array} \overset{O}{\underbrace{}} \\ \begin{array}{c} 1 \text{ equiv } N \\ \hline \\ 2a, \text{ NaOH, MeCN, 80 °C} \\ 79 \% \end{array} \overset{O}{\underbrace{}} \\ \begin{array}{c} Ph \overset{O}{\underbrace{}} \\ Ph \\ \hline \\ 3a \\ O \end{array} \overset{O}{\underbrace{}} \\ \begin{array}{c} 0 \\ O \\ Bu \\ \hline \\ 3a \\ O \end{array} \overset{O}{\underbrace{}} \\ \end{array} \end{array}$$

Scheme 1. One-pot cyclopropanation.

worthy that the reaction proceeds without the formation of byproducts, and simple aqueous workup affords pure cyclopropane (¹H NMR spectroscopy, LCMS) that does not necessarily require further purification.

A stoichiometric "one-pot" cyclopropanation reaction offers significant advantages over other similar methods as it precludes the necessity to generate and isolate the ylide precursor (1) in a separate step. The general utility of this

they are a potential source of chirality for an enantioselective cyclo-

propanation reaction. The chiral

quaternary ammonium salt 11 was readily formed in situ from the

addition of phenacyl bromide (8)

to amine 9.^[14] Subsequent addition of acrylate 2a and sodium hydroxide (2 equiv), and stirring the reac-

tion at 80°C for 24 h generated

cyclopropane (+)-**3a** in 61% yield (Table 3). The enantiomeric ratio

(e.r.) from this initial reaction was

91.5:8.5 (83 % *ee*), as determined by chiral HPLC.^[15] While this result

displayed excellent enantioselectivity we were concerned for the

enantiomeric integrity of the cyclo-

propane in the reaction mixture. By

monitoring the e.r. over the course of the reaction, we observed an

erosion in the enantiomeric purity

of the cyclopropane (Table 3, entry 1). However, reducing the

amount of base in the reaction to

1.3 equivalents generated (+)-3a in

57% yield with an e.r. of 97:3 (94% *ee*). Furthermore, using dihy-

drocinchonine derivative 10 under our modified reaction conditions formed the opposite enantiomer, (-)-3a, in 58% yield and an e.r. of

97:3 (94% ee). To the best of our

knowledge these are the first exam-

ples of an enantioselective cyclo-

propanation reaction based on a

chiral ammonium ylide. We are

currently screening the range of alkenes and other chiral tertiary

Table 2: Scope of reaction.



[a] α-Chlorocarbonyl, DABCO (1.0 equiv), MeCN (0.25 м), room temperature, 30 min, then alkene (1 equiv), base (1.5 equiv), 80 °C. [b] Yield of isolated product after chromatography. [c] Determined by ¹H NMR spectroscopy. [d] Preformed salt **1** was used in this reaction.

process was demonstrated across a range of substrates which established the efficacy of this new and versatile cyclopropane methodology (Table 2). The reaction worked well for a range of alkenes, with acrylates **2a** and **2b**, enone **2c**, acrylamide **2d**, and vinyl sulfone **2e** all forming only the *trans*-substituted 1,2cyclopropanes **3a–3e** in good to excellent yield.^[12] Acrylonitrile (**2f**) gave a mixture of cyclopropanes (**3f**; 2.3:1, *trans:cis*) in good yield, although some polymerization was observed. Acrolein (**2g**) also gave a separable mixture of cyclopropanes (**3g**; 4.3:1, *trans:cis*) in 40% yield when the preformed salt **1** was used. Ammonium ylide species bearing other functional groups were also investigated and produced cyclopropanes in good yield (entries 8 and 9, Table 2). Ketone **7** formed spiro[2.4]heptane derivative **3i** as a single diastereomer when reacted with 3 equivalents of **2c**.

The simplicity of the one-pot ammonium ylide approach to cyclopropanation should allow the use of any tertiary amine.^[13] DABCO (**5**) was chosen as a model tertiary amine to mimic the core structure of the cinchona alkaloids because Table 3: One-pot enantioselective cyclopropanation.

O Br		R ₃ N (1 equiv), SCO ₂ tBu 2a			Î. N	
	Ph 1 equiv)	NaOH, MeCN	l, 80 °C, 24	h Pł	√ ~/ `C 3a	O ₂ tBu
Ent	ry ^[a] Chiral amine	equiv NaOH	<i>t</i> [h]	$Yield^{[b]}$	e.r. (<i>ee</i>)	[c]
1	9	2	8	10%	97:3 (9 [,]	4) (+)
			16	46%	92:8 (8	4) (+)
			24	61%	91.5:8.5	5 (83) (+)
2	9	1.3	24	57%	97:3 (9-	4) (+)
3	10	1.3	24	58%	97:3 (9	4) (-)
				7	11	
			OMe H		OMe H-	
Ň	l _≫ ∕″ 9		10		Ň,	

Communications



Scheme 2. Proposed catalytic cycle for cyclopropanation.

amines to investigate the scope of this new enantioselective process.

With an efficient stoichiometric one-pot cyclopropanation reaction in hand, we next investigated a catalytic process. Examination of a possible reaction mechanism suggests that the amine 12 should be released at the end of the reaction when the cyclopropane ring forms (Scheme 2).^[16] Thus, in principle it should be possible to use the tertiary amine in catalytic quantities without fear of a competing background reaction. Accordingly, stirring a mixture of phenacyl chloride (4), 2c (3 equiv),^[17] sodium carbonate (1.2 equiv) and DAB-CO (0.2 equiv) in acetonitrile at 80°C for 24 h produced cyclopropane 3c in 82% yield. tert-Butyl acrylate (2a) and vinyl sulfone 2e also produce cyclopropanes 3a and 3e, respectively, under catalytic conditions (Table 4). In the absence of DABCO little or no reaction takes place which proves that the tertiary amine is required as a catalyst. We are currently exploring the scope of the reaction to develop a general catalytic cyclopropanation reaction based on ammonium vlides.

In summary, we have developed a practical and general "one-pot" cyclopropanation process, mediated by a tertiary

Table 4: Initial results for a catalytic cyclopropanation.^[a]



[a] Phenacyl chloride, DABCO (0.2 equiv), alkene (1.5 equiv), Na₂CO₃ (1.2 equiv), MeCN (0.25 M), 80 °C, 24 h. [b] Yield of isolated product after chromatography. [c] Determined by ¹H NMR spectroscopy.
 [d] 3 equiv Of 2c was used.

amine via an ammonium ylide intermediate, by using readily available starting materials. We have demonstrated that this reaction is both diastereo- and enantioselective and that the reaction can be made catalytic.^[18] We are currently exploring the scope of an asymmetric process with the goal of developing a catalytic enantioselective cyclopropanation reaction. Furthermore, application of this methodology to an intramolecular cyclopropanation system and to the synthesis of small-ring heterocycles is also being explored.

Received: October 14, 2002 [Z50353]

- For reviews, see: a) J. Salaun, *Chem. Rev.* **1989**, *89*, 1247; b) A.-H. Li, L.-X. Dai, V. K. Aggarwal, *Chem. Rev.* **1997**, *97*, 2341;
 c) A. Pfaltz in *Comprehensive Asymmetric Catalysis II* (Eds.: E. N. Jacobsen, A. Pfaltz, H. Yammamoto), Springer, Berlin, **1999**, pp. 513–538.
- [2] For recent developments, see: W. A. Donaldson, *Tetrahedron* 2001, 57, 8589, and references therein. For other examples see ref. [1].
- [3] E. J. Corey, M. Chaykovsky, J. Am. Chem. Soc. 1965, 87, 1353;
 E. J. Corey, M. Jautelat, J. Am. Chem. Soc. 1967, 89, 3912.
- [4] S. Hanessian, D. Andreotti, A. Gomtsyan, J. Am. Chem. Soc. 1995, 117, 10393.
- [5] S. Ye, Z.-Z. Huang, C.-A. Xia, Y. Tang, L.-X. Dai, J. Am. Chem. Soc. 2002, 124, 2432.
- [6] V. K. Aggarwal, A. Thompson, R. V. H. Jones, *Chem. Commun.* 1997, 1785 and ref. [7]. See also A. Solladie-Cavallo, A. Diep-Vohuule, T. Isarno, *Angew. Chem.* 1998, 110, 1824; *Angew. Chem. Int. Ed.* 1998, 37, 1689.
- [7] a) V. K. Aggarwal, E. Alonso, G. Fang, M. Ferrara, G. Hynd, M. Porcelloni, *Angew. Chem.* 2001, *113*, 1482; *Angew. Chem. Int. Ed.* 2001, *40*, 1433; b) V. K. Aggarwal, H. W. Smith, G. Hynd, R. V. H. Jones, R. Fieldhouse, S. E. Spey, *J. Chem. Soc. Perkin Trans. 1* 2000, 3267.
- [8] I. Zugravescu, M. Petrovanu, *N-Ylid Chemistry*, McGraw-Hill, New York, 1976.
- [9] a) S. S. Bhattacarjee, H. Ila, H. Junjappa, Synthesis 1982, 301; b) A. Jonczyk, A. Konarska, Synlett 1999, 1085; for other examples, see c) N. H. Vo, C. J. Eyermann, C. N. Hodge, *Tetrahedron Lett.* 1997, 38, 7951; d) A. M. Shestopalov, V. P. Litvinov, L. A. Rodinovskaya, Y. A. Sharanin, Zh. Org. Khim. 1991, 27, 146.
- [10] When Na₂CO₃ is used as the base, no reaction is observed. When NaOH is used, decomposition is
 - observed. [11] All compounds were characterized by ¹H and ¹³C NMR spectroscopy, high-resolution mass spectrometry and infrared spectroscopy.
 - [12] trans-Stereochemistry was confirmed in 3a by hydrolysis of the tert-butyl ester and comparison with an authentic sample, and in 3e by X-ray crystal-structure analysis.
 - [13] We also found that using quinuclidine in place of DABCO produced 3a in similar yield.
 - [14] Phenacyl chloride did not react with the more hindered chiral amine.
 - [15] See supporting information for details chiral HPLC analysis.
 - [16] No cyclopropane formation was observed when the reaction was

© 2003 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim 1433-7851/03/4207-0830 \$ 20.00+.50/0

830

Angew. Chem. Int. Ed. 2003, 42, No. 7



carried out with styrene instead of the electron-deficient alkenes which suggests that the reaction does not proceed through a carbene mechanism.

- [17] In the case of methyl vinyl ketone, 3.0 equiv was used because of its volatility. In the other cases only 1.5 equiv of the alkene was used.
- [18] For an overview of organocatalysis, see: P. I. Dalko, L. Moisan, Angew. Chem. 2001, 113, 3840; Angew. Chem. Int. Ed. 2001, 40, 3726.



Principles and Practice of Heterogeneous Catalysis

J. M. THOMAS, The Master's Lodge, Peterhouse, Cambridge, UK; W. J. THOMAS, University of Bath, UK

3-527-29239-X 1996 XIX, 669 pp 375 figures 25 tables. SC. € 55.00*/£32.50

Catalysis occupies a pivotal position in the physical and biological sciences. As well as being the mainstay of the chemical industry, it is the means of effecting many laboratory syntheses and the root cause of all enzymatic processes.

This book is an eminently readable introduction to the fundamental

principles of heterogeneous catalysis. Written by worldrenowned experts, it explains the vocabulary, grammar and literature of catalysis from the laboratoryoriented model study through to the operating plant.

Didactically skillful and using many lucidly designed figures, the authors present an insightful exposition of all important concepts, new developments and techniques in this rapidly advancing field.

WILEY-VCH

* The €-prices refer to Germany only!

73901111 v