


Transformation of Primary Carboxamides to N-(*t*-Butoxycarbonyl)amines Using CuBr₂-LiOBu^{*t*}

Jun-ichi YAMAGUCHI, Kensuke HOSHI, and Takeshi TAKEDA*

Department of Applied Chemistry, Tokyo University of Agriculture and Technology, Koganei, Tokyo 184

The reaction of primary carboxamides with the copper(II) reagent prepared from copper(II) bromide and lithium *t*-butoxide gave the corresponding N-(*t*-butoxycarbonyl)amines in good to high yields.

Recently we reported the oxidations of alcohols¹⁾ and amines²⁾ with CuBr₂-LiOBu^{*t*}, in which bromo(*t*-butoxy)copper(II) is assumed to serve as an active species. These oxidations are formally regarded as removal of hydrogens from adjacent two atoms. The alternative mode of dehydrogenation which would be promoted by the copper(II) oxidizing agent is 1,1-elimination of hydrogens. Then we have examined such oxidations, and describe here the transformation of primary carboxamides (1) to N-(*t*-butoxycarbonyl)amines (BOC-amines) (2) (Eq. 1).



When benzamide (1a) was treated with 2.2 equiv. of CuBr₂-LiOBu^{*t*}(3), the BOC-amine 2a was produced in 36% yield (run 1). It was found that 2a was obtained in good yields by the use of more than 3 equiv. of the oxidizing agent 3 (runs 2 and 3). On the basis of these results, we speculated that the present transformation would consist of the formation and Curtius rearrangement of nitrene intermediate 5 and the addition of bromo(*t*-butoxy)copper(II) (4) to isocyanate 6 (Eq. 2). In fact, 2a was produced in 84% yield by the reaction of phenyl isocyanate with 3 (1.1 equiv.) in THF at room temperature, whereas the treatment with *t*-butanol under the similar reaction conditions gave no addition product.

As shown in Table 1, various BOC-amines (2) were obtained in good to high yields by the reaction of primary carboxamides (1). In the case of the reaction of 3-phenylpropionamide (1g), however, no desired product was obtained, but the formation of insoluble materials was observed presumably due to the polymerization of 2-phenylethyl isocyanate (6g). In order to accelerate the addition of 4 to 6g, the use of a large excess of 3 and higher reaction temperature was examined. Although the reaction carried out at 45 °C gave the BOC-amine 2g in 20% yield along with *t*-butyl 3-phenylpropionate (7g) (run 10), 7g was preferentially produced under reflux (run 11).

A typical experimental procedure is as follows: To a THF (4 ml) solution of lithium *t*-butoxide (2.1 mmol) was added copper(II) bromide (491 mg, 2.1 mmol) at room temperature under argon, and the mixture was stirred for 15 min. Then a THF (2 ml) solution of cyclohexanecarboxamide (64 mg, 0.5 mmol) was added to the reaction mixture. After being stirred for 4 h, the reaction was quenched by addition of 3.5% aqueous solution of NH₃. The organic materials were extracted with CH₂Cl₂, and the extract was dried over Na₂SO₄. After removal

(Received May 6, 1993)