with the Mukaiyama reagent.<sup>13</sup> Irradiation at 350 nm for 18 h of a solution of the glycal 1a and 5 equiv of dibenzyl azodi-carboxylate (DBAD)<sup>14</sup> in cyclohexane gave a single [4 + 2]cycloadduct 2a (TLC, <sup>1</sup>H NMR, HPLC) in 73% yield (Scheme I) (for a typical procedure see ref 15). Glycals 1b, 6, 10, and 14 could be converted into their corresponding cycloadducts (2b, 7, 11, 15) in similar yields (Scheme I).

The diastereofacial selectivity of the cycloaddition is apparently controlled by the stereochemistry at C-3 and was assigned on the basis of NOE studies and by conversion of the adducts 2a, 2b, 7, and 15 into known compounds. That the product was a [4 +2] cycloadduct vs. a [2 + 2] was confirmed by the presence of the characteristic C=N band of dihydrooxadiazines in the IR spectra ( $\sim 1670 \text{ cm}^{-1}$ ).<sup>9</sup>

Treatment of the cycloadducts with a catalytic amount of *p*-toluene sulfonic acid in methanol gave the corresponding methyl glycosides (3a, 3b, 8, 12, 16) in quantitative yield. The opening of the cycloadducts by methanol occurred exclusively with inversion at C-1.17 Hydrogenolysis of the protected hydrazines gave the 2-amino glycosides (4, 9, 13, 17) in high yields (Scheme I). Manipulation of the protecting groups of the amines 4, 9, and 17 gave cited compounds which possessed identical characteristics (<sup>1</sup>H NMR, mp,  $[\alpha]_D$ ) with those reported in the literature, <sup>17-22</sup> thus confirming the stereochemical assignments made on the basis of NOE data. The amino furanoside 13 is a derivative of the hereto unreported methyl 2-amino-2-deoxy- $\alpha$ -D-idofuranoside.

There are several features of the transformations described above that bear comment. Substitution of water for methanol in opening of the cycloadduct would yield the lactol which could be further transformed via its aldehyde and/or hydroxyl moieties. Also noteworthy is that the nitrogen functionality is obtained directly in a protected form obviating the need for protection if one wishes to use the product as a synthetic intermediate.

In summary, 2-deoxy-2-amino glycosides can be prepared simply and in high yield by the cycloaddition of DBAD on the appropriate glycal. Due to the simplicity and mildness of the reaction conditions this reaction should be readily extendable to non-carbohydrate cases. Such studies are currently under way and will be reported in due course.

Acknowledgment. We thank Professor S. Danishefsky for helpful suggestions in the preparation of this manuscript.

Supplementary Material Available: Analytical and spectral data for 4, 7-9, 13, and 18 (4 pages). Ordering information given on any current masthead page.

(16) All new compounds cited in this article gave satisfactory elemental analysis

(17) Similar results were observed with the equivalent phenyloxazoline; see: Jacquinet, J. C.; Sinay, P. *Carbohydr. Res.* **1974**, *32*, 101–114. (18) Methyl 2-amino-2-deoxy-α-D-arabinofuranoside (**4**); mp 78–81 °C

(thy lacetate); iti. 75-77 °C (thyl acctate),  $[\alpha]^{22}_{D}$  +135° (c 0.4, methanol); iti.<sup>20</sup>  $[\alpha]_{D}$  +128° (c 1.0, methanol).

(19) Montgomery, J. A.; Thorpe, M. C.; Clayton, S. D.; Thomas, H. J. Carbohydr. Res. 1974, 32, 404-407

(20) Buchanan, J. G.; Clark, D. R. Carbohydr. Res. 1977, 57, 85-96. (21) Methyl 2-amino-2-deoxy-5,6-O-(1-methylethylidene)-B-D-glucofuranoside (18) (supplementary material) as a white solid; mp 60 °C (ether),  $[\alpha]^{22}_{D} - 58.8^{\circ}$  (c 0.4, methanol); lit.<sup>17</sup> mp 61-62 °C (ether),  $[\alpha]^{20}_{D} - 60.5^{\circ}$  (c

## Binuclear Platinum(II) Photochemistry. Rates of Hydrogen Atom Transfer from Organometallic Hydrides to Electronically Excited Pt<sub>2</sub>(P<sub>2</sub>O<sub>5</sub>H<sub>2</sub>)<sub>4</sub><sup>4-</sup>

Antonin Vlček, Jr.,\*1 and Harry B. Gray\*

Contribution No. 7487, Arthur Amos Noyes Laboratory California Institute of Technology Pasadena, California 91125 Received September 22, 1986

Previous investigations have established that the triplet  $d\sigma^* p\sigma$ excited states of  $d^8-d^8$  binuclear complexes ( ${}^3M_2^*$ , M = Rh, Ir, Pt) are readily oxidized by one-electron acceptors.<sup>2-5</sup> A related but largely unexplored reaction for these  ${}^{3}M_{2}^{*}$  states is hydrogen atom abstraction,  $RH + {}^{3}M_{2}^{*} \rightarrow R^{*} + {}^{*}M_{2}H$ , a step that was proposed by Roundhill<sup>6</sup> to be a primary process in the Pt<sub>2</sub>photocatalyzed conversion of isopropyl alcohol to acetone and dihydrogen (Pt<sub>2</sub> = Pt<sub>2</sub>(P<sub>2</sub>O<sub>5</sub>H<sub>2</sub>)<sub>4</sub><sup>4-</sup>). Support for Roundhill's proposal has come from work in our laboratory<sup>5</sup> that has shown that <sup>3</sup>Pt<sub>2</sub>\* will abstract hydrogen atoms from a large number of different substrates.

Because of the potential importance of H atom transfer in hydrocarbon and related small-molecule activation processes, we are engaged in studies involving systematic variations in the substrate donor to  ${}^{3}Pt_{2}^{*}$ : our aim is to elucidate the factors that control the reaction rates. Here we report the results of kinetic studies of H atom transfers from organometallic hydrides to  ${}^{3}\text{Pt}_{2}^{*}$ :

$${}^{3}\text{Pt}_{2}^{*} + \text{R}_{3}\text{EH} \xrightarrow{\kappa_{H}} {}^{\bullet}\text{Pt}_{2}\text{H} + \text{R}_{3}\text{E}^{\bullet}$$
 (1)

where E = Si, Ge, Sn and R = alkyl or phenyl. Observation of  $Pt_2H_2$  as a reaction product  $^7$  together with the measured kinetic isotope effect (vide infra) indicates that H atom transfer, (1), is the primary step involved in the photoreactivity of  $Pt_2$  with  $R_3EH$ .

The rate constants (eq 1) were measured from the quenching of the  ${}^{3}\text{Pt}_{2}^{*}$  emission ( $\tau_{0} = 10 \ \mu\text{s}$ ) by R<sub>3</sub>EH in degassed acetonitrile solution at 298 K, employing a Nd-YAG laser (excitation at 355 nm, 8-ns fwhm). Linear Stern-Volmer plots following the equation  $\tau_0/\tau = 1 + k_{\rm H}\tau_0[R_3 \text{EH}]$  ( $\tau$  and  $\tau_0$  are the excited-state lifetimes in the presence and absence of the hydride) were obtained.

Because the rate constants for diffusion of the reactants and for encounter-pair dissociation are not expected to vary appreciably in the  $R_3EH$  series, the  $k_H$  values should be a good measure of the reactivity of <sup>3</sup>Pt<sub>2</sub>\*. It is apparent from the data in Table I that the reactivity of <sup>3</sup>Pt<sub>2</sub>\* toward R<sub>3</sub>EH decreases in the order Sn > Ge >> Si. This trend qualitatively parallels that of hydrogen abstraction by tert-butoxy radicals, although the latter is much less pronounced (Table I). The observed rate constants decrease as the E-H bond energy increases (Sn < Ge < Si).<sup>8</sup> The general tendency of R<sub>3</sub>EH species to undergo radical reactions decreases in the same order as  $k_{\rm H}$ . For example, addition to olefins and reduction of organic halides usually take place by radical mechanisms for both Sn and Ge; however, for Si, a catalyzed polar mechanism is preferred.<sup>9</sup> The triphenyl derivatives react with

<sup>(13)</sup> Mukaiyama, T.; Usui, M.; Shimada, E. Chem. Lett. 1975, 1045-1048. (14) Boshagen, H.; Ullrich, J. Chem. Ber. 1959, 92, 1478-1480. Kenner,

G. W.; Stedman, R. J. J. Chem. Soc. 1952, 2089-2094. Also available from CTC organics, Atlanta, GA 30315.

<sup>(15)</sup> To a solution of 1a (417 mg, 1.5 mmol) in cyclohexane (3 mL) was added the DBAD (2.26 g, 7.58 mmol, 5 equiv). The suspension was solub-lized by the addition of  $CH_2Cl_2$  (approximately 500  $\mu$ L) and then irradiated at 350 nm (approximately 18 h). Evaporation and subsequent flash chromatography (30% ether in hexanes) gave 638 mg (73%) of the cycloadduct **2a** ( $R_f = 0.4$  40% ether in hexanes,  $[\alpha]^{22}_D + 106.6^\circ$  (c 1.3, CHCl<sub>3</sub>).

 $<sup>[\</sup>alpha_1^{>_D} - _{3.6}$  (c 0.4, methanol); it. <sup>1</sup> mp  $_{0}^{-1} - _{2}^{-2}$  (c ener);  $[\alpha_1^{>_D} - _{00.5}^{-6}$  (c 1.0, methanol). Also the benzamide derivative gave white needles; mp 134 °C (ether, petroleum ether),  $[\alpha]_D^{2-} - 7.0^{\circ}$  (c 0.5, methanol); lit.<sup>17</sup> mp 134 °C (ethyl acetate, petroleum ether),  $[\alpha]_D^{-} - 7.3^{\circ}$  (methanol). (22) Methyl *N*-acetyl-3,4,6-tri-*O*-acetyl-2-amino-2-deoxy- $\beta$ -D-gluco-pyranoside; mp 161–162 °C,  $[\alpha]_D^{22} - 23.3^{\circ}$  (c 1.0, methanol); lit. mp 163–164 °C,  $[\alpha]_D^{22} - 24.0^{\circ}$  (c 1.0, methanol). See: Kuhn, R.; Kirschenlohr, W. *Chem. Ber.* 1953, 86, 1331–1333.

<sup>(1)</sup> Permanent address: The J. Heyrovský Institute of Physical Chemistry and Electrochemistry, Czechoslovak Academy of Sciences, Vlašská 9, 118 40 Prague, Czechoslovakia

<sup>(2)</sup> Milder, S. J.; Goldbeck, R. A.; Kliger, D. S.; Gray, H. B. J. Am. Chem. Soc. 1980, 102, 6761-6764.

<sup>(3)</sup> Che, C.-M.; Butler, L. G.; Gray, H. B. J. Am. Chem. Soc. 1981, 103, 7796-7797.

<sup>(4)</sup> Marshall, J. L.; Stobart, S. R.; Gray, H. B. J. Am. Chem. Soc. 1984, 106, 3027-3029

<sup>(5)</sup> Marshall, J. L.; Stiegman, A. E.; Gray, H. B. In Excited States and Reactive Intermediates; Lever, A. B. P., Ed.; ACS Symposium Series 307; American Chemical Society: Washington, DC, 1986; pp 166–176. (6) Roundhill, D. M. J. Am. Chem. Soc. 1985, 107, 4354–4356. (7) The dihydride was identified by its absorption spectrum ( $\sigma \rightarrow d\sigma^*$  at 314 nm)<sup>5</sup> and by <sup>31</sup>P NMR ( $\delta$  27,9, J(P,H) = 7,3 Hz, <sup>1</sup>J(P,Pt)  $\approx$  2180 Hz).

For E = Sn or Ge, another reaction product,  $Pt_2(R_3E)_2$ , is formed (Vlček, A.,

<sup>(8)</sup> The gas-phase (298 K) E-H bond energies of (CH<sub>3</sub>)<sub>3</sub>EH species have been reported: 90 (Si), 82 (Ge), 74 kcal mol<sup>-1</sup> (Sn). Jackson, R. A. J. Organomet. Chem. 1979, 166, 17-19.

Table I. Rate Constants  $(k_{\rm H})^a$  for Hydrogen Atom Transfer Reactions

H atom donor	<sup>3</sup> Pt <sub>2</sub> *	t-BuO*	t-Bu*
Et <sub>3</sub> SiH	$2.0 \times 10^{4}$	5.7 × 10 <sup>6 b</sup>	
PhySiH	$1.6 \times 10^{5}$	$1.1 \times 10^{7 b}$	
Ph₃GeH	$2.9 \times 10^{7}$	$8.9 \times 10^{7 b}$	
Ph <sub>3</sub> SnH	$1.0 \times 10^{8}$	$4.0 \times 10^{8 b}$	$3.1 \times 10^{6 d}$
Bu <sub>3</sub> SnH	$1.2 \times 10^{7}$	$2.2 \times 10^{8 c}$	$7.4 \times 10^{5 d}$
Bu <sub>3</sub> SnD	6.9 × 10 <sup>6</sup>	$1.8 \times 10^{8}$ c	$2.7 \times 10^{5 d}$

<sup>a</sup>Rate constants in M<sup>-1</sup> s<sup>-1</sup>. Values for <sup>3</sup>Pt<sub>2</sub>\* measured in acetonitrile at 298 K, t-BuO' reactions studied in 1:2 (v/v) benzene/t-Bu<sub>2</sub>O<sub>2</sub>. <sup>b</sup>Chatgilialoglu, C.; Ingold, K. U.; Lusztyk, I.; Nazran, A. S.; Scaiano, J. C. Organometallics 1983, 2, 1332-1335. T = 300 K. <sup>c</sup>Scaiano, J. C. J. Am. Chem. Soc. 1980, 102, 5399-5400. T = 295 K. <sup>d</sup>Carlsson, D. J.; Ingold, K. U. J. Am. Chem. Soc. 1968, 90, 7047-7055. T = 298 K in benzene.

Table II. Hydrogen Atom Transfer from Bu<sub>3</sub>SnH to Electronically Excited Acceptors

acceptor	excited state	$E_{T}^{a}$	k <sub>H</sub> <sup>c</sup>
acetone	n <b>π*</b>	78	$2 \times 10^{8  d,g}$
benzophenone	·nπ*	69	$4.7 \times 10^{7  e,g}$
2-acetylnaphthalene	$\pi\pi^*$	58	$2.0 \times 10^{6  e,g}$
Pt <sub>2</sub>	dσ*pσ	57.7 <sup>b</sup>	$1.2 \times 10^{7f}$
1-naphthaldehyde	$\pi\pi^*$	56.4	$1.1 \times 10^{6  e,g}$
biacetyl	nπ*	55	$1.5 \times 10^{7  e,g}$

<sup>a</sup> Triplet energies in kcal mol<sup>-1</sup>; data from ref 13, p 290, and from: Murov, S. L. Handbook of Photochemistry; Marcel Dekker: New York, 1973; pp 3-21. <sup>b</sup> Heuer, W. B.; Totten, M. D.; Rodman, G. S.; Hebert, E. J.; Tracy, H. J.; Nagle, J. K. J. Am. Chem. Soc. 1984, 106, 1163-1164. <sup>c</sup>Rate constants in M<sup>-1</sup> s<sup>-1</sup> measured at room temperature. <sup>d</sup> In *n*-hexane. <sup>e</sup> In benzene. <sup>f</sup> In acetonitrile. <sup>g</sup>Reference 14, p 93.

<sup>3</sup>Pt<sub>2</sub>\* faster than the trialkyls, which is analogous to rates found with both tert-butoxy and tert-butyl radicals (Table I).

The kinetic isotope effect measured for the reaction of <sup>3</sup>Pt<sub>2</sub>\* with Bu<sub>3</sub>SnH,  $k_2(H)/k_2(D) = 1.7$ , is in accord<sup>10</sup> with H atom transfer involving a linear Pt-H-Sn transition state with negligible charge transfer.<sup>12-14</sup> This value lies between those found for the reaction of Bu<sub>3</sub>SnH with alkyl<sup>15</sup> (2.3 for Me<sup>•</sup> and *n*-Bu<sup>•</sup>, 1.9 for Et\*) and t-BuO\* radicals (1.2, Table I). Relative to the zero-point energy value,<sup>10</sup> the isotope effect is only slightly lower for the H atom transfer to  ${}^{3}Pt_{2}^{*}$  from Bu<sub>3</sub>SnH than from the  $\alpha$ -(C-H) bond in PhCH(OH)CH<sub>3</sub> (4.7).<sup>16</sup>

Interestingly, the rate constants for H atom transfer are at least 2 orders of magnitude lower than those for electron transfer<sup>3,17</sup> at comparable driving forces.<sup>18</sup> A severe orientation requirement related to the formation of a linear Pt-H-E transition state is the likely explanation of this difference.

Rate constants for H atom transfer from Bu<sub>3</sub>SnH to different excited triplet acceptors are set out in Table II. It is interesting that the reactivity of <sup>3</sup>Pt<sub>2</sub>\* toward Bu<sub>3</sub>SnH is comparable to that

of the  $n\pi^*$  excited states of ketones with similar triplet energies, being greater than that of  $\pi\pi^*$  excited states of organic carbonyls. All  $d^8-d^8$  complexes possess two open axial coordination sites. Excitation of Pt<sub>2</sub> strongly activates these sites, producing a highly energetic ( $E_T = 57.7$  kcal mol<sup>-1</sup>) species with an unpaired electron in an axially localized  $d\sigma$ -antibonding orbital. It would appear that this localized  $d\sigma^*$  electron plays the same role in the reactivity of <sup>3</sup>Pt<sub>2</sub>\* as a nonbonding, oxo-localized electron does in the chemistry<sup>13,14</sup> of excited organic carbonyls.

Acknowledgment. We thank N. J. Turro and D. C. Smith for helpful comments. This research was supported by National Science Foundation Grant CHE84-19828.

## Total Synthesis and Absolute Configuration of 7,20-Diisocyanoadociane

E. J. Corey\* and Plato A. Magriotis

Department of Chemistry, Harvard University Cambridge, Massachusetts 02138 Received September 8, 1986

7,20-Diisocyanoadociane (1 or mirror image), a marine natural product from Adocia sp.,<sup>1</sup> is noteworthy in terms of structure, unprecedented biosynthesis,<sup>1,2</sup> and key position in a growing class of diterpenoid isocyanides.<sup>3</sup> The synthesis of 1 has been a conspicuous problem which has been taken as a challenge by a number of research groups. Reported herein is the first synthesis of this interesting perhydropyrene by an enantioselective route which permits assignment of the absolute configuration indicated in 1.4

The acid chloride 2, readily available in two steps from (1R,2S, 5R)-(-)-menthol and glutaric anhydride (heating at 90 °C for 10 h followed by reaction with oxalyl chloride in benzene at 23 °C, 75% overall), was converted into vinyl ketone 3 (1.1 equiv of vinyltri-n-butylstannane in tetrahydrofuran (THF) with 0.34 mol % of Pd(PPh<sub>3</sub>)<sub>4</sub> at 70 °C for 2 h, 90% yield).<sup>5,6</sup> Enone 3 was transformed into the corresponding ethylene ketal (90% overall yield) by a new method consisting of two steps: (1) treatment of a 1 M solution of 3 in  $CHCl_2$  with (phenylseleno)trimethylsilane<sup>7</sup> (1.3 equiv), ethylene glycol (5 equiv), and I<sub>2</sub> (0.025 equiv) at 65 °C for 4 h to form the  $\beta$ -(phenylseleno) ethylene ketal (99%); (2) oxidation of selenide to selenoxide (1.5 equiv of m-chloroperbenzoic acid in CH2Cl2 at -20 °C for 15 min) followed by addition of dimethyl sulfide (0.8 equiv) and diisopropylamine (3 equiv) and warming to 60 °C to effect elimination (6.5 h at 60 °C), and finally gradient elution chromatography on silica gel (sg)(hexane-ether). Attempts to prepare this ketal by conventional acid-catalyzed direct ketalization were not successful.

The conversion of the ethylene ketal of 3 to diester 4 depended on new methodology for enantioselective and diastereoselective Michael addition which has recently been reported.<sup>8</sup> The ester enolate of the ketal 3 was generated at -78 °C in THF using 1.1 equiv of lithium diisopropylamide (LDA) for 45 min, methyl crotonate (1.1 equiv, E isomer) was added, the reaction mixture was quenched (HOAc) after 1 h at -78 °C, and the product was isolated by extractive workup and sg flash chromatography. The major Michael adduct (80% yield) was, as expected, the desired threo isomer 4 (threo/erythro ratio 8:1, ca. 60% ee).<sup>8,9</sup> Removal

(3) (a) Kazlauskas, R.; Murphy, P. T.; Wells, R. J.; Blount, J. F. Tetra-hedron Lett. 1980, 21, 315. (b) Faulkner, D. J. Tetrahedron 1977, 33, 1421.

(8) Corey, E. J.; Peterson, R. T. Tetrahedron Lett. 1985, 26, 5025.

<sup>(9)</sup> Armitage, D. A.; Riviere, P.; Riviere-Baudet, M.; Satge, J.; Davies, A. G.; Smith, P. J. In Comprehensive Organometallic Chemistry; Wilkinson, G., Stone, F. G. A., Abel, E. W., Eds.; Pergamon Press: Oxford, 1982; Vol. 2, pp 1-203, 399-627.

<sup>(10)</sup> Taking into account the low Sn-H force constant in R<sub>3</sub>SnH species ( $\nu$ (Sn-H) = 1837 cm<sup>-1</sup> in (CH<sub>3</sub>)<sub>3</sub>SnH<sup>11</sup>,  $k_f \simeq 2 \times 10^2$  N m<sup>-1</sup>), the value of the zero-point energy isotope effect for Sn-H bond breaking may be estimated<sup>12</sup> a 2.6 this is more defined. mated<sup>12</sup> as 3.6; this is much smaller than the analogous value for a typical C-H bond (6.9). The observed isotope effect is quite large, as it accounts for about 47% of the zero-point energy value.

<sup>(11)</sup> Nakamoto, K. Infrared and Raman Spectra of Inorganic and Coordination Compounds, 3rd ed.; Wiley: New York, 1978; p 380-381. (12) Melander, L.; Saunders, W. H., Jr. Reaction Rates of Isotopic

Molecules: Wiley: New York, 1980; pp 4-45, 129-170. (13) Turro, N. J. Modern Molecular Photochemistry; Benjamin/Cum-

mings: Menlo Park, CA, 1978; pp 363-385. (14) Scaiano, J. C. J. Photochem. 1973/1974, 2, 81-118.

<sup>(15)</sup> Chatgilialoglu, C.; Ingold, K. U.; Scaiano, J. C. J. Am. Chem. Soc. 1981, 103, 7739-7742.

<sup>(16)</sup> Harvey, E. L.; Gray, H. B., unpublished results.

<sup>(17)</sup> Peterson, J. R.; Kalyanasundaram, K. J. Phys. Chem. 1985, 89, 2486-2492

<sup>(18)</sup> Assuming a  $Pt_2$ -H dissociation energy of 60 kcal mol<sup>-1</sup> (Pearson, R. G. Chem. Rev. **1985**, 85, 41-49), the driving force for H atom transfer is roughly estimated to be  $(60 - D_{\rm EH} + E_{\rm T}) \ge 30$  kcal mol<sup>-1</sup>.

<sup>(1)</sup> Baker, J. T.; Wells, R. J.; Oberhänsli, W. E.; Hawes, G. B. J. Am. Chem. Soc. 1976, 98, 4010.

<sup>(2)</sup> Garson, M. J. J. Chem. Soc., Chem. Commun. 1986, 35.

<sup>(</sup>c) Faulkner, D. J. Nat. Prod. Rep. 1984, 1, 551.
(4) The original X-ray crystallographic determination of structure<sup>1</sup> did not allow assignment of absolute configuration. (5) (a) Kosugi, M.; Shimizu, Y.; Migita, T. Chem. Lett. 1977, 1423. (b)

The reaction byproduct, Bu<sub>3</sub>SnCl, was removed by washing an ether extract with 50% aqueous KF solution.

<sup>(6)</sup> All reactions involving air-sensitive materials were conducted under an inert (N2 or Ar) atmosphere.

<sup>7)</sup> Detty, M. R.; Seidler, M. D. J. Org. Chem. 1981, 46, 1283