

A NEW PROCESS FOR THE ENANTIOSELECTIVE SYNTHESIS OF CHIRAL α -ARYLOXY- AND α -HYDROXY ACIDS

E. J. Corey and John O. Link

Department of Chemistry, Harvard University, Cambridge, Massachusetts, 02138


Summary: Chiral trichloromethyl carbinols **3**, readily available by catalytic enantioselective reduction of trichloromethyl ketones, are converted with inversion of configuration into chiral α -aryloxy and α -hydroxy carboxylic acid derivatives.

The reduction of a series of trichloromethyl ketones **1** by catecholborane in the presence of oxazaborolidine **2** as catalyst¹ produces the corresponding chiral secondary trichloromethyl carbinols **3** with high enantioselectivity.² This is an exceedingly useful process since the carbinols **3** can be converted in two simple steps (reaction with aqueous $\text{NaN}_3\text{-NaOH}$ and subsequent reduction with $\text{H}_2\text{-Pd/C}$) to α -amino acids which are obtained in enantiomerically pure form by simple recrystallization (Scheme I).² We describe herein another valuable application of chiral trichloromethyl carbinols, specifically to the enantioselective synthesis of chiral α -hydroxy acids via the corresponding *p*-methoxyphenyl ether derivatives, themselves useful building blocks for synthesis.³

Table I summarizes the results obtained for the reduction of a representative set of four trichloromethyl ketones using 1.5 equiv of catecholborane and 0.1 equiv of catalyst **2** in toluene or methylene chloride solution to give the chiral alcohols **3**.² The new process for the transformation of **3** into chiral α -aryloxy acids and α -hydroxy acids is outlined in Scheme II.

The reaction of chiral trichloromethyl carbinols (**3**) with *p*-methoxyphenol in basic aqueous dimethoxyethane at ambient temperature provided the corresponding α -*p*-anisloxy acid derivatives **4** with clean inversion of configuration by way of an intermediate 1,1-dichlorooxirane² as shown in Scheme II.^{4,5} The yields of **4** in the case of primary alkyl substitution, $\text{R} = n\text{-C}_5\text{H}_{11}$ (77%), and $\text{R} = \text{C}_6\text{H}_5\text{CH}_2\text{CH}_2$ (84%), were higher than for secondary alkyl substitution, $\text{R} = \text{cyclohexyl}$ (62%).^{5b} As expected the transformation of **3** to **4** for $\text{R} = \text{tertiary butyl}$ was quite inefficient (10% yield) because of the extreme degree of steric retardation of displacement for a neopentyl system.

Scheme I



Recrystallization of the acids **4**, $R = n\text{-C}_5\text{H}_{11}$, $\text{C}_6\text{H}_5\text{CH}_2\text{CH}_2$, or cyclohexyl, from ether–hexane or hexane afforded each α -*p*-anisloxy acid in 100% enantiomeric purity as determined by HPLC analysis using a chiral column.⁴ The various acids were converted in 98–99% yield to the corresponding methyl esters **5** using dimethyl sulfate at pH 7–8 in a phase transfer system. Finally, these esters were treated with ceric ammonium nitrate in dimethylformamide–water at $-20\text{ }^\circ\text{C}$ (30 min) and $0\text{ }^\circ\text{C}$ (30 min) to give the corresponding enantiomerically pure α -hydroxy acid methyl esters in 86–90% yield.⁶ It is important to carry out this deprotection at lower temperatures to avoid competing nitration of the *p*-anisloxy group in **5**. In addition, the oxidative deprotection of the methyl esters **5** is advantageous since it avoids the problem of oxidative decarboxylation which is observed in the reaction of the free carboxylic acids **4** with ceric ammonium nitrate.


In conclusion, a highly effective process for the synthesis of optically pure α -hydroxy acids has been developed which has the advantage of employing a catalytic step for the introduction of chirality and a chiral reagent which can be recovered efficiently. The new process should allow access to a large series of useful chiral α -hydroxy acids and esters.⁷

The following procedures are representative.

(*R*)-4-Phenyl-1,1,1-trichloro-2-butanol. Addition of 4-phenyl-1,1,1-trichloro-2-butanol (1.59 g, 7.4 mmol) to a solution of preformed oxazaborolidine catalyst **2**² (3.7 mL, 0.74 mmol, 0.2 *M* in toluene, 0.1 equiv), cooling to $-78\text{ }^\circ\text{C}$ and dropwise addition of freshly distilled catecholborane (5.5 mL, 11 mmol, 2 *M* in toluene) over 10 min with vigorous stirring afforded a white precipitate. At 4 h the colorless solution was homogeneous and after 12 h at $-78\text{ }^\circ\text{C}$ the reaction was quenched with methanolic HCl (1.8 mL, 0.5 *M*) and allowed to warm to $23\text{ }^\circ\text{C}$. Partial concentration *in vacuo* afforded a fine white precipitate ((*S*)- α , α -diphenylprolinol•HCl) which was recovered by filtration. The filtrate was diluted with 40 mL of ether, washed with pH 13 buffer until colorless (6 x 20 mL), then brine (3 x 10 mL), dried over magnesium sulfate and concentrated *in vacuo* to afford 1.54 g (96%) of (*R*)-4-phenyl-1,1,1-trichloro-2-butanol as a colorless solid after chromatography through a short plug of silica gel (sg) (10 : 1 hexane–ethyl acetate); mp $53\text{--}5\text{ }^\circ\text{C}$; $[\alpha]_{\text{D}}^{24} +49.4^\circ$ ($c = 2.39$, CHCl_3); 95% ee ^1H NMR (270 MHz, CDCl_3) δ 7.4–7.2 (m, 5H), 4.0 (ddd, 1H, $J = 9.3, 5.4, 1.6$ Hz), 3.0 (m, 1H), 2.8 (d, 1H, $J = 5.4$ Hz, -OH), 2.75 (m, 1H), 2.4 (m, 1H), 2.0 (m, 1H); FTIR (neat) 3400, 1455 cm^{-1} ; CIMS 289 $[\text{M}+\text{Cl}]^-$; HRMS: calcd. for $[\text{C}_{10}\text{H}_{11}\text{Cl}_3\text{O}+\text{Cl}]^-$: 288.9535; found: 288.9517.

(*S*)-2-*p*-Anisyloxy-4-phenylbutanoic acid. To a mixture of (*R*)-4-phenyl-1,1,1-trichloro-2-butanol (1.06 g, 4.18 mmol), and *p*-methoxyphenol (1.04 g, 8.36 mmol), in dimethoxyethane (DME) (24 mL) was added water (15.9 mL) and then (dropwise) aqueous NaOH (2.33 mL, 24.9 mmol, 10.75 *M*) over 10 min with vigorous stirring at $23\text{ }^\circ\text{C}$ (the solution remained homogeneous). After 12 h the DME was removed at 20 torr and the mixture was diluted with sat. NaHCO_3 (50 mL) and washed with CH_2Cl_2 (3 x 20 mL) to remove excess *p*-methoxyphenol. The CH_2Cl_2 was extracted with sat. NaHCO_3 (2 x 10 mL). The combined aqueous extracts were acidified to *ca.* pH 4 with solid KH_2PO_4 , extracted with ether (40 mL), then acidified to pH 1 (12 *M* HCl) and extracted with ether (2 x 20 mL); the ether was dried (MgSO_4) and concentrated *in vacuo* to afford (*S*)-2-*p*-anisloxy-4-phenylbutanoic acid (1.01 g, 84%) as a colorless solid after filtration through a 3 cm sg plug with 9 : 1 toluene–ethyl acetate - 3% acetic acid as eluent. Recrystallization from ether–hexane afforded a 92% recovery of optically pure acid; mp $81.5\text{--}82.5\text{ }^\circ\text{C}$; $[\alpha]_{\text{D}}^{24} -31.9^\circ$ ($c = 1.6$, CHCl_3); ^1H NMR (270 MHz, CDCl_3) δ 7.3–7.2 (m, 5H), 6.8 (s, 4H), 4.6 (dd, $J = 7.5, 5.1$ Hz, 1H), 3.8 (s, 3H), 2.9

Scheme II



(m, 2H), 2.2 (m, 2H); FTIR (neat) 3400-2700, 1723, 1507 cm⁻¹; FAB MS: 286 [M]⁺; HRMS: calcd. for [C₁₇H₁₈O₄]⁺: 286.1205; found: 286.1208.

(S)-Methyl-2-*p*-anisyoxy-4-phenylbutanoate. To a vigorously stirred mixture of (S)-2-*p*-anisyoxy-4-phenylbutanoic acid (573 mg, 2.0 mmol), CH₂Cl₂ (1.5 mL), pH 7 aqueous buffer (1.8 mL), NaOH (186 μL, 2.0 mmol, 10.75 M), and benzyltriethylammonium chloride (62 mg, 0.2 mmol), was added dimethyl sulfate (285 μL, 3.0 mmol) at 23 °C. After 6 h NaOAc (272 mg, 2.0 mmol) was added to consume unreacted dimethyl sulfate and the mixture was stirred for 1 h. The mixture was diluted with 10 mL of pH 7

Table I. Enantioselective Reduction of α,α,α-Trichloromethyl Ketones

R in RCOCCl ₃	solvent	temperature, °C	
		(time, h)	3, % ee ^b
<i>n</i> -C ₅ H ₁₁	toluene	-60 ^a (12)	95 ^c
C ₆ H ₅ (CH ₂) ₂	toluene	-78 (12)	95 ^c
<i>c</i> -C ₆ H ₁₁	CH ₂ Cl ₂	-20 ^a (48)	92 ^d
<i>t</i> -C ₄ H ₉	toluene	-20 ^a (56)	98 ^c

^aThese reactions were initiated at -78 °C and brought to the indicated temperature after 1 h. ^bAlcohol % ee determined by Chiracel (Diacel Co.) HPLC analysis with the following columns: ^cChiracel AD. ^dChiracel OJ. ^eChiracel OD analysis on the benzoate ester.

buffer and extracted with ether (2 x 20 mL), dried (MgSO₄), and concentrated *in vacuo*; chromatography on a short sg column (10:1 hexane–ethyl acetate) afforded optically pure (*S*)-methyl-2-*p*-anisyoxy-4-phenylbutanoate as a colorless oil (589 mg, 98%); $[\alpha]_D^{24}$ -39.5° (*c* = 1.33, CHCl₃); ¹H NMR (270 MHz, CDCl₃) δ 7.3–7.2 (m, 5H), 6.8 (s, 4H), 4.5 (dd, *J* = 7.9, 4.6 Hz, 1H), 3.8 (s, 3H), 3.7 (s, 3H), 2.9 (m, 2H), 2.2 (m, 2H); FTIR (neat) 1755, 1734, 1506 cm⁻¹; EIMS: 300 [M]⁺; HRMS: calcd. for [C₁₈H₂₀O₄]⁺: 300.1361; found: 300.1349.

(*S*)-Methyl-2-hydroxy-4-phenylbutanoate. A solution of (*S*)-methyl-2-*p*-anisyoxy-4-phenylbutanoate (569 mg, 1.90 mmol, in 1.7 mL of 4:1 DMF - H₂O) was added dropwise by cannula over 15 min to a cold (-23 °C) well-stirred solution of ceric ammonium nitrate (5.2 g, 9.5 mmol) in 4:1 DMF-H₂O (13 mL). After reaction for 30 min at -23 °C the solution was maintained at 0 °C for 30 min and then diluted with 10 mL each of 1:1 ether–hexane and water with stirring at 0 °C. Water (30 mL) was added and the mixture was extracted with 1:1 ether–hexane (3 x 20 mL). The organic extracts were washed (10 mL of 0.5 M Na₂CO₃), dried (MgSO₄), and concentrated *in vacuo* and the resultant orange oil was chromatographed on sg with 5:1 hexane–ethyl acetate to afford optically pure⁴ (*S*)-methyl-2-hydroxy-4-phenylbutanoate (330 mg, 90%) as a colorless oil; $[\alpha]_D^{24}$ +32.4° (*c* = 2.1, CHCl₃), lit.^{6b} $[\alpha]_D^{25}$ -32.5° (CHCl₃) (*R*) enantiomer; ¹H NMR (270 MHz, CDCl₃) δ 7.3–7.2 (m, 5H), 4.2 (dd, 1H, *J* = 7.9, 4.0), 3.75 (s, 3H), 2.8 (m, 2H), 2.1 (m, 1H), 1.9 (m, 1H); IR (neat) 3450, 1734 cm⁻¹; EIMS: 194 [M]⁺; HRMS: calcd. for [C₁₁H₁₄O₃]⁺: 194.0943; found: 194.0939.

References and Notes

- (a) Corey, E. J.; Bakshi, R. K.; Shibata, S. *J. Am. Chem. Soc.* **1987**, *109*, 5551-5553. (b) Corey, E. J.; Bakshi, R. K.; Shibata, S.; Chen, C. P.; Singh, V. K. *J. Am. Chem. Soc.* **1987**, *109*, 7925-7926. (c) Corey, E. J.; Shibata, S.; Bakshi, R. K. *J. Org. Chem.* **1988**, *53*, 2861-2863. (d) Corey, E. J. in *Proceedings 31st National Organic Chemistry Symposium of the American Chemical Society*; 1989; p 1. (e) Corey, E. J.; Link, J. O. *Tetrahedron Lett.* **1989**, *30*, 6275-6278. (f) Corey, E. J.; Bakshi, R. K. *Tetrahedron Lett.* **1990**, *31*, 611-614. (g) Corey, E. J. *Pure Appl. Chem.* **1990**, *62*, 1209-1216.
- Corey, E. J.; Link, J. O. *J. Am. Chem. Soc.* **1992**, *114*, 1906.
- For other pathways for the enantioselective synthesis of α-hydroxy acids by oxazaborolidine catalyzed enantioselective reduction see ref. 1f.
- Analysis for enantiomeric purity was performed using a Chiracel OD analytical column (0.46 mm, 25 cm) with isopropyl alcohol in hexane for elution (10% *i*-PrOH for R = *n*-C₅H₁₁ and R = C₆H₅(CH₂)₂; 1.5% *i*-PrOH for R = cyclohexyl). In each case the major enantiomer (*S*) eluted more rapidly than the minor (*R*) enantiomer. The retention times observed at a flow rate of 1 mL per min were as follows (major, minor in min): **5**, R = *n*-C₅H₁₁ (5.0, 6.5); **5**, R = *c*-C₆H₁₁ (7.3, 8.3); **6**, R = C₆H₅(CH₂)₂ (7.9, 10.5).
- (a) Yields with *p*-anisoxide as the nucleophile are significantly higher than those employing direct hydrolysis to the α-hydroxy acid with hydroxide (BnEt₃NCI, NaOH, H₂O, CH₂Cl₂, **3** R = C₆H₅(CH₂)₂; 40%). Further, the *p*-anisyoxy group provides selective hydroxyl protection and also imparts UV activity for convenient TLC and HPLC detection. See Corey, E. J.; Barcza, S.; Klotmann, G. *J. Am. Chem. Soc.* **1969**, *91*, 4782-4786 for the synthesis of α-aryloxy carboxylic acids from trichloromethyl carbinols and phenoxides. (b) For R = *c*-C₆H₁₁ 4 equiv of *p*-methoxyphenol and 8 equiv of NaOH were used.
- (a) **6**, R = *n*-C₅H₁₁ $[\alpha]_D^{24}$ +12.8° (*c* = 2.35, CHCl₃), lit. $[\alpha]_D^{23}$ +10.7° (*c* = 4.57, CHCl₃): Grieco, P. A.; Takigawa, T.; Vedananda, T. R. *J. Org. Chem.* **1985**, *50*, 3111-3115. (b) **6**, R = C₆H₅(CH₂)₂ $[\alpha]_D^{24}$ +32.4° (*c* = 2.1, CHCl₃), lit. $[\alpha]_D^{25}$ -32.5° (CHCl₃) (*R*) enantiomer: Satoh, T.; Onda, K.-i.; Yamakawa, K. *Tetrahedron Lett.* **1990**, *31*, 3567-3570. (c) **6**, R = *c*-C₆H₁₁ $[\alpha]_D^{24}$ +34.9° (*c* = 1.49, CHCl₃), lit. $[\alpha]_D^{20}$ -31.3° (*c* = 2.36, CHCl₃) (*R*) enantiomer: Ko, K. Y.; Frazee, W. J.; Eliel, E. L. *Tetrahedron*. **1984**, *40*, 1333-1343.
- This research was assisted financially by generous grants from the National Institutes of Health and the National Science Foundation.