Chemical Properties of Dodecahedrane. **Monofunctionalization Reactions**

Leo A. Paquette,* Jeffrey C. Weber, and Tomoshige Kobayashi¹

> Evans Chemical Laboratories, The Ohio State University Columbus, Ohio 43210

Received September 21, 1987

The I_h symmetry of the pentagonal dodecahedrane (1) qualifies it as the most highly symmetric of all organic molecules. One end result of this aesthetically appealing feature is the identity of all 20 constituent methine units. Since disclosure was made of the first synthesis of this spherical C₂₀H₂₀ system,^{2,3} significant improvements have been realized in the Paquette approach^{4,5} such that the compound can currently be obtained in 25-50-mg lots if desired. Since differently substituted dodecahedranes are of interest in their own right and as precursors to a host of interesting molecules, a systematic investigation of the chemical reactivity of 1 has been undertaken. Reported herein are key observations concerning the direct monofunctionalization of this unusual hydrocarbon.

The geometry^{6,7} and rigid conformational characteristics⁸ of 1 are recognized to require almost ideal sp³ hybridization at carbon. Development of positive charge at any site is not expected to be met with ready accommodation of planarity. Moreover, the dodecahedryl cation (DDH⁺) clearly lacks the normal stabilizing effect of alkyl substitution,⁹ since no C-C (or C-H) bond can be oriented trans to the vacant orbital that is strongly directed to the molecular exterior. For these reasons, the generation of DDH+ was expected to be akin to the behavior exhibited by the perhydrotriquinacene system, whose spectacular deviation from the norm¹⁰ (109-fold rate retardation in solvolysis of the 10-tosylate)¹¹ has been documented. The situation is quite unlike that available in adamantane,¹² and indeed very significant differences have been found between adamantane and 1.13

Stirring 1 in neat bromine at room temperature¹⁴ overnight led quantitatively to monobromide 2a. Although 2a was unreactive toward K₂CO₃ or AgNO₃ in refluxing aqueous tetrahydrofuran,¹⁵

- (2) (a) Ternansky, R. J.; Balogh, D. W.; Paquette, L. A. J. Am. Chem. Soc. 1982, 104, 4503. (b) Paquette, L. A.; Ternansky, R. J.; Balogh, D. W.; Kentgen, G. Ibid. 1983, 105, 5446.
- (3) An alternative route to 1 from pagodane has recently been described: Fessner, W.-D.; Murty, B. A. R. C.; Worth, J.; Hunkler, D.; Fritz, H.; Prinzbach, H.; Roth, W. D.; Schleyer, P. v. R.; McEwen, A. B.; Maier, W. F. Angew. Chem., Int. Ed. Engl. 1987, 26, 452.
- (4) (a) Paquette, L. A.; Miyahara, Y.; Doecke, C. W. J. Am. Chem. Soc. 1986, 108, 1716. (b) Paquette, L. A.; Miyahara, Y. J. Org. Chem. 1987, 52, 1265
 - (5) Kobayashi, T.; Miyahara, Y.; Weber, J. C., unpublished results.
- (6) (a) Paquette, L. A.; Balogh, D. W.; Usha, R.; Kountz, D.; Christoph, G. G. Science (Washington, D. C.) 1981, 211, 575. (b) Christoph, G. G.; Engel, P.; Usha, R.; Balogh, D. W.; Paquette, L. A. J. Am. Chem. Soc. 1982,
- 104, 784
- (7) Gallucci, J. C.; Doecke, C. W.; Paquette, L. A. J. Am. Chem. Soc. 1986, 108, 1343.
 - (8) Ermer, O. Angew. Chem., Int. Ed. Engl. 1977, 6, 411.
 - (9) Hoffmann, R. J. Chem. Phys. 1964, 40, 2480.
- (10) Paquette, L. A.; Meehan, G. V.; Marshall, S. J. J. Am. Chem. Soc. 1969, 91, 6779
- (11) Bingham, R. C.; Schleyer, P. v. R. J. Am. Chem. Soc. 1971, 93, 3189. (12) (a) Fort, R. C.; Schleyer, P. v. R. Chem. Ber. 1964, 64, 277. (b) Engler, E. M.; Schleyer, P. v. R. In MTP International Review of Scien Vol. 5, Alicyclic Compounds; Parker, W., Ed.; Butterworths: London, 1973;
- Chapter 7. (13) Space limitations preclude substantive discussion of these issues here. However, detailed comparison will be made later in a full paper
- (14) The bromination of adamantane is reported to require heating at the reflux temperature for 4.5 h: Stetter, H.; Schwarz, M.; Hirschhorn, A. Chem.
- Ber. 1959, 92, 1629. (15) Bromoadamantane is converted into adamantanol (95%) during 4 h under these conditions: ref 12a, eq 23.



conversion to chloride 2b (91%) and fluoride 2c (89%) proved possible with appropriate Lewis acid catalysis (FeCl₃, CH₂Cl₂, room temperature, 4 h; AgBF₄, CH₂Cl₂, Et₂O, room temperature, 20 h).¹⁶ The sluggish response of DDH-Br to ionization was also encountered in our inability to achieve Koch-Haaf carboxylation under standard conditions.^{14,17} However, when recourse was made instead to bubbling CO into magic acid solutions of 2a,¹⁸ access was successfully gained to 3a. Direct treatment with CH_2N_2 gave in 60% yield the methyl ester 3b, the structural features of which were corroborated by X-ray crystallographic analysis.¹⁹ Alcohol 3c and aldehyde 3d were derived conventionally from 3b. Exposure of the methyl ester to a large excess of Me₂AlNH₂ according to Weinreb,20 but with gentle warming, provided amide 3e in quantitative yield after 24 h.

Still more varied carbon functionalization was achieved by other avenues. For example, warming a mixture of 2a and FeCl₃ in benzene for 3 h furnished the phenyl derivative 4 (94%). Direct methylation was most conveniently realized by exposure of 2a to Me_3Al in hexane²¹ (room temperature 24 h, 90%). The identity of 3f with material prepared by total synthesis²² was accomplished spectroscopically.

Investigation of the controlled oxygenation of 1 was successful from several directions. Dodecahedranol (5b) was obtained in 76% yield by treatment of 1 with Pb(OAc)₄ in a CF₃COOH-CH₂Cl₂ solvent system containing LiCl²³ followed directly by saponification of the trifluoroacetate 5a. Alternatively, 5a is available by reaction of 2b with AgOCOCF₃ in CF₃COOH (room temperature, 18 h, 96%). A related protocol (AgOTf, CH₃OH-CH₂Cl₂ (1:1), room temperature, 18 h) was utilized to prepare methyl ether 5c (71% from 2b). Dissolution of 2a into magic acid at -78 °C and quenching of the cation solution in cold methanol²⁴ delivered 5c with almost equal efficiency (61%). Treatment of 1 with commercial $NO_2^+BF_4^-$ in $CHCl_3^{25}$ gave rise to nitrate ester **5d** (84%) rather than the nitro compound.

Finally, formation of a nitrogen derivative was achieved by again making recourse to the triflate.²⁶ Suspension of 2a and AgOTf

- (18) (a) Souma, Y.; Sano, H. Bull. Chem. Soc. Jpn. 1976, 49, 3335. (b) Yoneda, N.; Fukuhara, T.; Takahashi, Y.; Suzuki, A. Chem. Lett. 1983, 17. (19) Taylor, R. T., unpublished results.
 - (20) Basha, A.; Lipton, M.; Weinreb, S. M. Tetrahedron Lett. 1977, 4171.
 - (21) Kennedy, J. P. J. Org. Chem. 1970, 35, 532.
- (22) (a) Paquette, L. A.; Ternansky, R. J.; Balogh, D. W. J. Am. Chem. Soc. 1982, 104, 4502. (b) Paquette, L. A.; Ternansky, R. J.; Balogh, D. W.; Taylor, W. J. Ibid. 1983, 105, 5441.
- (23) Jones, S. R.; Mellor, J. M. J. Chem. Soc., Perkin Trans. 1, 1976, 2576.
- (24) (a) Olah, G. A.; Lukas, J. J. Am. Chem. Soc. 1968, 90, 933. (b) Olah, G. A.; et al. Ibid. 1985, 107, 2764.
 - (25) Olah, G. A.; Lin, H. C. J. Am. Chem. Soc. 1971, 93, 1259.
- (26) Takeuchi, K.; Moriyama, T.; Kinoshita, T.; Tachino, H.; Okamoto, K. Chem. Lett. 1980, 1395.

0002-7863/88/1510-1303\$01.50/0 © 1988 American Chemical Society

⁽¹⁾ The Ohio State University Postdoctoral Fellow, 1986-1988.

⁽¹⁶⁾ Fry, A. J.; Migron, Y. Tetrahedron Lett. 1979, 3357.

⁽¹⁷⁾ Koch, H.; Haaf, W. Org. Synth. 1964, 44, 1.

Table I. Monosubstituted Dodecahedranes

		¹³ C δ	m/z (M ⁺)	
substituent	mp, °C	(quart C)	calcd	obsd
Br	>240	96.98	340.0650	340.0628
Cl	>280	104.39	294.1175	294.1140
F	>260	137.64	278.1487	278.1476
СООН	>250	b	304.1463	304.1449
COOMe	192-193	84.54	319 ^c	319
CH ₂ OH	>250	81.86	290.1681	290.1659
СНО	>250	90.00	288.1514	288.1519
CONH ₂	>250	85.61	303.1623	303.1640
C ₆ H ₅	200-201	84.42	336.1870	336.1874
OCOCF,	183-185ª	126.47	258.1408 ^d	258.1407
ОН	>250	115.99	276.1515	276.1482
ONO ₂	220	129.89	259°	259
NHCOCH ₃	>250	95.77	317.1780	317.1772
				1) 1/2 14

^bVery low solubility. ^cCI (M + 1). ^d(M⁺ -^a Dec 220 °C. CF_3COOH). $(M^+ - ONO_2)$.

in hot acetonitrile and subsequent hydration delivered 6 (100%).

Acknowledgment. We are grateful to the National Institutes of Health for their generous financial support of this research program (Grant AI-11490).

Chemical Properties of Dodecahedrane. The Dodecahedryl Cation and 1,16-Dodecahedryl Dication. Regioselective Difunctionalization of the Sphere^{1a}

George A. Olah* and G. K. Surya Prakash

Donald P. and Katherine B. Loker Hydrocarbon Research Institute and Department of Chemistry University of Southern California Los Angeles, California 90089

Tomoshige Kobayashi^{1b} and Leo A. Paquette*

Evans Chemical Laboratories, The Ohio State University Columbus, Ohio 43210

Received September 21, 1987

The challenge of defining practical synthetic routes for arrival at dodecahedrane (1a) has been met in two laboratories.^{2,3} The preceding report defines for the first time various protocols capable of delivering a wide range of monofunctionalized dodecahedranes.⁴ Practical exploitation of this chemistry rests heavily, although not exclusively, on transient generation and efficient trapping of the monocation. Several empirical observations made in the course of these studies implicated for $C_{20}H_{19}^+$ (2) a lability that was not at all expected.⁵ For this reason, we sought to generate and examine 2 under long-life conditions. Herein is described the direct spectral observation of 2 in superacid solution, an analysis of its stability and capacity for degenerate isomerization, and the remarkable ease of its conversion to the 1,16-dication 4.

Careful dissolution of 5 mg of chlorododecahedrane $(1b)^4$ in 1 mL of SO₂ClF containing \sim 250 mg of SbF₅ at -78 °C in a 5-mm NMR tube gave a pale yellow-colored solution. The 200



MHz ¹H NMR spectrum^{6a} of this solution at -70 °C revealed a set of three absorptions at δ 4.64 (br, 3 H), 3.05 (br, 7 H), and 2.59 (br, 9 H) as seen in Figure 1A^{6b} The peaks did not split further at 500 MHz. The 50 MHz ¹³C spectrum consisted of six absorptions at 363.9 (s), 81.1 (d), 64.4 (d), 64.1 (d), 63.0 (d), and 60.9 (d) ppm, clearly indicating formation of the static dodecahedryl cation (2). Ion 2 was similarly obtained from dodecahedranol (1c),⁴ although more slowly. The limited solubility of dodecahedrane necessitated that the superacidic medium be warmed to 0 °C for dissolution to occur. Under these conditions, smooth conversion to 2 also materialized.

We considered that the dodecahedryl cation could undergo rapid hydrogen scrambling (through 1,2-hydride shifts) similar to that observed in cyclopentenyl cation 7, a process that would render

all the carbon and hydrogen atoms equivalent. However, no such degenerate process was observed as indicated by the lack of change in the ¹H NMR line shapes, even when solutions of 2 were allowed to warm to 0 °C. Thus, the lower barrier for such degenerate rearrangement is approximately 15 kcal/mol. The findings presented below suggest that the actual barrier is not apt to be much higher however.

Upon standing in the superacidic solution for 6-7 h at -50 °C, monocation 2 is slowly and irreversibly transformed into a new species displaying much simplified NMR spectra. Only two absorptions are seen at δ 4.74 (br, 6 H) and 3.23 (br, 12 H) in

^{(1) (}a) Considered as Stable Carbocations, Part 271 at the University of Southern California. (b) The Ohio State University Postdoctoral Fellow, 1986-1988.

 ^{(2) (}a) Ternansky, R. J.; Balogh, D. W.; Paquette, L. A. J. Am. Chem.
Soc. 1982, 104, 4503. (b) Paquette, L. A.; Ternansky, R. J.; Balogh, D. W.;
Kentgen, G. Ibid. 1983, 104, 5446.
(3) Fessner, W.-D.; Murty, B. A. R. C.; Worth, J.; Hunkler, D.; Fritz, H.;
Prinzbach, H.; Roth, W. D.; Schleyer, P. v. R.; McEwen, A. B.; Maier, W.

F. Angew. Chem., Int. Ed. Engl. 1987, 26, 452

⁽⁴⁾ Paquette, L. A.; Weber, J. C.; Kobayashi, T., preceding paper in this issue

⁽⁵⁾ Unpublished observations of J. C. Weber and T. Kobayashi at The Ohio State University,

^{(6) (}a) The NMR spectra were recorded on a Varian Associates Model VXR-200 NMR spectrometer equipped with a variable temperature broad band probe. (b) The observed relative intergral area of the three absorptions of 3:7:9 in 2 indicates that the apical hydrogen atom at the 16-position which is farthest from the positive charge experiences more deshielding ($\delta = 3.05$) than the other belt hydrogens ($\delta = 2.59$) which are much closer to the positive charge. This may be due to a unique cage effect in the dodecahedrane skeleton.