

[Chem. Pharm. Bull.]
34(5)1968—1979(1986)

Components of *Broussonetia kazinoki* SIEB. I. Structures of Two New Isoprenylated Flavans and Five New Isoprenylated 1,3-Diphenylpropane Derivatives^{1,2)}

JUNKO IKUTA (née MATSUMOTO),^a YOSHIO HANO,^a TARO NOMURA,^{*,a}
YOSHIYUKI KAWAKAMI,^b and TADASHI SATO^b

Faculty of Pharmaceutical Sciences, Toho University, ^a Miyama, Funabashi,
Chiba 274, Japan and Tsukuba Research Laboratory, Eisai Company,^b
Tokodai, Toyosato, Tsukuba, Ibaraki 300-26, Japan

(Received September 27, 1985)

Two new isoprenylated flavans, kazinols E and H, and five new isoprenylated 1,3-diphenylpropane derivatives, kazinols C, D, F, G, and K, were isolated from the extract of the root bark of *Broussonetia kazinoki* SIEB. (Japanese name, Himekōzo, Moraceae). On the basis of chemical and spectral evidence, the structures of kazinols E and H were shown to be **3** and **6**, respectively, and the structures of kazinols C, D, F, G, and K to be **1**, **2**, **4**, **5**, and **7**, respectively.

Keywords—*Broussonetia kazinoki*; Moraceae; flavan; 1,3-diphenylpropane; ¹H-NMR; ¹³C-NMR; nuclear Overhauser effect

Previously we reported the structure determination of a series of natural Diels–Alder type adducts as well as isoprenylated flavonoids isolated from the root bark of cultivated mulberry tree and the Chinese crude drug “Sang-Bai-Pi” (Japanese name “Sōhakuhi”), the latter imported from the People’s Republic of China.³⁾ Furthermore we reported the phenolic components of *Cudrania tricuspidata* (CARR.) BUR.⁴⁾ and *Broussonetia papyrifera* (L.) VENT.,^{2,5)} both of which belong to the family Moraceae. In the course of our studies on the phenolic constituents of Moraceae plants, we studied the root bark of *Broussonetia kazinoki* SIEB. (Japanese name “Himekōzo”), which is a deciduous tree distributed over Korea, China, Taiwan, and Japan. Its cortex has been used as a raw material for paper and also as a Chinese crude drug, as has the cortex of *B. papyrifera* (L.) VENT. (Japanese name “kazinoki”).⁶⁾ In this paper, we report the structure determination of two isoprenylated flavans, kazinols E and H, and of five isoprenylated 1,3-diphenylpropane derivatives, kazinols C, D, F, G, and K, obtained from the root bark of *B. kazinoki* SIEB.

The dried root bark of *B. kazinoki* SIEB. was extracted successively with *n*-hexane and benzene. The *n*-hexane extract was fractionated sequentially by column chromatography and preparative thin-layer chromatography (TLC) to give kazinols C (**1**), D (**2**), E (**3**), G (**5**), H (**6**), and K (**7**). A part of the benzene extract was also fractionated sequentially by preparative TLC to give kazinol F (**4**).

Kazinol C (**1**) was obtained as an oily substance, $M^+ = 464.2924$, $C_{30}H_{40}O_4$, exhibiting a positive ferric chloride test and sodium molybdate test.⁷⁾ The infrared (IR) spectrum of **1** suggested the presence of hydroxyl groups [3650 (sh), 3600 (sh), 3530 (br) cm^{-1}] and aromatic rings [1625, 1600 (sh) cm^{-1}]. Treatment of **1** with dimethyl sulfate and potassium carbonate in acetone gave a tetramethyl ether (**1a**). Acetylation of **1** with acetic anhydride in pyridine gave a triacetate (**1b**) and a tetraacetate (**1c**). The ultraviolet (UV) spectrum of **1** showed absorption maxima at 218 ($\log \epsilon = 4.59$) and 286 nm (3.97), which indicated the presence of an unconjugated aromatic system.^{2,8)} The proton nuclear magnetic resonance (¹H-NMR)




Fig. 1

spectrum of **1** indicated the presence of two 3,3-dimethylallyl groups, a 1,1-dimethylallyl group, and three aromatic protons: δ 1.66, 1.70, 1.72, 1.78 (each 3H, s), 3.23 (2H, d, $J=7$ Hz), 3.34 (2H, d, $J=7$ Hz), 4.96 (1H, t, $J=7$ Hz), 5.12 (1H, t, $J=7$ Hz); δ 1.38 (6H, s), 5.25 (1H, d, $J=10$ Hz), 5.31 (1H, d, $J=18$ Hz), 6.16 (1H, dd, $J=10$ and 18 Hz); δ 6.30 (1H, s), 6.61 (1H, s), 6.93 (1H, s). The 1H -NMR spectrum also indicated the presence of a 1,3-disubstituted propane moiety: δ 1.82 (2H, quintet, $J=7$ Hz), 2.55 (2H, t, $J=7$ Hz), 2.59 (2H, t, $J=7$ Hz). The mass spectrum (MS) of **1** showed significant fragments⁹⁾ at m/z 408 ($M^+ - C_4H_8$), 396 ($M^+ - C_5H_8$), 340 ($396 - C_4H_8$), 273 ($M^+ - 191$), 259 (**8**), 245 (**9**), 205 ($M^+ - 259$), 203 ($259 - C_4H_8$), 191 (**10**), 123 ($191 - C_5H_8$). From the above results, it was suggested that kazinol C is a 1,3-diphenylpropane derivative having two isoprenyl groups and two hydroxyl groups in one of the phenyl moieties, and an isoprenyl group and two hydroxyl groups in the other




Fig. 2. NOE Results for Kazinol C (CDCl_3)
%: Increase of signal area.




Fig. 3. LSPD Measurement of Kazinol C

- a) $^{13}\text{C}\{-^1\text{H}\}$ Complete decoupling.
- b) $^{13}\text{C}\{-^1\text{H}\}$ Non-decoupling.
- c) Irradiated at δ 3.3 ($\text{C}_{7''}\text{-H} \times 2$ and $\text{C}_{12''}\text{-H} \times 2$).
- d) Irradiated at δ 6.61 ($\text{C}_{2''}\text{-H}$).

phenyl moiety. In the carbon-13 nuclear magnetic resonance (^{13}C -NMR) studies, the carbon atoms of **1** were assigned by the off-resonance decoupling technique as well as by comparing the ^{13}C -NMR spectrum of **1** with those of model compounds^{2,10)} and by the long-range selective ^1H decoupling (LSPD) technique (Table I). In the ^{13}C -NMR spectrum of **1**, the chemical shift values of the carbon atoms at the C-3'' and -4'' positions were similar to those of the corresponding carbon atoms of 3',4'-dioxxygenated flavonoids.^{2,11)} Based on the biogenetic analogy to the 1,3-diphenylpropane derivatives obtained from the plants of *Broussonetia* species and the spectral data (MS and ^1H -NMR),^{2,12)} the A ring of **1** seems to be a 2',4'-dihydroxy-5'-isoprenylated phenyl moiety. The substitution pattern of the A and B rings was also supported by the nuclear Overhauser effects (NOE), which were observed as follows: (1) at the C-3 protons (δ 2.55) when the aromatic proton at C-2'' (δ 6.61) was irradiated, (2) at the C-12'' protons (δ 3.23) and the C-2'' proton when the C-3 protons were irradiated, (3) between the protons at C-12'' and those at the C-7'' (δ 3.34) and C-3 positions when the C-12'' protons (δ 3.23) were irradiated, (4) between the protons at C-7'' and those at C-12'' when the C-7'' protons (δ 3.34) were irradiated, and (5) between the C-1 protons (δ 2.59) and the C-6' proton (δ 6.93) when the C-1 protons (δ 2.59) were irradiated (Fig. 2). These results suggest that the B ring is a 3'',4''-dihydroxy-5'',6''-bis(3,3-dimethylallyl)phenyl moiety and that the A ring is a 2',4'-dihydroxy-5'-(1,1-dimethylallyl)phenyl moiety. The substitution pattern of the B ring was further supported by the result of the LSPD technique and the chemical shift values of methoxyl carbons of the tetramethyl ether (**1a**). The LSPD technique yielded the following observations. When the C-2'' proton (δ 6.61) was weakly irradiated, the signals of the C-3'' (δ 141.7) and C-4'' (δ 140.5) carbon atoms changed. When the C-12'' (δ 3.23) and C-7'' (δ 3.34) protons were weakly irradiated, the C-4'' carbon signal changed, while the C-2'' (δ 114.2) and C-3'' carbon signals showed no change (Fig. 3). On the other hand, it was reported that the signal of the di-*ortho*-substituted methoxyl carbon nucleus appears at δ ca. 60 ppm, while that of the mono-*ortho*-substituted methoxyl carbon nucleus appears at ca. 55 ppm.¹³⁾ In the case of **1a**, the signals of

TABLE I. ^{13}C -NMR Chemical Shifts (ppm) 1—7

	1	2	4	5	7		3	6
C-1	32.9 ^{a)}	32.7 ^{a)}	33.6 ^{a)}	33.1 ^{a)}	32.6 ^{a)}	C-2	75.0	74.8
C-2	29.7	29.7	30.4	29.7	29.5	C-3	25.6	25.7
C-3	31.6 ^{a)}	31.7 ^{a)}	33.1 ^{a)}	31.8 ^{a)}	31.4 ^{a)}	C-4	29.9	29.8
C-1'	120.7	120.5	120.5	120.0	119.5	C-4a	113.9	113.3
C-2'	153.2	153.7 ^{b)}	157.3 ^{b)}	153.6 ^{b)}	152.8 ^{b)}	C-5	127.0	127.0
C-3'	105.0	104.8	103.4	104.9	104.3	C-6	125.1	124.9
C-4'	153.2	153.3 ^{b)}	156.6 ^{b)}	153.1 ^{b)}	152.6 ^{b)}	C-7	153.5 ^{a)}	153.9 ^{a)}
C-5'	124.4	124.0	107.3	124.5	123.7	C-8	105.6	105.8
C-6'	127.7	127.6	131.1	127.6	127.0	C-8a	155.1 ^{a)}	155.4 ^{a)}
C-1''	133.0	132.0	132.7	131.2	132.3	C-1'	131.0	131.3
C-2''	114.2	113.1	114.6	111.1	115.0	C-2'	111.3	112.9
C-3''	141.7	143.0	143.0	143.7	141.9	C-3'	142.4	143.3
C-4''	140.5	139.2	142.2	142.2	137.0	C-4'	141.7	139.3
C-5''	130.5	119.8	130.4	130.7	119.2	C-5'	130.2	119.8
C-6''	127.0	128.9	127.7	126.5	126.2	C-6'	127.0	126.4
C-7'	39.8	39.8		39.8	39.6	C-9	39.8	39.9
C-8'	27.3	27.3		27.3	27.1	C-10	27.2	27.4
C-9'	27.3	27.3		27.3	27.1	C-11	27.2	27.4
C-10'	148.5	148.6		148.3	147.8	C-12	148.3	148.5
C-11'	113.1	112.7		113.1	112.5	C-13	113.0	113.8
C-7''	27.8 ^{b)}	20.3	28.1 ^{c)}	27.7 ^{c)}	119.6	C-14	27.2 ^{b)}	120.0
C-8''	122.7 ^{c)}	33.2	124.9 ^{d)}	123.3 ^{d)}	129.8	C-15	122.8 ^{c)}	130.8
C-9''	133.3 ^{d)}	74.0	130.7 ^{e)}	131.5 ^{e)}	75.5	C-16	132.4 ^{d)}	76.3
C-10''	18.0	26.6	18.0	17.9 ^{f)}	27.5	C-17	17.9	27.7
C-11''	25.7	26.6	25.8	25.7	27.5	C-18	25.6	28.1
C-12''	26.1 ^{b)}	27.3	26.2 ^{c)}	25.6 ^{c)}	26.7	C-19	25.6 ^{b)}	26.5
C-13''	124.4 ^{c)}	123.6	125.8 ^{d)}	124.4 ^{d)}	123.6	C-20	124.1 ^{c)}	123.9
C-14''	130.9 ^{d)}	131.0	130.7 ^{e)}	131.2 ^{e)}	130.2	C-21	131.4 ^{d)}	132.2
C-15''	18.0	17.9	18.0	18.0 ^{f)}	17.9	C-22	17.9	18.2
C-16''	25.7	25.7	25.8	25.7	25.5	C-23	25.6	25.7
Solv.	A	A	B	A	A		A	A

Solvent: A, CDCl_3 ; B, acetone- d_6 . a—f) Assignments may be interchanged in each column.

TABLE II. Acetylation Shifts for C_3 -, C_6 -, C_{10} -, C_{11} - and C_2 -H of **1b** and **1c**

	C_3 -H	C_6 -H	C_{10} -H	C_{11} -H		C_2 -H
1b	6.50	7.10	6.20	5.33	5.38	6.90
1c	6.79	7.26	5.99	5.00	5.03	6.91
Δ	-0.23	-0.16	+0.21	+0.33	+0.35	-0.01

Measured in CDCl_3 (ppm).

methoxyl carbons appeared at δ 55.4 ($\times 2$), 55.7, and 60.6 ppm. This results suggests that one of the methoxyl groups is di-*ortho*-substituted. The substitution pattern of the A ring was further supported by observation of the acetylation shift of the A ring proton signals of **1b** and **1c**. Comparison of the ^1H -NMR spectra of **1**, **1b**, and **1c** indicates that the acetylation of the hydroxyl group of **1b** caused a remarkable downfield shift of the proton at C-3' position and an upperfield shift of the protons at C-10' and -11' (Table II). This result suggests that the 1,1-dimethylallyl group is located at the C-5' position. From the above results, the formula **1** was suggested for the structure of kazinol C.

The formula (1) was further supported by the following results. A mixture of **1** and hydrochloric acid in methanol was kept at 60 °C for 2 h. The reaction mixture was purified by preparative TLC to give **1d**, **1e**, **1f**, and **1g**. Compound **1e** was identical with kazinol D, and the structures of the reaction products were supported by the spectral data. The MS of **1d** gave the MS which showed the molecular ion peak at m/z 464. The ^1H -NMR spectrum of **1d** indicated the presence of a 3,3-dimethylallyl group and a 2,2-dimethyldihydropyran ring system, δ 1.70 (6H, s), 3.17 (2H, d, $J=7$ Hz), 4.92 (1H, t, $J=7$ Hz); δ 1.32 (6H, s), 1.80 (2H, t, $J=7$ Hz), 2.45—2.75 (2H, m), as well as a 1,3-diphenylpropane moiety, δ 1.60—1.90 (2H, m), 2.58 (2H, t, $J=7$ Hz), 2.64 (2H, t, $J=7$ Hz). The presence of a 2-methyl-3,3-dimethyl-5,6-disubstituted dihydrobenzofuran moiety¹⁴⁾ and an aromatic proton was indicated: δ 1.25 (3H, s), 1.38 (3H, d, $J=7$ Hz), 1.66 (3H, s), 4.30 (1H, q, $J=7$ Hz), 6.21 (1H, s), 6.75 (1H, s); δ 6.62 (1H, s). From the above results, the structure (**1d**) was suggested. The MS of **1f** showed the molecular ion peak at m/z 496. The ^1H -NMR of **1f** showed the presence of a 1,1-dimethylallyl group, a 2,2-dimethyldihydropyran ring system, and a 3-methyl-3-methoxybutyl moiety: δ 1.42 (6H, s), 5.26 (1H, d, $J=10$ Hz), 5.29 (1H, d, $J=18$ Hz), 6.16 (1H, dd, $J=10$ and 18 Hz); δ 1.34 (6H, s), 1.82 (2H, t, $J=7$ Hz), 2.30—2.80 (8H, m, overlapping the signals of the protons at C-1, -3, -7'', and -12''); δ 1.23 (6H, s), 1.50—1.95 (4H, m, overlapping the signals of protons at C-2 and -13''), 3.26 (3H, s). From the above results, the structure **1f** was suggested. The MS of **1g** showed the molecular ion peak at m/z 496. The ^1H -NMR spectrum indicated the presence of a 3,3-dimethylallyl group, a 1,1-dimethylallyl group, and a 3-methyl-3-methoxybutyl moiety: δ 1.67, 1.72 (each 3H, s), 3.21 (2H, d, $J=7$ Hz), 4.96 (1H, t, $J=7$ Hz); δ 1.39 (6H, s), 5.24 (1H, d, $J=10$ Hz), 5.29 (1H, d, $J=18$ Hz), 6.15 (1H, dd, $J=10$ and 18 Hz); δ 1.20 (6H, s), 1.60—1.90 (4H, m, overlapping the signals of protons at C-2 and -8'' (or -13'')), 2.40—2.80 (6H, m, overlapping the signals of protons at C-1, -3, and -7'' (or -12'')), 3.28 (3H, s). From these results, two possible structures (**1g** and **1g'**) were suggested. Comparison of the chemical shift values of the 3,3-dimethylallyl group of **1g** and the values of the relevant groups of **1**, **1d**, and **2** (**1e**) indicated that the structure **1g** seemed more probable than **1g'**. The ^1H -NMR and IR spectra of the flavan compound **3**, which was obtained by treatment of **1** with 2,3-dichloro-5,6-dicyanobenzoquinone (DDQ), were identical with those of kazinol E (**3**). All these results indicate that the structure of kazinol C is represented by formula **1**.


Kazinol D (**2**) was obtained as an oily substance, $M^+ = 464.2927$, $\text{C}_{30}\text{H}_{40}\text{O}_4$, exhibiting a positive ferric chloride test, but a negative sodium molybdate test.⁷⁾ The IR spectrum of **2** suggested the presence of hydroxyl groups [3615 , 3570 , 3500 cm^{-1}] and aromatic rings [1625 , 1600 cm^{-1}]. Treatment of **2** with dimethyl sulfate and potassium carbonate in acetone gave a trimethyl ether (**2a**). The UV spectrum of **2** showed absorption maxima at 221 (4.67), 230 (infl. 4.07), and 283 nm (3.91). As described in the experimental section, the ^1H -NMR spectrum of **2** indicated the presence of a 3,3-dimethylallyl group, a 1,1-dimethylallyl group, and a 2,2-dimethyldihydropyran ring system. The spectrum also indicated the presence of three aromatic protons: δ 6.28 (1H, s), 6.63 (1H, s), 6.91 (1H, s). The MS of **2** showed significant fragments at m/z 259, 245, and 191 (**10**),⁹⁾ and the spectrum was similar to that of **1**. These results indicated that kazinol D is a 1,3-diphenylpropane derivative having a 3,3-dimethylallyl group, a 1,1-dimethylallyl group, and a 2,2-dimethyldihydropyran ring system. In the ^{13}C -NMR spectrum of **2**, the chemical shift values of the carbon atoms of the 1,3-diphenylpropane skeleton as well as of a 3,3-dimethylallyl group and a 1,1-dimethylallyl group were similar to those of the relevant carbon atoms of **1** except that of the carbon atom at C-5''. In the ^{13}C -NMR spectrum of **2a**, the signals of the methoxyl carbon atoms appeared at δ 55.4 ($\times 2$) and 56.0 ppm. This result suggests that all the methoxyl groups of **2a** are mono-*ortho*-substituted.¹³⁾ Kazinol D was derived from kazinol C by treatment with hydrochloric acid solution. All these results indicated that the structure of kazinol D is represented by formula **2**.

Kazinol E (**3**) was obtained as colorless needles, mp 147 °C, $M^+ = 462.2803$, $\text{C}_{30}\text{H}_{38}\text{O}_4$,

$[\alpha]_D^{18} + 0.30^\circ$, exhibiting a positive ferric chloride test and sodium molybdate test.⁷⁾ The IR spectrum of **3** indicated the presence of hydroxyl groups [3590 (sh), 3550, 3490 cm^{-1}] and aromatic rings [1625, 1585 cm^{-1}]. The UV spectrum showed absorption maxima at 224 (4.92), 232 (infl. 4.36), and 288 nm (3.91).⁸⁾ As described in the experimental section, the ^1H -NMR spectrum of **3** showed the presence of two 3,3-dimethylallyl groups and a 1,1-dimethylallyl group. The spectrum also showed five aliphatic proton signals and three aromatic proton signals: δ 1.70—2.10 (2H, m), 2.62—2.88 (2H, m), 4.80—5.10 (1H, m); δ 6.31 (1H, s), 6.73 (1H, s), 6.83 (1H, s). The MS of **3** showed significant fragments at m/z 272 (**11**) and 191 (**10**).¹⁵⁾ In the ^{13}C -NMR spectrum of **3**, the carbon atoms were assigned by the off-resonance decoupling technique as well as by comparing the ^{13}C -NMR spectrum of **3** with those of model compounds (**1**, **2**, kazinol A²⁾ and B²⁾) (Table I). ($\pm$)-Kazinol E was derived from **1** by treatment with DDQ. On the basis of the circular dichroism (CD) spectrum, **3** was concluded to have the (*S*)-configuration at the C-2 position.¹⁶⁾ From the above results, kazinol E is represented by the formula **3**.

Kazinol F (**4**) was obtained as colorless needles, mp 108—109°C, $M^+ = 396.2301$, $\text{C}_{25}\text{H}_{32}\text{O}_4$, exhibiting a positive ferric chloride test, sodium molybdate test,⁷⁾ and Gibbs test. The IR spectrum of **4** suggested the presence of hydroxyl groups and aromatic rings. Treatment of **4** with dimethyl sulfate and potassium carbonate in acetone gave a tetramethyl ether (**4a**). The UV spectrum showed absorption maxima at 282 (3.91), 286 nm (sh 3.89). As described in the experimental section, the ^1H -NMR spectrum of **4** indicated the presence of two 3,3-dimethylallyl groups and a 1,3-diphenylpropane moiety. The ^1H -NMR spectrum also indicated the presence of ABC-type aromatic protons, and an aromatic proton: δ 6.28 (1H, dd, $J = 2$ and 8 Hz), 6.39 (1H, d, $J = 2$ Hz), 6.88 (1H, d, $J = 8$ Hz); δ 6.57 (1H, s). The MS of **4** showed significant fragments⁹⁾ at m/z 259 (**8**), 245 (**9**), and 123 (**12**). In the ^{13}C -NMR spectrum of **4**, the chemical shift values of the carbon atoms of the 1,3-diphenylpropane skeleton as well as of two 3,3-dimethylallyl groups were similar to those of the relevant carbon atoms of **1** except those of the carbon atoms at C-2', -4', -5', and -6' (Table I). In the ^{13}C -NMR spectrum of **4a**, the signals of the methoxyl carbon atoms appeared at δ 55.0, 55.1, 55.5, and 60.3 ppm, suggesting that one of the four methoxyl groups is di-*ortho*-substituted.¹³⁾ The substitution pattern of the B ring of **4** was supported by the NOE: (1) at the C-3 protons (δ 2.47) when the aromatic proton at C-2'' (δ 6.58) was irradiated, (2) between the protons at C-12'' and those at C-13'' (δ 4.96), C-3 and C-7'' when C-12'' protons (δ 3.21) were irradiated, and (3) between the protons at C-7'' and those at C-12'', C-13'' and C-8'' (δ 5.09) when the C-7'' protons (δ 3.34) were irradiated (Fig. 4). These results suggest that the B ring is a 3'',4''-dihydroxy-5'',6''-bis(3,3-dimethylallyl)phenyl moiety. When kazinol F was treated with DDQ in dry ether, a flavan derivative (**13**) was obtained, the ^1H -NMR and IR spectra of which were identical with those of kazinol I isolated from *Broussonetia papyrifera* (L.) VENT.¹⁷⁾ From the above results, kazinol F is represented by formula **4**.

Kazinol G (**5**) was obtained as an oily substance, $M^+ = 532.3536$, $\text{C}_{35}\text{H}_{48}\text{O}_4$, exhibiting a positive ferric chloride test, but a negative sodium molybdate test.⁷⁾ The IR spectrum of **5** suggested the presence of hydroxyl groups and aromatic rings. The UV spectrum showed absorption maxima at 221 (sh 4.49) and 285 nm (3.88), similar to those of **1**. The ^1H -NMR spectrum of **5** indicated the presence of two 3,3-dimethylallyl groups, a 3,3-dimethylallyloxy moiety, a 1,1-dimethylallyl group, three aromatic protons, and 1,3-diphenylpropane moiety: δ 1.66 (6H, s), 1.70, 1.71 (each 3H, s), 3.24 (2H, d, $J = 6$ Hz), 3.36 (2H, d, $J = 6$ Hz), 4.98 (1H, t, $J = 6$ Hz), 5.11 (1H, t, $J = 6$ Hz); δ 1.74, 1.78 (each 3H, s), 4.51 (2H, d, $J = 7$ Hz), 5.47 (1H, t, $J = 7$ Hz); δ 1.36 (6H, s), 5.27 (1H, d, $J = 11$ Hz), 5.32 (1H, d, $J = 18$ Hz), 6.16 (1H, dd, $J = 11$ and 18 Hz); δ 6.30 (1H, s), 6.59 (1H, s), 6.94 (1H, s); δ 1.84 (2H, dd, $J = 7$ and 8 Hz), 2.58 (2H, t, $J = 8$ Hz), 2.60 (2H, t, $J = 7$ Hz). The ^1H -NMR spectrum also indicated the presence of three protons of hydroxyl groups as follows: δ 4.61, 5.62, 5.72 (each 1H, br s). The MS of **5** showed



The chemical structure shows a complex polycyclic molecule with several substituents and coupling constants. The structure includes a central ring system with various functional groups and substituents. The following table lists the chemical shifts and coupling constants for the protons in the structure:

Proton Label	Chemical Shift (ppm)	Coupling Constant (Hz)
1.36	1.36	21%
1.84	1.84	3%
2.60	2.60	8%
2.58	2.58	10%
3.24	3.24	2%
3.36	3.36	3%
6.59	6.59	6%
4.51	4.51	4%

significant fragments⁹) at m/z 259 (9) and 191 (10). In the ¹³C-NMR spectrum of **5**, the chemical shift values of the carbon atoms of the 1,3-diphenylpropane skeleton as well as of two 3,3-dimethylallyl groups and a 1,1-dimethylallyl group were similar to those of the relevant carbon atoms of **1** except those of the carbon atoms at C-1'', -2'', -3'', and -4''. These results suggest that the structure of kazinol G is represented by the formula **5**. This assumption was further supported by the NOE: (1) at the C-17'' protons (δ 4.51) when the aromatic proton at C-2'' (δ 6.59) was irradiated, (2) at the C-2'' and C-12'' (δ 3.24) protons when the C-3 protons (δ 2.58) were irradiated, (3) at the C-7'' protons (δ 3.36) when the C-12'' protons were irradiated, (4) between the protons at C-7'' and those at C-12'' when the C-7'' protons (δ 3.36) were irradiated, (5) between the protons at C-2 (δ 1.84), that at C-6' (δ 6.94) and that at C-2'' when the C-2 protons (δ 1.84) were irradiated, (6) between the C-1 protons (δ 2.60) and the C-6' proton when the C-1 protons (δ 2.60) were irradiated, (7) between the C-6' proton, the C-1 protons and the methyl protons at C-7' (δ 1.36) when the C-6' proton (δ 6.94) was irradiated, and (8) between the C-7' methyl protons and the C-6' proton when the C-7' methyl protons (δ 1.36) were irradiated (Fig. 5). All these results indicated that the structure of kazinol G is represented by **5**.

Kazinol K (**7**) was obtained as an oily substance, $M^+ = 462.2778$, $C_{30}H_{38}O_4$, exhibiting a positive ferric chloride test, but a negative sodium molybdate test.⁷⁾ The IR spectrum of **7** suggested the presence of hydroxyl groups and aromatic rings. Treatment of **7** with dimethyl sulfate and potassium carbonate in acetone gave a trimethyl ether (**7a**). The UV spectrum was similar to that of **6**, showing absorption maxima at 226 (4.45), 266 (sh 4.00), 277 (4.18), 285 (4.16), and 324 nm (3.34). As described in the experimental section, the 1H -NMR spectrum of




Fig. 6. NOE Results for Kazinol K (CDCl_3)
%: Increase of signal area.

7 indicated the presence of a 3,3-dimethylallyl group, a 1,1-dimethylallyl group, a 2,2-dimethylpyran ring system, 1,3-diphenylpropane moiety, and three aromatic protons. The MS of 7 showed a significant fragment at m/z 191 (10).⁹ In the ^{13}C -NMR spectrum of 7 the chemical shift values of the carbon atoms of a 1,3-diphenylpropane skeleton as well as of a 3,3-dimethylallyl group and a 1,1-dimethylallyl group were similar to those of the relevant carbon atoms of 1 except those of the carbon atoms at C-4'' and -5''. In the ^{13}C -NMR spectrum of 7a, the signals of the methoxyl carbon atoms appeared at δ 55.5 ($\times 2$), and 56.4 ppm. This result suggests that all the methoxyl group of 7a are mono-*ortho*-substituted.¹³ The substitution pattern of the B ring of 7 was supported by the NOE: (1) at the C-3 protons (δ 2.55) when the aromatic proton at C-2'' (δ 6.67) was irradiated, (2) at the C-3, C-13'' (δ 4.97), and C-7'' (δ 6.49) protons when the C-12'' protons (δ 3.24) were irradiated, and (3) between the C-7'' proton and the C-12'', C-13'', and C-8'' (δ 5.62) protons when the C-7'' proton (δ 6.49) was irradiated (Fig. 6). The IR and ^{13}C -NMR spectra of the flavan derivative (6), which was obtained by treatment of 7 with DDQ, were identical with those of kazinol H. From the above results, kazinol K is represented by the formula 7.

It is noteworthy that the new compounds described in this paper have a unique substitution pattern in the B ring, that is a 3,4-dihydroxy-5,6-bis(3,3-dimethylallyl)phenyl moiety, and also that kazinol C (1) and two other compounds (3, 5) have two 3,3-dimethylallyl groups and a 1,1-dimethylallyl group in the molecule.

Experimental

All melting points are uncorrected. ^1H -NMR and ^{13}C -NMR spectra were measured with tetramethylsilane (TMS) as an internal reference. Chemical shifts are expressed in ppm downfield from TMS and coupling constants (J) in Hz. Abbreviations: s=singlet, d=doublet, t=triplet, q=quartet, m=multiplet, br=broad, sh=shoulder, infl.=inflection. The following instruments were used: melting points, Yazawa micromelting point apparatus (hot-stage type); UV spectra, Hitachi 340 UV spectrometer; IR spectra, Hitachi 260-30 IR spectrometer; ^1H -NMR spectra, Hitachi R-900 FT NMR spectrometer, JEOL JNM 4H-100 and GX-400 FT NMR spectrometers and Varian XL-200 FT NMR spectrometer; ^{13}C -NMR spectra, JEOL GX-270, GX-400, and Hitachi R-900 FT NMR spectrometers; optical rotation, JASCO DIP-4; CD spectra, JASCO J-20 ORD spectrometer. For TLC and preparative TLC, Wakogel B-5FM and B-5F were used, and for column chromatography, Wakogel C-200.

Isolation of Kazinols C (1), D (2), E (3), F (4), G (5), H (6), and K (7)—The dried root bark (1.0 kg) of *Broussonetia kazinoki* SIEB., collected in the vicinity of Shimokawara, Moroyama Town, Saitama Prefecture in May 1984, was finely cut and extracted with *n*-hexane and then with benzene. Evaporation of the *n*-hexane extract and the benzene extract to dryness yielded 28.0 g and 44.0 g of residue, respectively. The *n*-hexane extract (28.0 g) was chromatographed on silica gel (230 g) with benzene- $(\text{CH}_3)_2\text{CO}$ as an eluent, each fraction being monitored by TLC. One of the fractions eluted with benzene (1370 mg) was fractionated by preparative TLC (solvent system, CHCl_3 : Et_2O =12:1, *n*-hexane: Et_2O =1:1) to give kazinol E (3, 650 mg). One of the fractions eluted with benzene (1314 mg) was fractionated by preparative TLC (*n*-hexane: Et_2O =2:1, AcOEt : CHCl_3 =2:50, AcOEt :*n*-hexane=1:3) to give kazinol G (5, 52 mg). One of the fractions eluted with benzene (746 mg) was fractionated by preparative TLC (*n*-hexane: Et_2O =2:1, AcOEt :*n*-hexane=1:3, CHCl_3 only) to give kazinol H (6, 17 mg). The fractions eluted with benzene containing 1% $(\text{CH}_3)_2\text{CO}$ were evaporated to give the residue (1.68 g), which was fractionated by preparative TLC ($(\text{CH}_3)_2\text{CO}$:*n*-hexane=1:3, *n*-hexane: Et_2O =1:1, CHCl_3 : Et_2O =4:1) to give kazinol K (7, 176 mg). One of the fractions eluted with benzene containing 3% $(\text{CH}_3)_2\text{CO}$ (1728 mg) was fractionated by preparative TLC (benzene: Et_2O =3:1, CHCl_3 : Et_2O =5:1, benzene: AcOEt =4:1) to give kazinol C (1, 420 mg).

The same fraction (1728 mg) was also fractionated by preparative TLC (benzene:Et₂O=3:1, CHCl₃ only, CHCl₃:AcOEt=3:1) to give kazinol D (2, 370 mg). A part of the benzene extract (2.1 g) was fractionated by preparative TLC (benzene:Et₂O=3:1, *n*-hexane:Et₂O=1:3, CHCl₃:(CH₃)₂CO=5:2) to give kazinol F (4, 2 mg).

Kazinol C (1)—Compound 1 was obtained as an oily substance. FeCl₃ test (brown), Na₂MoO₄ test (orange). UV $\lambda_{\text{max}}^{\text{EtOH}}$ nm (log ϵ): 218 (4.59), 286 (3.97). IR $\nu_{\text{max}}^{\text{CHCl}_3}$ cm⁻¹: 3650 (sh), 3600 (sh), 3530 (br), 1640 (sh), 1635 (sh), 1625, 1600 (sh). High-resolution MS, Calcd for C₃₀H₄₀O₄ (M⁺), *m/z* 464.2925. Found: *m/z* 464.2924. MS: *m/z* 464 (M⁺), 408 (M⁺ - C₄H₈), 396 (M⁺ - C₅H₈), 340 (396 - C₄H₈), 273 (M⁺ - 191), 259 (8), 245 (9), 205 (M⁺ - 259), 203 (259 - C₄H₈), 191 (10), 123 (191 - C₅H₈). ¹H-NMR (CDCl₃, 100 MHz) δ : 1.38 (6H, s, C₇-CH₃ × 2), 1.66, 1.70, 1.72, 1.78 (each 3H, s, C₉- and C₁₄-CH₃), 1.82 (2H, quintet, *J*=7, C₂-H × 2), 2.55 (2H, t, *J*=7, C₃-H × 2), 2.59 (2H, t, *J*=7, C₁-H × 2), 3.23 (2H, d, *J*=7, C₁₂-H × 2), 3.34 (2H, d, *J*=7, C₇-H × 2), 4.96 (1H, t, *J*=7, C₁₃-H), 5.12 (1H, t, *J*=7, C₈-H), 5.25 (1H, d, *J*=10, C₁₁-H), 5.31 (1H, d, *J*=18, C₁₁-H), 5.33 (2H, brs, OH × 2, disappeared on addition of D₂O), 5.66, 5.78 (each 1H, brs, OH, disappeared on addition of D₂O), 6.16 (1H, dd, *J*=10, 18, C₁₀-H), 6.30 (1H, s, C₃-H), 6.61 (1H, s, C₂-H), 6.93 (1H, s, C₆-H). ¹³C-NMR (22.6 MHz): Table I.

Kazinol C Tetramethyl Ether (1a)—A mixture of 1 (101 mg), (CH₃)₂SO₄ (0.5 ml) and K₂CO₃ (5 g) in (CH₃)₂CO (30 ml) was refluxed for 1 h, and treated as usual. The product was purified by preparative TLC (Et₂O:*n*-hexane=1:5) to give the tetramethyl ether (1a, 40 mg). The compound 1a was obtained as an oily substance. FeCl₃ test (negative). MS: *m/z* 520 (M⁺). ¹³C-NMR (CDCl₃, 22.6 MHz) δ : 18.0 (C₁₀- and C₁₅-), 25.7 (C₁₁- and C₁₆-), 25.9 (C₁₂-), 27.5 (C₈- and C₉-), 27.7 (C₇-), 30.2 (C₂), 31.8 (C₁), 33.5 (C₃), 40.0 (C₇-), 55.4 (OCH₃ × 2), 55.7 (OCH₃), 60.6 (OCH₃), 96.8 (C₃-), 109.4 (C₁₁-), 111.4 (C₂-), 121.8 (C₁-), 124.0 (C₈-), 124.3 (C₁₃-), 128.2 (C₅-), 128.4 (C₆-), 130.7 (C₅-), 130.9 (C₆- and C₁₄-), 134.5 (C₉-), 136.8 (C₁-), 145.5 (C₃-), 148.6 (C₁₀-), 150.6 (C₄-), 156.6 (C₂-), 157.3 (C₄-).

Acetylation of Kazinol C (1)—Kazinol C (1, 30 mg) was acetylated with Ac₂O (1.5 ml) and pyridine (0.5 ml) at room temperature for 3 min. The reaction mixture was treated as usual, and the product was purified by preparative TLC (Et₂O:*n*-hexane=1:1) to give a triacetate (1b, 19.0 mg) and a tetraacetate (1c, 11.8 mg). Kazinol C triacetate (1b) was obtained as an oily substance. MS: *m/z* 590 (M⁺). ¹H-NMR (CDCl₃, 200 MHz) δ : 1.41 (6H, s, C₇-CH₃ × 2), 1.66 (12H, s, C₉-CH₃ × 2 and C₁₄-CH₃ × 2), 1.84 (2H, quintet, *J*=7, C₂-H × 2), 2.23, 2.26, 2.28 (each 3H, s, OAc), 2.53 (2H, t, *J*=7, C₁-H × 2), 2.61 (2H, t, *J*=7, C₃-H × 2), 3.26 (2H, d, *J*=7, C₁₂-H × 2), 3.29 (2H, d, *J*=7, C₇-H × 2), 4.99 (2H, t, *J*=7, C₈- and C₁₃-H), 5.33 (1H, d, *J*=10, C₁₁-H), 5.38 (1H, d, *J*=18, C₁₁-H), 6.20 (1H, dd, *J*=10, 18, C₁₀-H), 6.56 (1H, s, C₃-H), 6.90 (1H, s, C₂-H), 7.10 (1H, s, C₆-H). Kazinol C tetraacetate (1c) was obtained as an oily substance. MS: *m/z* 632 (M⁺). ¹H-NMR (CDCl₃, 200 MHz) δ : 1.41 (6H, s, C₇-CH₃ × 2), 1.67 (6H, s, C₉- and C₁₄-CH₃), 1.70 (6H, s, C₉- and C₁₄-CH₃), 1.87 (2H, quintet, *J*=7, C₂-H × 2), 2.20, 2.23, 2.27, 2.28 (each 3H, s, OAc), 2.60 (2H, t, *J*=7, C₃-H × 2), 2.64 (2H, t, *J*=7, C₁-H × 2), 3.26 (2H, d, *J*=7, C₁₂-H × 2), 3.30 (2H, d, *J*=7, C₇-H × 2), 4.99 (2H, t, *J*=7, C₈- and C₁₃-H), 5.00 (1H, d, *J*=10, C₁₁-H), 5.03 (1H, d, *J*=18, C₁₁-H), 5.99 (1H, dd, *J*=10, 18, C₁₀-H), 6.79 (1H, s, C₃-H), 6.91 (1H, s, C₂-H), 7.26 (1H, s, C₆-H).

Acidic Treatment of Kazinol C (1) (Formation of Kazinol D (2) and Other Compounds from 1)—A mixture of 1 (52.0 mg) and 35% HCl (3 ml) in MeOH (15 ml) was kept at 60 °C for 2 h, and treated as usual. The product was purified by preparative TLC (*n*-hexane:AcOEt=2:1) to give 1d (4.5 mg), 1e (14.4 mg), 1f (9.6 mg), and 1g (13.0 mg). Compound 1d was obtained as an oily substance. MS: *m/z* 464 (M⁺), 259, 245, 205, 191. ¹H-NMR (CDCl₃, 100 MHz) δ : 1.25 (3H, s, C₇-CH₃), 1.32 (6H, s, C₉-CH₃ × 2), 1.38 (3H, d, *J*=7, C₁₀-CH₃), 1.66 (3H, s, C₇-CH₃), 1.70 (6H, s, C₁₄-CH₃ × 2), 1.80 (2H, t, *J*=7, C₈-H × 2), 1.60—1.90 (2H, m, C₂-H × 2), 2.45—2.75 (2H, m, C₇-H × 2), 2.58 (2H, t, *J*=7, C₃-H × 2), 2.64 (2H, t, *J*=7, C₁-H × 2), 3.17 (2H, d, *J*=7, C₁₂-H × 2), 4.30 (1H, q, *J*=7, C₁₀-H), 4.72 (1H, brs, OH), 4.92 (1H, t, *J*=7, C₁₃-H), 5.44 (1H, brs, OH), 6.21 (1H, s, C₃-H), 6.62 (1H, s, C₂-H), 6.75 (1H, s, C₆-H). Compound 1e thus obtained was identical with kazinol D (2) on the basis of IR and ¹H-NMR spectral comparisons. Compound 1f was obtained as an oily substance. MS: *m/z* 496 (M⁺), 464, 396, 259, 245, 217, 205, 191, 123. ¹H-NMR (CDCl₃, 100 MHz) δ : 1.23 (6H, s, C₁₄-CH₃ × 2), 1.34 (6H, s, C₉-CH₃ × 2), 1.42 (6H, s, C₇-CH₃ × 2), 1.50—1.95 (4H, m, C₂-H × 2 and C₁₃-H × 2), 1.82 (2H, t, *J*=7, C₈-H × 2), 2.30—2.80 (8H, m, C₁-H × 2, C₃-H × 2, C₇-H × 2, and C₁₂-H × 2), 3.26 (3H, s, C₁₄-OCH₃), 5.26 (1H, d, *J*=10, C₁₁-H), 5.29 (1H, d, *J*=18, C₁₁-H), 6.16 (1H, dd, *J*=10, 18, C₁₀-H), 6.28 (1H, s, C₃-H), 6.64 (1H, s, C₂-H), 6.92 (1H, s, C₆-H). Compound 1g was obtained as an oily substance. MS: *m/z* 496 (M⁺), 464, 259, 245, 217, 205, 191, 123. ¹H-NMR (CDCl₃, 100 MHz) δ : 1.20 (6H, s, C₉-CH₃ × 2), 1.39 (6H, s, C₇-CH₃ × 2), 1.67, 1.72 (each 3H, s, C₁₄-CH₃), 1.60—1.90 (4H, m, C₂-H × 2 and C₈-H × 2), 2.40—2.80 (6H, m, C₁-H × 2, C₃-H × 2, and C₇-H × 2), 3.21 (2H, d, *J*=7, C₁₂-H × 2), 3.28 (3H, s, C₉-OCH₃), 4.96 (1H, t, *J*=7, C₁₃-H), 5.24 (1H, d, *J*=10, C₁₁-H), 5.29 (1H, d, *J*=18, C₁₁-H), 5.50 (1H, brs, OH), 5.77 (2H, brs, OH × 2), 6.15 (1H, dd, *J*=10, 18, C₁₀-H), 6.29 (1H, s, C₃-H), 6.63 (1H, s, C₂-H), 6.92 (1H, s, C₆-H).

Treatment of Kazinol C (1) with DDQ [Formation of Kazinol E (3) from 1]—A mixture of 1 (9.8 mg) and DDQ (1.5 mg) in dry benzene (1.5 ml) was kept at room temperature for 18 h. The reaction product was purified by preparative TLC (*n*-hexane:Et₂O=1:1) to give an amorphous powder (3, 3.9 mg). Compound 3 thus obtained was identical with kazinol E on the basis of MS, IR, ¹H-NMR, and ¹³C-NMR spectral comparisons.

Kazinol D (2)—Compound 2 was obtained as an oily substance. FeCl₃ test (brown), Na₂MoO₄ test (negative). UV $\lambda_{\text{max}}^{\text{EtOH}}$ nm (log ϵ): 221 (4.67), 230 (infl. 4.07), 283 (3.91). IR $\nu_{\text{max}}^{\text{CHCl}_3}$ cm⁻¹: 3615, 3570, 3500, 1645, 1625, 1600. High-resolution MS, Calcd for C₃₀H₄₀O₄ (M⁺), *m/z* 464.2925. Found: *m/z* 464.2927. MS: *m/z* 464 (M⁺), 396 (M⁺ - C₅H₈), 340 (396 - C₄H₈), 273 (M⁺ - 191), 259, 245, 205 (M⁺ - 259), 203 (259 - C₄H₈), 191 (10), 123 (191 - C₅H₈). ¹H-

NMR (CDCl₃, 100 MHz) δ : 1.32 (6H, s, C₉-CH₃ × 2), 1.39 (6H, s, C₇-CH₃ × 2), 1.66, 1.71 (each 3H, s, C₁₄-CH₃), 1.60–1.90 (2H, m, C₂-H × 2), 1.80 (2H, t, $J=7$, C₈-H × 2), 2.45–2.75 (6H, m, C₁-H × 2, C₃-H × 2 and C₇-H × 2), 3.18 (2H, brd, $J=7$, C₁₂-H × 2), 4.92 (1H, brt, $J=7$, C₁₃-H), 5.23 (1H, d, $J=10$, C₁₁-H), 5.27 (1H, d, $J=18$, C₁₁-H), 6.14 (1H, dd, $J=10, 18$, C₁₀-H), 6.28 (1H, s, C₃-H), 6.63 (1H, s, C₂-H), 6.91 (1H, s, C₆-H). ¹³C-NMR (22.6 MHz): Table I.

Kazinol D Trimethyl Ether (2a)—A mixture of **2** (100 mg), (CH₃)₂SO₄ (0.5 ml) and K₂CO₃ (5 g) in (CH₃)₂CO (30 ml) was refluxed for 2 h, and treated as usual. The product was purified by preparative TLC (*n*-hexane:CHCl₃ = 1:1) to give kazinol D trimethyl ether (**2a**, 82 mg). Compound **2a** was obtained as an oily substance. FeCl₃ test (negative). MS: m/z 506 (M⁺). ¹³C-NMR (CDCl₃, 22.6 MHz) δ : 17.9 (C₁₅), 20.6 (C₇), 25.7 (C₁₆), 26.6 (C₁₀ and C₁₁), 27.5 (C₈, C₉, and C₁₂), 30.0 (C₂), 31.9 (C₃), 33.2 (C₁ and C₈), 39.9 (C₇), 55.4 (OCH₃ × 2), 56.0 (OCH₃), 73.0 (C₉), 96.7 (C₃), 109.4 (C₁₁), 110.9 (C₂), 120.4 (C₅), 121.8 (C₁), 123.5 (C₁₃), 128.1 (C₅), 128.4 (C₆), 129.8 (C₆), 131.0 (C₁₄), 131.3 (C₁), 141.8 (C₄), 146.8 (C₃), 148.5 (C₁₀), 156.5 (C₂), 157.2 (C₄).

Kazinol E (3)—Compound **3** was crystallized from CHCl₃ to give colorless needles, mp 147 °C, $[\alpha]_D^{18} +0.30^\circ$ ($c=0.41$, CHCl₃). FeCl₃ test (dark green). Na₂MoO₄ test (orange). UV $\lambda_{\max}^{\text{EtOH}}$ nm (log ϵ): 224 (4.92), 232 (infl. 4.36), 288 (3.91). IR $\nu_{\max}^{\text{CHCl}_3}$ cm⁻¹: 3590 (sh), 3550, 3490, 1625, 1585. High-resolution MS, Calcd for C₃₀H₃₈O₄ (M⁺), m/z 462.2768. Found: m/z 462.2803. MS: m/z 462 (M⁺), 406 (M⁺ - C₄H₈), 394 (M⁺ - C₅H₈), 272 (11), 271, 191 (10). ¹H-NMR (CDCl₃, 100 MHz) δ : 1.38 (6H, s, C₉-CH₃ × 2), 1.65 (9H, s, C₁₆-CH₃ × 2 and C₂₁-CH₃ or C₁₆-CH₃ and C₂₁-CH₃ × 2), 1.76 (3H, s, C₁₆-CH₃ or C₂₁-CH₃), 1.70–2.10 (2H, m, C₃-H × 2), 2.62–2.88 (2H, m, C₄-H × 2), 3.31 (4H, brd, $J=7$, C₁₄-H × 2 and C₁₉-H × 2), 4.80–5.10 (3H, m, C₂, C₁₅, and C₁₉-H), 5.17 (1H, d, $J=10$, C₁₃-H), 5.21 (1H, d, $J=18$, C₁₃-H), 5.60 (1H, brs, OH), 5.75 (2H, brs, OH × 2), 6.07 (1H, dd, $J=10, 18$, C₁₂-H), 6.31 (1H, s, C₈-H), 6.73 (1H, s, C₂-H), 6.83 (1H, s, C₅-H). ¹³C-NMR (22.6 MHz): Table I. CD spectrum: $[\theta]_{295} -99$, $[\theta]_{279} +66$ ($c=5.3 \times 10^{-4}$, MeOH).

Kazinol F (4)—Compound **4** was crystallized from benzene to give colorless needles, mp 108–109 °C. FeCl₃ test (dark green). Na₂MoO₄ test (orange). Gibbs test (blue). UV $\lambda_{\max}^{\text{EtOH}}$ nm (log ϵ): 282 (3.91), 286 (sh 3.89). IR $\nu_{\max}^{\text{KBr}}$ cm⁻¹: 3510, 3380, 3320 (sh), 1610. High-resolution MS, Calcd for C₂₅H₃₂O₄ (M⁺), m/z 396.2299. Found: m/z 396.2301. MS: m/z 396 (M⁺), 340 (M⁺ - C₄H₈), 259 (8), 245 (9), 123 (12). ¹H-NMR ((CD₃)₂CO, 400 MHz) δ : 1.65 (6H, s, C₉- and C₁₄-CH₃), 1.71, 1.73 (each 3H, s, C₉- or C₁₄-CH₃), 1.77 (2H, dd, $J=7, 8$, C₂-H × 2), 2.49 (2H, t, $J=8$, C₃-H × 2), 2.59 (2H, t, $J=7$, C₁-H × 2), 3.23 (2H, d, $J=6$, C₁₂-H × 2), 3.36 (2H, d, $J=7$, C₇-H × 2), 4.97 (1H, t, $J=6$, C₁₃-H), 5.10 (1H, t, $J=7$, C₈-H), 6.28 (1H, dd, $J=2, 8$, C₅-H), 6.39 (1H, d, $J=2$, C₃-H), 6.57 (1H, s, C₂-H), 6.88 (1H, d, $J=8$, C₆-H). ¹³C-NMR (22.6 MHz): Table I.

Kazinol F Tetramethyl Ether (4a)—A mixture of **4** (30 mg), (CH₃)₂SO₄ (0.2 ml), and K₂CO₃ (5 g) in (CH₃)₂CO (30 ml) was refluxed for 2 h, and treated as usual. The product was purified by preparative TLC (benzene only) to give kazinol F tetramethyl ether (**4a**, 19 mg). Compound **4a** was obtained as an oily substance. FeCl₃ test (negative). MS: m/z 452 (M⁺). ¹³C-NMR (CDCl₃, 22.5 MHz) δ : 17.9 (C₁₀ and C₁₅), 25.5 (C₁₁), 25.6 (C₁₆), 25.8 (C₁₂), 27.6 (C₇), 29.7 (C₂), 31.4 (C₃), 33.2 (C₁), 55.0, 55.1, 55.5, 60.3 (each OCH₃), 98.2 (C₃), 103.5 (C₅), 111.0 (C₂), 122.7 (C₁), 123.4 (C₈), 123.7 (C₁₃), 129.3 (C₅), 130.1 (C₆), 130.3 (C₆), 130.4 (C₁₄), 133.8 (C₉), 136.1 (C₁), 144.8 (C₃), 149.8 (C₄), 157.7 (C₂), 158.4 (C₄).

Treatment of Kazinol F (4) with DDQ—A mixture of **4** (20 mg) and DDQ (3 mg) in dry Et₂O (3 ml) was kept at room temperature for 18 h. The reaction product was purified by preparative TLC (*n*-hexane:(CH₃)₂CO = 2:1) to give **13**. Compound **13** thus obtained was identical with kazinol I isolated from *Broussonetia papyrifera* (L.) VENT. On the basis of MS, IR, ¹H-NMR spectral comparisons.

Kazinol G (5)—Compound **5** was obtained as an oily substance. FeCl₃ test (brown). Na₂MoO₄ test (negative). UV $\lambda_{\max}^{\text{EtOH}}$ nm (log ϵ): 221 (sh 4.49), 285 (3.88). IR $\nu_{\max}^{\text{CHCl}_3}$ cm⁻¹: 3600, 3540, 3480, 1625, 1610 (sh). High-resolution MS, Calcd for C₃₅H₄₈O₄ (M⁺), m/z 532.3536. Found: m/z 532.3536. MS: m/z 532 (M⁺), 464 (M⁺ - C₃H₈), 408 (464 - C₄H₈), 259 (8), 191 (10), 147, 123. ¹H-NMR (CDCl₃, 400 MHz) δ : 1.36 (6H, s, C₇-CH₃ × 2), 1.66 (6H, s, C₉- and C₁₄-CH₃), 1.70 (3H, s, C₉- or C₁₄-CH₃), 1.71 (3H, s, C₉- or C₁₄-CH₃), 1.74, 1.78 (each 3H, s, C₁₉-CH₃), 1.84 (2H, dd, $J=7, 8$, C₂-H × 2), 2.58 (2H, t, $J=8$, C₃-H × 2), 2.60 (2H, t, $J=7$, C₁-H × 2), 3.24 (2H, d, $J=6$, C₁₂-H × 2), 3.36 (2H, d, $J=6$, C₇-H × 2), 4.51 (2H, d, $J=7$, C₁₇-H × 2), 4.61 (1H, brs, OH), 4.98 (1H, t, $J=6$, C₁₃-H), 5.11 (1H, t, $J=6$, C₈-H), 5.27 (1H, d, $J=11$, C₁₁-H), 5.32 (1H, d, $J=18$, C₁₁-H), 5.47 (1H, t, $J=7$, C₁₈-H), 5.62 (1H, brs, OH), 5.72 (1H, brs, OH), 6.16 (1H, dd, $J=11, 18$, C₁₀-H), 6.30 (1H, s, C₃-H), 6.59 (1H, s, C₂-H), 6.94 (1H, s, C₆-H). ¹³C-NMR (100.4 MHz): Table I.

Kazinol H (6)—Compound **6** was obtained as an oily substance, $[\alpha]_D^{18} +0.53^\circ$ ($c=0.94$, MeOH). FeCl₃ test (brown). Na₂MoO₄ test (negative). UV $\lambda_{\max}^{\text{EtOH}}$ nm (log ϵ): 229 (4.60), 265 (infl. 3.87), 277 (4.05), 286 (4.05), 320 (3.28). IR $\nu_{\max}^{\text{CHCl}_3}$ cm⁻¹: 3550, 3490, 1660, 1595, 1585. High-resolution MS, Calcd for C₃₀H₃₆O₄ (M⁺), m/z 460.2612. Found: m/z 460.2617. MS: m/z 460 (M⁺), 445 (M⁺ - CH₃), 270 (14), 255 (270 - CH₃), 191 (10), 123 (191 - C₃H₈). ¹H-NMR (CDCl₃, 90 MHz) δ : 1.43 (9H, s, C₉-CH₃ × 2 and C₁₆-CH₃), 1.46 (3H, s, C₁₆-CH₃), 1.67, 1.69 (each 3H, s, C₂₁-CH₃), 1.86–2.18 (2H, m, C₃-H × 2), 2.70–2.95 (2H, m, C₄-H × 2), 3.31 (2H, brd, $J=7$, C₁₉-H × 2), 4.99 (1H, t, $J=7$, C₂₀-H), 4.99–5.13 (1H, m, C₂-H), 5.25 (1H, d, $J=10$, C₁₃-H), 5.29 (1H, d, $J=18$, C₁₃-H), 5.36 (1H, brs, OH), 5.63 (1H, d, $J=10$, C₁₆-H), 5.64 (1H, brs, OH), 6.17 (1H, dd, $J=10, 18$, C₁₂-H), 6.36 (1H, s, C₈-H), 6.49 (1H, d, $J=10$, C₁₄-H), 6.90 (1H, s, C₂-H), 6.96 (1H, s, C₅-H). ¹³C-NMR (67.8 MHz): Table I.

Kazinol K (7)—Compound 7 was obtained as an oily substance. FeCl_3 test (brown). Na_2MoO_4 test (negative). UV $\lambda_{\text{max}}^{\text{EtOH}}$ nm (log ϵ): 226 (4.45), 266 (sh, 4.00), 277 (4.18), 285 (4.16), 324 (3.34). IR $\nu_{\text{max}}^{\text{CHCl}_3}$ cm^{-1} : 3610, 3550, 3480, 1625, 1600. High-resolution MS, Calcd for $\text{C}_{30}\text{H}_{38}\text{O}_4$ (M^+), m/z 462.2768. Found: m/z 462.2778. MS: m/z 462 (M^+), 447 ($\text{M}^+ - \text{CH}_3$), 191 (10). $^1\text{H-NMR}$ (CDCl_3 , 400 MHz) δ : 1.40, 1.42 (each 6H, s, C_7 - $\text{CH}_3 \times 2$ and C_9 - $\text{CH}_3 \times 2$), 1.66, 1.71 (each 3H, s, C_{14} - CH_3), 1.82 (2H, quintet, $J=8$, C_2 -H $\times 2$), 2.55 (2H, t, $J=8$, C_3 -H $\times 2$), 2.60 (2H, t, $J=8$, C_1 -H $\times 2$), 3.24 (2H, d, $J=6$, C_{12} -H $\times 2$), 4.97 (1H, t, $J=6$, C_{13} -H), 5.16 (1H, br s, OH, disappeared on addition of D_2O), 5.26 (1H, dd, $J=1, 11$, C_{11} -H), 5.31 (1H, dd, $J=1, 18$, C_{11} -H), 5.40 (1H, br s, OH, disappeared on addition of D_2O), 5.62 (1H, d, $J=10$, C_8 -H), 5.78 (1H, br s, OH, disappeared on addition of D_2O), 6.16 (1H, dd, $J=11, 18$, C_{10} -H), 6.30 (1H, s, C_3 -H), 6.49 (1H, d, $J=10$, C_7 -H), 6.67 (1H, s, C_2 -H), 6.94 (1H, s, C_6 -H). $^{13}\text{C-NMR}$ (100.4 MHz): Table I.

Kazinol K Trimethyl Ether (7a)—A mixture of 7 (20 mg), $(\text{CH}_3)_2\text{SO}_4$ (0.2 ml), and K_2CO_3 (5 g) in $(\text{CH}_3)_2\text{CO}$ (30 ml) was refluxed for 2 h, and treated as usual. The product was purified by preparative TLC (benzene only) to give kazinol K trimethyl ether (7a, 12 mg). Compound 7a was obtained as an oily substance. FeCl_3 test (negative). MS: m/z 504 (M^+). $^{13}\text{C-NMR}$ (CDCl_3 , 67.8 MHz) δ : 17.9 (C_{15}), 25.6 (C_{16}), 26.8 (C_{12}), 27.4 (C_8 , C_9 , C_{10} , and C_{11}), 29.9 (C_2), 31.8 (C_3), 33.1 (C_1), 39.9 (C_7), 55.5 ($\text{OCH}_3 \times 2$), 56.4 (OCH_3), 74.9 (C_9), 96.8 (C_3), 109.3 (C_{11}), 113.8 (C_2), 120.2 (C_5), 120.6 (C_7), 121.6 (C_1), 123.9 (C_{13}), 127.6 (C_6), 128.0 (C_5), 128.2 (C_6), 130.7 (C_8 and C_{14}), 132.5 (C_1), 140.3 (C_4), 146.3 (C_3), 148.5 (C_{10}), 156.4 (C_2), 157.1 (C_4).

Treatment of Kazinol K with DDQ [Formation of Kazinol H (6) from 7]—A mixture of 7 (97 mg) and DDQ (15 mg) in dry benzene (15 ml) was kept at room temperature for 36 h. The reaction product was purified by preparative TLC ($(\text{CH}_3)_2\text{CO} : n\text{-hexane} = 1 : 3$) to give 6 (6.6 mg). Compound 6 thus obtained was identical with authentic 6 on the basis of MS, IR, and $^{13}\text{C-NMR}$ comparisons.

Acknowledgement We are grateful to Emeritus Prof. H. Hara, University of Tokyo, to Emeritus Prof. M. Ikuse, Toho University, and to Prof. A. Sugaya, Faculty of Pharmaceutical Sciences, Josai University, for valuable advice on the identification of the material. We are also grateful to Prof. S. Sakai, Faculty of Pharmaceutical Sciences, Chiba University, for high-resolution mass spectroscopy, to Dr. S. Urano, Tokyo Metropolitan Institute of Gerontology, for $^1\text{H-NMR}$ measurement (200 MHz), and to Prof. I. Yokoe, Faculty of Pharmaceutical Sciences, Josai University, for CD spectral measurement.

References and Notes

- 1) This paper forms Part XXXII of "Constituents of the Cultivated Mulberry Tree." Part XXXI: reference 2.
- 2) J. Ikuta (née Matsumoto), Y. Hano, and T. Nomura, *Heterocycles*, **23**, 2835 (1985).
- 3) a) T. Nomura, *Kagaku No Ryoiki*, **36**, 596 (1982) and references cited therein; b) T. Nomura and T. Fukai, *Heterocycles*, **15**, 1531 (1981); c) T. Nomura, Abstracts of Papers, The 20th Symposium on Phytochemistry, Tokyo, January 1984, p. 1 and references cited therein.
- 4) a) T. Fujimoto, Y. Hano, and T. Nomura, *Planta Med.*, **50**, 218 (1984); b) T. Fujimoto, Y. Hano, T. Nomura, and J. Uzawa, *ibid.*, **50**, 161 (1984); c) T. Fujimoto and T. Nomura, *ibid.*, **1985**, 190.
- 5) J. Matsumoto, T. Fujimoto, C. Takino, M. Saitoh, Y. Hano, T. Fukai, and T. Nomura, *Chem. Pharm. Bull.*, **33**, 3250 (1985).
- 6) a) "Zhong Yao Do Ci Dian," Vol. 2, ed. by Jian Su New Medical, Shanghai, 1977, p. 1502; b) K. Akamatsu, "Wakanyaku," Ishiyaku Shuppan, Tokyo, 1970, p. 511.
- 7) a) K. Takeda and K. Hayashi, "Shokubutsu Shikiso," ed. by K. Hayashi, Yokendo, Tokyo, 1980, p. 179; b) T. Swain and J. L. Goldstein, "Methods in Polyphenol Chemistry," ed. by J. B. Pridham, Pergamon Press, Oxford, 1964.
- 8) a) K. S. Saini and S. Ghosal, *Phytochemistry*, **23**, 2415 (1984); b) F. Gomez, L. Quijiano, G. Garcia, J. S. Calderon, and T. Rios, *ibid.*, **22**, 1305 (1983).
- 9) R. B. Filho, M. F. F. Leite, and O. R. Gottlieb, *Phytochemistry*, **12**, 417 (1973).
- 10) M. E. L. de Almeida, R. B. Filho, M. V. von Bülow, J. J. L. Correa, O. R. Gottlieb, J. G. S. Maia, and M. S. da Silva, *Phytochemistry*, **18**, 1015 (1979).
- 11) a) E. Wenkert and H. E. Gottlieb, *Phytochemistry*, **16**, 1811 (1977); b) K. R. Markham and B. Ternai, *Tetrahedron*, **32**, 2607 (1976).
- 12) a) M. Takasugi, M. Anetai, T. Masamune, A. Shirata, and K. Takahashi, *Chem. Lett.*, **1980**, 339; b) M. Takasugi, Y. Kumagai, S. Nagao, M. Anetai, T. Masamune, A. Shirata, and K. Takahashi, *ibid.*, **1980**, 1459; c) M. Takasugi, N. Niino, M. Anetai, T. Masamune, A. Shirata, and K. Takahashi, *ibid.*, **1984**, 689; d) *Idem*, *ibid.*, **1984**, 693.
- 13) K. S. Dhami and J. B. Stothers, *Can. J. Chem.*, **44**, 2855 (1966).
- 14) a) H. D. Locksley, I. Moore, and F. Sheinmann, *J. Chem. Soc. (C)*, **1966**, 2265; b) F. D. Monache, B. Botta, M. Nicolletti, J. S. de B. Coelho, and F. D. de A. Lyra, *J. Chem. Soc., Perkin Trans. 1*, **1981**, 484; c) H. D. Locksley, A. J. Quillinan, and F. Scheinmann, *J. Chem. Soc. (C)*, **1971**, 3804.

-
- 15) O. R. Gottlieb, *Israel J. Chem.*, **16**, 45 (1977).
 - 16) G. Cardillo, L. Merlini, G. Nasini, and P. Salvadori, *J. Chem. Soc. (C)*, **1971**, 3967.
 - 17) Kazinol I (**13**) was isolated from the cortex of *Broussonetia papyrifera* (L.) VENT., the structure of which is represented by the formula **13** (J. Matsumoto, S. Sekiguchi, M. Someya, and T. Nomura, Abstracts of Papers, the 105th Annual Meeting of the Pharmaceutical Society of Japan, Kanazawa, April 1985, p. 503).