

reflection of [3]1 and (020) of [4]1 correspond closely to molecular lengths. Absence of diffraction in this region from [n]1 is due to the lack of uniform chain length in the polymer. Reflections (110) of [3]1 and (100) of [4]1 reveal a separation of  $\sim 5.2$  Å between parallel molecular axes. The polymer has a strong peak at this  $d$  value, indicating that the staffs of varying lengths pack in a similar manner. Reflections (006) of [3]1 and (080) of [4]1 arise from intrastaff repetition of the bicyclic cages; in the polymer,  $d \approx 3 \frac{1}{3}$  Å. The comparison leaves no doubt that poly-([1.1.1]propellane) indeed is a mixture of [n]staffanes with larger values of  $n$ , as deduced originally<sup>1</sup> from the CP-MAS <sup>13</sup>C NMR spectrum.

**Acknowledgment.** This research was supported by the National Science Foundation (Grant DMR 8807701). K.H. is grateful to the Alexander von Humboldt Foundation for a Feodor Lynen fellowship.

### Rearrangement Approach to Bridgehead Substitution of 1-Methoxybicyclo[2.2.2]oct-5-en-2-ones

Tadao Uyehara,\* Kiyoshi Osanai, Mika Sugimoto, Ichiro Suzuki, and Yoshinori Yamamoto\*

Department of Chemistry, Faculty of Science  
Tohoku University, Sendai 980, Japan

Received May 8, 1989

Bicyclo[2.2.2]oct-5-en-2-ones (1) are potential compounds which are convertible into [5-5] fused-ring systems (2) by the triplet sensitized photochemical process, oxa-di- $\pi$ -methane rearrangement,<sup>1</sup> and also into [6-6] fused-ring systems (4) via vinylation followed by the oxy-Cope rearrangement of the exo-alcohols (3).<sup>2</sup> In connection with our studies on rearrangement strategies from bridged polycyclic compounds to [m-n] fused-ring natural products,<sup>3</sup> we were interested in principal preparation of compounds 1. Straightforward routes to 1 seem to involve Diels-Alder reaction of 1,3-cyclohexadienes and ketene equivalents.<sup>4</sup> However, selective preparation of even a simply substituted 1,3-cyclohexadiene generally requires many steps.<sup>5</sup> Furthermore, the mode of substituents of the bicyclic ketones, thus obtained, depends upon the regioselectivity of the Diels-Alder reaction that we cannot reverse easily. To compensate these weak points, we must develop a new method for introduction of a substituent at the desired position, including the bridgehead carbon, of the bicyclic system. We wish to report herein formal bridgehead substitution of 1-methoxybicyclo[2.2.2]oct-5-en-2-ones (5), derived stereoselectively from anisoles by the Diels-Alder strategy including Birch reduction followed by selective isomerization into 1-methoxy-1,3-cyclohexadienes.<sup>6</sup>

(1) For recent reviews, see: Schaffner, K.; Demuth, M. *Modern Synthetic Method*; Scheffold, R., Ed.; Springer-Verlag: Berlin, 1986; Vol 4, pp 61-88, pp 89-124.

(2) (a) Berson, J. A.; Jones, M., Jr. *J. Am. Chem. Soc.* **1964**, *86*, 5017-5018, 5019-5021. (b) Gregson, P. G.; Mirrington, R. N. *J. Chem. Soc., Chem. Commun.* **1973**, 598-599. (c) Evans, D. A.; Golob, A. M. *J. Am. Chem. Soc.* **1975**, *97*, 4765-4766. (d) Gregson, R. P.; Mirrington, R. N. *Aust. J. Chem.* **1976**, *29*, 2037. (e) Evans, D. A.; Golob, A. M.; Mandel, N. S.; Mandel, G. S. *Ibid.* **1978**, *100*, 8170-8174. (f) Hirota, H.; Kitano, M.; Komatsubara, K.; Takahashi, T. *Chem. Lett.* **1987**, 2079-2080.


(3) Uyehara, T.; Yamada, J.; Ogata, K.; Kato, T. *Bull. Chem. Soc. Jpn.* **1985**, *58*, 211-216. Uyehara, T.; Yamada, J.; Kato, T.; Bohlmann, F. *Bull. Chem. Soc. Jpn.* **1985**, *58*, 861-867. Uyehara, T.; Kabasawa, Y.; Furuta, T.; Kato, T. *Bull. Chem. Soc. Jpn.* **1986**, *59*, 539-543. Uyehara, T.; Kabasawa, Y.; Kato, T. *Bull. Chem. Soc. Jpn.* **1986**, *59*, 2521-2528. Uyehara, T.; Yamada, J.; Furuta, T.; Kato, T.; Yamamoto, Y. *Tetrahedron* **1987**, *43*, 5605-5620. Uyehara, T.; Furuta, T.; Kabasawa, Y.; Yamada, J.; Kato, T.; Yamamoto, Y. *J. Org. Chem.* **1988**, *54*, 3669-3673.

(4) For a review, see: Ranganathan, S.; Ranganathan, D.; Mehrotra, A. K. *Synthesis* **1977**, 289-296.


(5) See, for example: reference 2d.

(6) Evans, D. A.; Scott, W. L.; Truesdale, L. K. *Tetrahedron Lett.* **1972**, 121-124. Birch, A. J.; Dastur, K. P. *Tetrahedron Lett.* **1972**, 4195-4198; *J. Chem. Soc., Perkin Trans. 1* **1973**, 1650-1652. Monti, S. A.; Chen, S.-C.; Yang, Y.-L.; Yuan, S.-S.; Bourgeois, O. P. *J. Org. Chem.* **1978**, *43*, 4062-4069.

Scheme I



Scheme II




Scheme II shows our rearrangement approach to the bridgehead substitution. The first step is a Lewis-acid-mediated pinacol-type transformation of 5 into the thermally more stable conjugated ketones 6. When 5a (X = Me, Y = H) was treated with  $\text{BF}_3 \cdot 2\text{CH}_3\text{OH}$  (1.2 equiv) in dry  $\text{CH}_2\text{Cl}_2$  at room temperature for 1-2 h, the desired ketone 6a was obtained in 90% yield along with 7a (3%) due to ether cleavage.<sup>7</sup> The second step is introduction of a desired bridgehead substituent by treatment with an organometallic reagent (RM) adding in a 1,2-manner. A reaction of 6a with MeLi gave the *endo*-alcohol 8a (R = X = Me, Y = H) in 72% yield. The final step is a pinacol-type rearrangement which proceeds stereoselectively. The alcohol 8a was converted into the bridgehead methyl ketone 1a in 52% yield by treatment with TsOH (0.1 equiv) in boiling benzene for 5 min. This outcome

(7) Similar conversion of a 1-methoxybicyclo[2.2.2]oct-5-en-2-one into a 1-hydroxybicyclo[3.2.1]oct-3-en-2-one, see: Monti, S. A.; Dean, T. R. *J. Org. Chem.* **1982**, *47*, 2679-2681.

**Table I.** Bridgehead Substitution of 1-Methoxybicyclo[2.2.2]oct-5-en-2-ones<sup>a</sup>

|    | substrate |    |                             | product |              | yield <sup>b</sup><br>(%) |
|----|-----------|----|-----------------------------|---------|--------------|---------------------------|
|    | X         | Y  | RM                          |         | R            |                           |
| 5a | Me        | H  | MeLi                        | 1a      | Me           | 65                        |
| 5a | Me        | H  | <i>n</i> -BuLi              | 1b      | <i>n</i> -Bu | 68                        |
| 5a | Me        | H  | <i>i</i> -PrLi <sup>c</sup> | 1c      | <i>i</i> -Pr | 46                        |
| 5a | Me        | H  | <i>t</i> -BuLi <sup>d</sup> | 1d      | <i>t</i> -Bu | 20                        |
| 5a | Me        | H  | PhLi                        | 1e      | Ph           | 66                        |
| 5a | Me        | H  | DIBAH                       | 1f      | H            | 44                        |
| 5b | H         | Me | MeLi                        | 1g      | Me           | 61                        |
| 5c | Me        | Me | DIBAH                       | 1h      | H            | 63                        |
| 5d | H         | H  | MeLi                        | 1i      | Me           | 46                        |

<sup>a</sup>All synthetic intermediates (6, 7, and 8) were used without purification. <sup>b</sup>Overall yields from 5. <sup>c</sup>In situ generated by sonication of a mixture of *i*-PrCl and Li in THF. <sup>d</sup>A mixture of 4-*tert*-butyl-5-methyl-1-methoxybicyclo[3.2.1]oct-2-ones was obtained from 6a in 47% yield.

**Chart I**

suggests that the acid-catalyzed dehydration and the bridge migration are not concerted, and a stable intermediate such as allyl cation 9 may be present. A simple model study of 9 indicates that the C-2 p atomic orbital overlaps preferentially the C-1–C-7  $\sigma$  bond (the two-carbon bridge) rather than the C-1–C-8  $\sigma$  bond (the one-carbon bridge).

The minor products, bridgehead hydroxy ketones 7 and the alcohols derived from 7, were also useful to prepare 1 in a similar manner. In practice, the transformation of 5 to 1 was carried out without chromatographic separation of synthetic intermediates (Table I). The mode of substitution of ketones 1f and 1h is contrary to that of Diels–Alder 1,4-adducts derived from the corresponding 1,3-cyclohexadienes and ketene equivalents. Ketones 1b–e are difficult to prepare selectively by means of simple Diels–Alder strategy.

The ketone 1h has been employed as the starting material for the formal total synthesis of (±)-gymnomitrol (13), a unique tricyclic sesquiterpene.<sup>8,9</sup> Sequential treatment of 1h with LDA, 3-iodopropene (THF–HMPA), LDA, and iodomethane (THF–HMPA) gave 10 stereoselectively in 72% overall yield.<sup>10</sup> The

(8) Isolation: Connolly, J. D.; Harding, A. E.; Thornton, I. M. S. *J. Chem. Soc., Chem. Commun.* 1970, 1320–1321; *J. Chem. Soc., Perkin Trans. 1* 1974, 2487–2493.

(9) Synthesis: (a) Coates, R. M.; Shah, S. K.; Mason, R. W. *J. Am. Chem. Soc.* 1979, 101, 6765–6767; 1982, 104, 2198–2208. (b) Buchi, G.; Chu, P.-S. *Ibid.* 1979, 101, 6767–6768; *Tetrahedron* 1981, 37, 4509–4513. (c) Welch, S. C.; Chayabunjonglerd, S. *J. Am. Chem. Soc.* 1979, 101, 6768–6769. Welch, S. C.; Chayabunjonglerd, S.; Parakasa, R. *J. Org. Chem.* 1980, 45, 4086–4093. (d) Han, Y.-K.; Paquette, L. A. *Ibid.* 1979, 44, 3731–3733. Paquette, L. A.; Han, Y.-K. *J. Am. Chem. Soc.* 1981, 103, 1831–1835. (e) Kodama, M.; Kurihara, T.; Sasaki, J.; Ito, S. *Can. J. Chem.* 1979, 57, 3343–3345.

oxa-di- $\pi$ -methane rearrangement of 10 proceeded efficiently in acetone to form 11 in 94% yield. Reduction of this cyclopropyl ketone by lithium in liquid ammonia gave the substituted diquinane 12 (75% yield) which had already been converted into (±)-gymnomitrol.<sup>9c</sup>

Thus, it is clear that this bridgehead substitution expands the utility of bridged polycyclic compounds for natural product syntheses.

**Acknowledgment.** Financial support by the Kurata Foundation is gratefully acknowledged.

**Supplementary Material Available:** Spectral (IR and <sup>1</sup>H NMR) and analytical data (elemental analysis) for 6a, 6c, 7a, 8a, 1c–h, and 10–12 (3 pages). Ordering information is given on any current masthead page.

(10) The <sup>1</sup>H NMR spectrum of 10 shows a singlet at  $\delta$  0.78 due to the *endo*-3-methyl protons which are shielded by the carbon–carbon double bond. Details are listed in the Supplementary Material.

### The Role of the Product in Asymmetric C–C Bond Formation: Stoichiometric and Catalytic Enantioselective Autoinduction


Albert H. Alberts\* and Hans Wynberg

Department of Chemistry, University of Groningen  
Nijenborgh 16, Groningen, The Netherlands 9747 AG

Received April 4, 1989

Evidence for aggregation of (alkali) metal enolates, alkoxides, and condensation products in the solid state and in solution has been rapidly accumulating in the past decade.<sup>1</sup> The organic moieties are mono- or bidentate ligands in these clusters prior to their liberation to monomeric products in the hydrolytic workup. Although C–C bond formation via condensation of organometallic reagents with carbonyl compounds is by far the most important synthetic procedure in organic chemistry (especially in the area of asymmetric induction with chiral auxiliary ligands), the stereochemical effect of the product acting as a ligand in intermediate complexes has not been systematically investigated.

We studied the effect of enriched (+)-(*R*)-1-phenyl-1-propanol-*d*<sub>1</sub> (+24.40),<sup>2</sup> 1D, on the addition of ethyllithium to benzaldehyde:



To a solution of 2 mmol of the alkyl lithium in benzene (filtered from LiBr, 5 mL, 0.402 M) was added 1 mmol of 1D at room temperature (all operations under dry nitrogen). Ethane was evolved in an exothermic reaction. The homogeneous solution was cooled in ice, and 1 mmol of freshly distilled benzaldehyde in 1 mL of dry benzene was added within 5 min with stirring. After 10 min the reaction mixture was subjected to a hydrolytic workup with dilute HCl and ether. The product, a mixture of 1D and 1H (94%), was purified by molecular distillation (0.1 mm, 100 °C bath temperature, repeated until constant rotation). The use of deuterated 1D allowed us to determine the ee of 1H separately in the mixture. The ratio of 1D vs 1H (found 100:86) was determined by integration of the methine proton in 1H at 5.1

(1) For a recent review: Seebach, D. *Angew. Chem.* 1988, 100, 1685.

(2) Prepared via the quinone-catalyzed addition of diethylzinc to benzaldehyde-*d* in 70% ee (according to Smaardijk, A. A.; Wynberg, H. *J. Org. Chem.* 1987, 52, 135). The optical properties of 1D and 1H are identical within experimental error. The relationship between specific rotation and various mixtures of 1D and racemic 1H is linear. All rotations measured at  $c = 100$ , in toluene, 21 °C, 578 nm.