



J. Serb. Chem. Soc. 79 (6) 637–648 (2014)
JSCS–4614

Structural, spectral and thermal properties of 2-(2-pyridylamino)pyridinium trihydrogen pyromellitate

DEJAN POLETI^{1*#}, JELENA ROGAN^{1#}, LIDIJA RADOVANOVIĆ^{2#}
and MARKO RODIĆ^{3#}

¹Department of General and Inorganic Chemistry, Faculty of Technology and Metallurgy,
University of Belgrade, Karnegijeva 4, 11000 Belgrade, Serbia, ²Innovation Center, Faculty
of Technology and Metallurgy, University of Belgrade, Karnegijeva 4, 11000 Belgrade,
Serbia and ³Faculty of Sciences, University of Novi Sad, Trg Dositeja Obradovića 3,
21000 Novi Sad, Serbia

(Received 7 June, revised 24 October, accepted 25 October 2013)

Abstract: The title compound, (Hdipya)(H₃pyr), where Hdipya is protonated 2,2'-dipyridilamine and H₃pyr is the monoanion of pyromellitic acid (H₄pyr), was obtained from a reaction mixture containing Zn(II) ions, dipya, Na₄pyr (in the mole ratio 2:2:1) and HNO₃. The products (micro- and single-crystalline) were characterized by X-ray structure determination, FT-IR spectroscopy and TG/DSC analysis. The most striking structural feature of (Hdipya)(H₃pyr) are short inter- and extremely short intramolecular hydrogen bonds. These bonds mutually connect cations and anions making thin layers parallel to the crystallographic (223) plane. In addition, quite unusual, linear proton-bound polymeric {H(H₂pyr)}_n were identified in the structure. The results of crystal structure determination are compared with FT-IR and TG/DSC data. The low position of the ν_{as}(COO) vibration at 1660 cm⁻¹ is in accordance with strong hydrogen bonding. This value could be used as a measure of the C=O bond order, which was calculated to be 1.81.

Keywords: pyromellitic acid; 2,2'-dipyridylamine; crystal structure; FT-IR spectra; TG/DSC analysis.

INTRODUCTION

Pyromellitic or 1,2,4,5-benzenetetracarboxylic acid (H₄pyr), one of 12 benzenepolycarboxylic acids, is used in the production of polyesters and polyamides. Anions of H₄pyr, in particular pyr⁴⁻, are well-known as very prospective ligands in coordination chemistry due to the presence of eight O atoms as potential ligating sites that can coordinate up to ten metal centers.¹ The crystal structure of

*Corresponding author. E-mail: dejan@tmf.bg.ac.rs

Serbian Chemical Society member.

doi: 10.2298/JSC130706115P

H₄pyr was published in 1971,² and earliest transition metal (TM) complexes with the tetra-anion of H₄pyr were published just after that.³ The first structurally characterized complexes were described at the beginning of the 1990s.⁴ Later, during structural studies of TM complexes, it was shown that the tetra-anion pyr⁴⁻ usually acts as a centrosymmetric bridging ligand and can build discrete,⁵ chain,⁶ layered,⁷ or framework⁸ structures. A survey of the Cambridge Structural Database, CSD,⁹ showed that the partially deprotonated forms: Hpyr³⁻, H₂pyr²⁻ and H₃pyr⁻ are, as a rule, not coordinated.

Binary and ternary TM complexes with anions of aromatic polycarboxylic acids have been studied for a long time.¹⁰ In line with this subject, attempts were made to prepare mixed ligand Mn(II) and Zn(II) complexes with pyr⁴⁻ and 2,2'-dipyridylamine (dipya) as an *N,N'* ligand. Surprisingly, the title compound containing only potential ligands was obtained instead. Such compounds can be classified into two major groups, as co-crystals (adducts) or organic salts (proton-transfer complexes),¹¹ but between them there are so-called "disordered solid forms",^{12,13} where hydrogen atoms are not undoubtedly located or they are disordered. Although such compounds are much less frequent in comparison with single-component organics or the corresponding complexes, the number of studied examples permanently increases since their possible importance for pharmaceutical chemistry was recognized.^{14,15} Even ferroelectric properties were found in some compounds.¹⁶ Therefore, it seemed worthwhile to study the obtained compound in detail.

In the CSD⁹ there is a very limited number of structurally characterized compounds concerning molecule of H₄pyr or its partially deprotonated ions. There are, however, several similar organic salts: (Hpy)(H₃pyr), where py is pyridine,¹³ (NMe₄)(H₃pyr),¹⁷ a series of H₂pyr²⁻ salts with different pyridinium derivatives,¹⁸ (cytosinium)₂(H₂pyr)·H₂O,¹⁹ and (Hbipy)₂(H₂pyr)(H₄pyr), where bipy is 2,2'-bipyridine, which was characterized structurally²⁰ and by FT-IR and NMR spectroscopy.²¹

On the other hand, dipya is well known as a chelating ligand. A search of the CSD revealed some 170 compounds containing dipya.⁹ Most of them were TM complexes, and there were no examples of uncoordinated dipya in its neutral form. At the same time, there were only eight compounds containing protonated dipya, which can be present as monoprotonated 2-(2-pyridylamino)pyridinium, Hdipya, or diprotonated 2,2'-iminodipyridinium(2+), H₂dipya. Such protonated forms are not coordinated, and in five out of eight cases tetrahalidometalates, [MX₄]²⁻ (M = Co, Cu or Hg; X = Cl or Br), were present as counter ions.⁹

Herein, crystal structure of the title compound, (Hdipya)(H₃pyr), together with its FT-IR spectral and thermal properties are described and correlated with each other.

EXPERIMENTAL

The starting reagents pyromellitic acid (> 95 %), 2,2'-dipyridylamine (98 %), Mn(II) nitrate solution, 45–50 % in dilute nitric acid (all Sigma–Aldrich) and $\text{Zn}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ (*p.a.* Kemika, Zagreb) were used without any purifications.

Preparation of the microcrystalline product. Into 100 cm³ of an aqueous solution containing 0.74 g (2.5 mmol) of $\text{Zn}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$, first 5 cm³ of HNO_3 solution ($c = 1 \text{ mol dm}^{-3}$), then 0.43 g of dipya (2.5 mmol) dissolved in 10 cm³ of an EtOH/H₂O mixture (3:1 volume ratio) was added. The pH value of this mixture was 2. Then, 50 cm³ of an aqueous solution of Na₄pyr (0.32 g; 1.25 mmol) was slowly added under stirring. The final pH value was 4. After standing for 24 h, the obtained precipitate was filtered and rinsed with H₂O, EtOH and Et₂O. Yield: 84.6 %; Anal. Calcd. for C₂₀H₁₅N₃O₈: C, 56.48; H, 3.55; N, 9.88 %. Found: C, 56.27, H, 3.62, N, 9.91 %; FT-IR (KBr, cm⁻¹): 3470 (*bw*), 3117 (*m*), 3097 (*m*), 3061 (*m*), 3024 (*m*), 2993 (*m*), 2920 (*w*), 1715 (*m*), 1660 (*s*), 1604 (*s*), 1562 (*s*), 1451 (*s*), 1311 (*w*), 1292 (*w*), 1270 (*w*), 1250 (*w*), 1170 (*m*), 1151 (*m*), 1005 (*m*), 955 (*m*), 901 (*m*), 771 (*s*), 748 (*s*), 594 (*m*), 559 (*w*), 532 (*m*). The compound was soluble in DMSO but insoluble in H₂O and EtOH.

A similar procedure using Mn(II) instead of Zn(II) ions resulted in a white microcrystalline powder in a comparable yield (78.9 %). As revealed by elemental analysis and FT-IR spectroscopy, this product was identical to the product obtained in the presence of Zn(II) ions.

Preparation of single-crystals. Into 50 cm³ of an aqueous suspension containing 0.25 g (1 mmol) of H₄pyr was added 0.17 g of dipya (1 mmol) dissolved in 5 cm³ of EtOH. Then, 50 cm³ of an aqueous solution containing 0.30 g (1 mmol) of $\text{Zn}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ was added dropwise under continuous stirring. The mixture was gently heated at 50 °C for 3 h, filtered to remove a small quantity of unreacted H₄pyr, and left standing under ambient conditions. The first crystals appeared after 24 h, but single-crystals of appropriate size were collected after 10 days.

Characterization. The FT-IR spectra were recorded on a Bomem MB-100 (Hartmann Braun) spectrophotometer (4000–600 cm⁻¹ region) using KBr pellets. The thermal properties of the compound were examined from room temperature up to 380 °C using an SDT Q600 TGA/DSC instrument (TA Instruments) in a dry nitrogen atmosphere (flow rate: 100 cm³ min⁻¹) at a heating rate of 20 °C min⁻¹. The sample mass was 8.14 mg.

Room temperature (22 °C) single-crystal X-ray diffraction data were collected on an Oxford Gemini S diffractometer equipped with CCD detector, using monochromatized Cu *K*α radiation ($\lambda = 1.5418 \text{ \AA}$). A multi-scan correction for absorption was applied. The structure was solved by direct methods (SIR97)²² and refined on F^2 by full-matrix least-squares using the programs SHELXL97²³ and WinGX.²⁴ All non-hydrogen atoms were refined anisotropically. The positions of the C- and N2-bonded H atoms were calculated geometrically and refined by the riding model with $U_{\text{iso}} = 1.2U(\text{C}, \text{N})$. H atoms from OH groups were found in ΔF maps and refined with the $U_{\text{iso}} = 1.5U(\text{O})$ constraint. The Hdipya cation was disordered over a centre of symmetry with congener atoms having 0.5 site occupancies; the individual pyridyl groups were refined using the FLAT constraint with equal displacement parameters for the corresponding A and B atoms. Attempts to determine the H atom that protonates dipya were unsuccessful, very likely because this atom was spread over four positions (N1A, N1B and their centrosymmetric counterparts, see Fig. 1. Selected crystal data and refinement results are listed in Table I. CCDC 948070 contains supplementary crystallographic data for this paper. This data can be obtained free of charge at <http://www.ccdc.cam.ac.uk/cgi-bin/catreq.cgi>.




Fig. 1. Molecular structure of (Hdipya)(H₃pyr) with atomic numbering scheme. Displacement ellipsoids are drawn at the 50 % probability level. Disordered Hdipya ions are marked by different shades of gray. Symmetry codes: (i) $1 - x, 1 - y, 1 - z$; (ii) $1 - x, 1 - y, -z$; (iii) $1 + x, 1 + y, -z$.

TABLE I. Crystal data and structure refinement for (Hdipya)(H₃pyr)

Empirical formula	C ₂₀ H ₁₅ N ₃ O ₈
<i>M</i> / g mol ⁻¹	425.35
Crystal system	Triclinic
Space group	<i>P</i> $\bar{1}$
<i>a</i> , <i>b</i> , <i>c</i> / Å	3.8268(2), 9.2726(6), 13.3494(8)
α , β , γ / °	74.761(6), 81.845(5), 82.935(6)
<i>V</i> / Å ³	450.59(5)
<i>Z</i>	1
ρ_c / g cm ⁻³	1.568
μ / mm ⁻¹	1.056
<i>F</i> (000)	220
Crystal dimensions, mm	0.09×0.10×0.31
θ_{\max} / °	72.12
Range of <i>h</i> , <i>k</i> , <i>l</i>	$-4 \leq h \leq 3, -11 \leq k \leq 11, -16 \leq l \leq 16$
Reflections collected	7605
Independent reflections	1770 (<i>R</i> _{int} = 0.0361)
Completeness, %	99.6
Data/restraints/parameters	1770/6/169
(Δ/σ) _{max}	0.001
<i>S</i>	1.116
<i>R</i> indices (<i>I</i> > 2σ(<i>I</i>))	<i>R</i> ₁ = 0.0533, <i>wR</i> ₂ = 0.1282
<i>R</i> indices (all data)	<i>R</i> ₁ = 0.0631, <i>wR</i> ₂ = 0.1341
$\Delta\rho_{\max}$, $\Delta\rho_{\min}$ (both in e Å ⁻³)	0.150, -0.205

RESULTS AND DISCUSSION

Synthesis

As described in the Experimental, the title compound was obtained during attempts to synthesize ternary Zn(II)–dipya–pyr complexes or their Mn(II) analogues. This demonstrates the difficulties in the preparation of such complexes, in spite of being known.⁹ A similar experience was already published by Ha,²⁵ when (Hdipya)₂[MnCl₄] was obtained instead of the expected Mn(II)–dipya complex, and it was not reported previously for Zn(II) complexes. Such behavior can be partly explained by an increase on the thermodynamic stability of the protonated dipya in respect to the neutral molecule,²⁶ but since dipya complexes of many TMs are known, the specific role of Zn(II)/Mn(II) ions and the pH of the reaction mixture should not be neglected.

Crystal structure

In the structure of (Hdipya)(H₃pyr), both the Hdipya and H₃pyr ions are positioned around the symmetry centre, and only half of each species belongs to the asymmetric unit (Fig. 1). Besides, Hdipya is disordered over a centre of symmetry with equal allocation of both entities. A comparable disorder was found in some organostannates,²⁶ and in the analogous (Hdipya)(Hpht), where Hpht is the hydrogen phthalate ion.²⁷ One of pyridyl N atoms has to be protonated, but it was not possible to locate the corresponding H atom. Both N atoms from the pyridyl rings could be protonated because N1A...O1 and N1B...O1 distances are 3.018(7) and 3.121(5) Å, respectively, *i.e.*, they could form hydrogen bonds. Therefore, the H atom is very likely spread over four possible positions.

The whole Hdipya is practically perfectly planar with a negligible dihedral angle between the mean-planes of the two pyridyl rings. In three known dipya polymorphs and five crystallographically different dipya molecules, this angle varied between about 4 and 40°. ²⁸ A very similar variety of angles was also found in some TMs–dipya complexes.¹⁰ The planarity of Hdipya might be explained by symmetry constraints or by extended delocalization of the π electrons. However, the influence of intramolecular (intra-H) and intermolecular (inter-H) hydrogen bonds could also be important. According to Haddad *et al.*,²⁹ weak intra-H bonding is also possible between the protonated and unprotonated N atom from adjacent pyridyl rings (Fig. 1, atoms N1A and N1B). As usual, pyridyl rings are less regular than benzene rings.

At first sight, the H₃pyr ion (Fig. 1) is fully protonated, *i.e.*, it should be described as a neutral H₄pyr molecule, not as a monoanion. However, the H2 atom is in special position (site symmetry: $\bar{1}$) and with a 0.5 site occupancy factor. The geometry data for the H₃pyr anion are listed in Table II. While the C–C bond distances are as expected, the angles, in particular C1–C2–C3 and C2–C3–C5, strongly deviate from the values found in H₄pyr·2H₂O (max. 121.7°).³⁰ This is the consequence of the stress caused by the presence of the

intra-H bond O2–H1 \cdots O3 and the formation of seven-membered, S(7), pseudo-rings. In respect to the aromatic ring, the C–COO groups are inclined in the opposite sides, with the corresponding angles of 13.3° for the C2C1O1O2 group and 73.6° for the C3C5O3O4 group. This intra-H bond is extremely short and clearly asymmetrical (Table III, Fig. 1). Very similar geometries were already found in H₂pyr and H₃pyr anions.^{13,18–20} and in some Hpht complexes.^{31,32}

TABLE II. Bond distances and angles in H₃pyr ion (For O–H bond distances, see Table III)

Bond distance, Å			
O1–C1	1.202(4)	O2–C1	1.297(3)
O3–C5	1.225(3)	O4–C5	1.251(3)
C1–C2	1.527(3)	C2–C3	1.406(3)
C2–C4	1.391(3)	C3–C4	1.391(3)
C3–C5	1.513(3)		
Bond angle, °			
O1–C1–O2	121.2(3)	O1–C1–C2	118.9(2)
O2–C1–C2	119.9(2)	C1–C2–C3	129.2(2)
C1–C2–C4	113.0(2)	C3–C2–C4	117.8(2)
C2–C3–C4	118.0(2)	C2–C3–C5	126.8(2)
C4–C3–C5	115.2(2)	C2–C4–C3	124.2(2)
C2–C4–H4	117.9(2)	C3–C4–H4	117.9(2)
O3–C5–O4	122.2(3)	O3–C5–C3	121.9(2)
O4–C5–C3	115.8(2)		

TABLE III. Geometry of hydrogen bonds in (Hdipya)(H₃pyr); Symmetry codes: (iv) $-x, -y + 2, -z + 1$; (v) $x - 1, y, z$

D–H \cdots A	$d(\text{D–H}) / \text{\AA}$	$d(\text{H}\cdots\text{A}) / \text{\AA}$	Angle (DHA) / °	$d(\text{D}\cdots\text{A}) / \text{\AA}$
O2–H1 \cdots O3	1.11(4)	1.34(4)	167(3)	2.435(3)
O4–H2 \cdots O4 ^{iv}	1.24 ^a	1.24 ^a	180 ^a	2.477(2)
N2–H2A \cdots O1 ^v	0.86	1.87	165	2.706(4)

^aAtom H2 is in a special position and its coordinates were fixed during refinement

Another, inter-H bond O4–H2 \cdots O4^{iv} ((iv) $-x, -y + 2, -z + 1$) is symmetrical and also very short, but a little longer than the intra-H bond (Table III). In comparison to typical O \cdots O distances found in related Hpht compounds,²⁷ the intra-H bond is slightly longer, whereas inter-H bond is shorter than expected. This shows how strong the inter-H bond is in this case. Being that the inter-H bond is symmetrical, H₃pyr could also be viewed as a polymeric proton-bound trihydrogen pyromellitate anion, {H(H₂pyr)[–]}_n (Fig. 2). Again, there is great similarity with Hpht compounds, in which proton-bound anions are also found.^{33–35} The main difference is that {H(H₂pyr)[–]}_n is polymeric, while [H(Hpht)₂][–] ions are only dimeric. This is because H₄pyr could be considered as developed from H₂pht by mirroring two COOH groups in the *ortho*-position to another pair of COOH groups. Thus, only H₃pyr[–] and H₂pyr^{2–} can make polymeric proton-bound anions,

while Hpht^- is restricted to dimeric forms.³⁶ To the best of our knowledge, a similar polyanion was found only in $(\text{NMe}_4)(\text{H}_3\text{pyr})$, described by Rodriguez-Cuamatzi *et al.*¹⁷ However, those chains were of a zig-zag type, while here the presented chains are linear (Fig. 2) and so could be regarded as unprecedented.




Fig. 2. Motif of hydrogen bonds (thin dark lines) in the (223) crystallographic plane. Color codes for atoms: H – white, C – light gray, N – dark gray and O – black.

In the structure, there is another hydrogen bond between the amine N2 atom and O1 (Fig. 1, Table III). Since all hydrogen bonds are concentrated in one plane, thin layers approximately parallel to the crystallographic (223) plane are formed (Fig. 2). The $\{\text{H}(\text{H}_2\text{pyr})^-\}_n$ and Hdipya ions are assembled in alternating rows making a non-rectangular grid. The layers are stacked together by short parallel-displaced π - π interactions. For example, all distances between the plane of one benzene ring and benzene C atoms from the nearest layer are 3.490(5) Å. Analogous layers also exist in $(\text{cytosinium})_2(\text{H}_2\text{pyr}) \cdot \text{H}_2\text{O}$,¹⁹ and in $(\text{NMe}_4)(\text{H}_3\text{pyr})$.¹⁷

Spectral and thermal properties

A sequence of weak and sometimes broad IR bands between 3500 and 3200 cm^{-1} is indicative for numerous hydrogen bonds. According to the literature,²¹ even weak and very broad bands centered at 2900, 2430 and 1900 cm^{-1} could also be ascribed to strong hydrogen bonding. In some cases, they could be extremely strong and broad.³⁷ Bands between about 3120 and 2990 cm^{-1} are characteristic for stretching N–H and C–H vibrations in secondary amines and aromatic rings.

It seems that the stretching N–H band was shifted from 3244 cm^{-1} in dipya to 3117 or 3097 cm^{-1} in (Hdipya)(H₃pyr), suggesting stronger hydrogen bonding.

In the “fingerprint” region, there are bands of different origin typical for substituted benzene rings and secondary amines; they agree with the bands in the individual components. Especially characteristic for dipya is a prominent doublet due to N–H out of plane bending at 771 and 748 cm^{-1} . Both vibrations were slightly shifted with respect to pure dipya (768 and 735 cm^{-1}) and very likely overlap with the corresponding C–H vibrations.³⁸

The most interesting part of the FT-IR spectrum is the 1720–1250 cm^{-1} region (Fig. 3), where many strong bands are expected: asymmetrical and symmetrical stretching vibrations of COOH and the corresponding COO groups, C–C and C–N stretching vibrations of aromatic rings and N–H bending in a secondary aromatic amine.³⁹ (Asymmetrical $\nu_{\text{as}}(\text{COOH})$ vibrations, characteristic for organic acids, are often labelled $\nu(\text{C}=\text{O})$ and called carbonyl, due to the presence of double C=O bond). Consistent to the literature,³⁷ $\nu_{\text{as}}(\text{COOH})$ and $\nu_{\text{s}}(\text{COOH})$ frequencies for H₄pyr are at 1710 and 1266 (as doublet at 1278 and 1255) cm^{-1} , respectively, and the present experimental data agree within ± 5 cm^{-1} with these values. These bands are broad and usually the most intense in the spectrum.³⁸ Surprisingly, the corresponding bands in the FT-IR spectrum of (Hdipya)(H₃pyr) are medium to weak, but one additional very prominent band at 1660 cm^{-1} appeared. At first sight, this band could be attributed to the secondary amine N–H bending group frequency, which is located at 1604 cm^{-1} in dipya. However, such high wavenumbers are observed only in the case of coordinated dipya and the band is typically very sharp.¹⁰ Therefore, the band at 1660 cm^{-1} should be ascribed to the $\nu_{\text{as}}(\text{COOH})$ or carbonyl C=O vibrations. Such a low position of $\nu_{\text{as}}(\text{COOH})$ is not expected,³⁸ and, together with no unambiguous identification of $\nu_{\text{s}}(\text{COOH})$ vibrations, has to be correlated to the extremely short and strong hydrogen bonds involving all O atoms from H₃pyr. In (Hbipy)₂(H₂pyr)(H₄pyr),^{20,21} just the opposite behavior was observed, $\nu_{\text{as}}(\text{COOH})$ is shifted to the higher wavenumbers (1726 cm^{-1}), which was explained by the absence of hydrogen bonds.

In the tetrathiafulvalene-*p*-chloranil (TTF–CA) complex, which is a widely studied charge-transfer organic compound,⁴⁰ the position of C=O vibrations was used to determine the degree of charge transfer from TTF to CA. In other words, $\nu_{\text{as}}(\text{COOH})/\nu(\text{C}=\text{O})$ could be related to the C=O bond order. For H₄pyr $\nu_{\text{as}}(\text{COOH})$ was at 1710 cm^{-1} ,³⁹ and this corresponds to a bond order of 2. In alkali metal salts, the COOH groups are ionized and, due to delocalization of the electrons in COO groups, the bond order should be 1.5. The available data for the two alkali metal salts K₄pyr³⁹ and Na₄pyr (this study) give an identical value, 1580 cm^{-1} for $\nu_{\text{as}}(\text{COO})$. From these values, it is easy to calculate the C=O bond order in (Hdipya)(H₃pyr), which was 1.81. By analogy to the “degree of charge transfer”,⁴¹ and should be correlated to the geometry of hydrogen bonding in (Hdipya)(H₃pyr).

However, the observed geometry of H_3pyr ions (Table II) does not enable a simple explanation of the calculated value. For O1–C1 and O2–C1 pair, bond distances can be judged as normal when compared to the H_4pyr and common COOH values.² Nevertheless, O3–C5 bond is longer than expected and O4–C5 bond distance corresponds to the typical values found in ionized COO groups.⁹ This could be a possible cause for the low value of $\nu_{as}(COOH)$, although an extensive coupling of vibrational modes should also be kept in mind.




Fig. 3. FT-IR spectrum (KBr pellets) of (Hdipya)(H_3pyr).

As shown in Fig. 4, the decomposition of (Hdipya)(H_3pyr) began at 237 °C (onset temperature) and ended at about 325 °C with a total mass loss. No melting




Fig. 4. TG/DSC curves of (Hdipya)(H_3pyr) under a dynamic N_2 atmosphere at a heating rate of 20 °C min^{-1} (exotherm up).

was observed and the decomposition started around the boiling point of dipya (222 °C), suggesting that this molecule escaped first. Although the TG curve decreased rapidly, the shape of the DSC curve indicated that the disintegration occurred in at least three, probably four, highly overlapped steps, which could not be resolved. The total molar decomposition enthalpy was relatively high, $\Delta_{\text{dec}}H_{\text{m}}^{\circ} = 461.2 \text{ kJ mol}^{-1}$. In comparison to TM complexes with dipya and some polycarboxylate ligands,¹⁰ the initial decomposition temperature of (Hdipya)(H₃pyr) was lower, very likely owing to the absence of coordinative bonds. In analogous systems (py)₂(H₂tpht), where H₂tpht is terephthalic acid, and (py)₃(H₃tms), where H₃tms is trimesic acid, the total degradation was found to be at 339 and 366 °C, respectively.¹³ There, however, loss of py molecules started just above room temperature, although the boiling point of py is 115 °C.

CONCLUSIONS

A new organic salt (Hdipya)(H₃pyr) was prepared unpredictably during efforts to obtain ternary Zn(II) or Mn(II) complexes with dipya and pyr ligands. Besides the higher stability of Hdipya with respect to dipya,²⁶ the presence of Zn²⁺ or Mn²⁺ ions, as well as a starting mixture pH value of 4 most likely affected the reaction mechanism and the formation of (Hdipya)(H₃pyr). The compound consisted of disordered Hdipya cations and H₃pyr anions connected by hydrogen bonds. There were short inter- and extremely short intra-H bonds. In this way, thin layers parallel to the (223) plane were formed. Due to the very strong and symmetrical inter-H bond, the H₃pyr anions could furthermore be regarded as polymeric {H(H₂pyr)⁻}_n anions. The position and shape of the characteristic bands revealed by FT-IR spectroscopy undoubtedly confirmed the structural findings in (Hdipya)(H₃pyr), and the value of $\nu_{\text{as}}(\text{COO})$ vibration was used for measuring the “degree of proton transfer”, which was calculated to be 1.81. The TG/DSC analysis proved that (Hdipya)(H₃pyr) decomposed in several overlapping steps and a total molar decomposition enthalpy of 461.2 kJ mol⁻¹ was calculated.

SUPPORTING INFORMATION

Crystallographic data can be obtained free of charge at <http://www.ccdc.cam.ac.uk/cgi-bin/catreq.cgi>.

Acknowledgement. This study was financially supported by the Ministry of Education, Science and Technological Development of the Republic of Serbia (Grant No. 45007).

ИЗВОД

СТРУКТУРНА, СПЕКТРАЛНА И ТЕРМИЧКА СВОЈСТВА
2-(2-ПИРИДИЛАМИНО)ПИРИДИНИЈУМ-ТРИХИДРОГЕНПИРОМЕЛИТАТАДЕЈАН ПОЛЕТИ¹, ЈЕЛЕНА РОГАН¹, ЛИДИЈА РАДОВАНОВИЋ² И МАРКО РОДИЋ³¹Капелара за општу и неорганску хемију, Технолошко-металушки факултет, Универзитет у Београду, Карнегијева 4, 11000 Београд, ²Иновациони центар Технолошко-металушког факултета, Универзитет у Београду, Карнегијева 4, 11000 Београд и ³Природно-математички факултет, Универзитет у Новом Саду, Три Досијеја Обрадовића 3, 21000 Нови Сад

Описано једињење, $(\text{Hdpya})(\text{H}_3\text{pyr})$, где је Hdpya протоновани 2,2'-дипиридил-амин, а H_3pyr моноанјон пиромелитне киселине (H_4pyr), добијено је из реакционе смеше која је садржала $\text{Zn}(\text{II})$ јоне, dpya, Na_4pyr (молски однос 2:2:1) и HNO_3 . Производи (микро- и монокристални) окарактерисани су рендгенском структурном анализом, FT-IR спектроскопијом и TG/DSC анализом. Врло необично структурно својство $(\text{Hdpya})(\text{H}_3\text{pyr})$ јесте присуство кратких интер- и екстремно кратких интрамолекуларних водоничних веза, које међусобно повезују катјоне и анјоне градећи танке слојеве паралелне кристалографској равни (223). Такође, у структури су нађени до сада веома ретки, линеарни полимерни анјони $\{\text{H}(\text{H}_2\text{pyr})^-\}_n$ настали повезивањем преко протона. Резултати рендгенске структурне анализе упоређени су са подацима добијеним FT-IR спектроскопијом и TG/DSC анализом. Ниска фреквенција $\nu_{\text{as}}(\text{COO})$ вибрације на 1660 cm^{-1} у сагласности је са постојањем јаким водоничних веза. Ова вредност искоришћена је за одређивање реда $\text{C}=\text{O}$ везе ("степен преноса протона") и израчунато је да он износи 1,81.

(Примљено 7. јуна, ревидирано 24. октобра, прихваћено 25. октобра 2013)

REFERENCES

1. Y.-G. Li, N. Hao, E.-B. Wang, Y. Lu, Ch.-W. Hu, L. Xu, *Eur. J. Inorg. Chem.* (2003) 2567
2. F. Takusagawa, K. Hirotsu, A. Shimada, *Bull. Chem. Soc. Jpn.* **44** (1971) 1274
3. a) T. Ota, H. Imai, *Technol. Rep. Kansai Univ.* **13** (1972) 25; b) T. Ota, H. Imai, *Technol. Rep. Kansai Univ.* **15** (1974) 61
4. a) B. T. Usubaliev, A. N. Shnulin, H. S. Mamedov, *Koord. Khim.* **8** (1982) 1532; b) D. L. Ward, D. C. Luehrs, *Acta Crystallogr., Sect. C: Cryst. Struct. Commun.* **39** (1983) 1370
5. a) D. Poleti, B. Prelesnik, R. Herak, Đ. Stojaković, *Acta Crystallogr., C* **44** (1988) 242; b) A. M. Atria, G. Corsini, L. Gonzalez, M. T. Garland, R. Baggio, *Acta Crystallogr., C* **65** (2009) m250
6. a) D. Poleti, Lj. Karanović, *Acta Crystallogr., C* **45** (1989) 1716; b) A. Majumder, V. Gramlich, G. M. Rosair, S. R. Batten, J. D. Masuda, M. S. El Fallah, J. Ribas, J.-P. Sutter, C. Desplanches, S. Mitra, *Cryst. Growth Des.* **6** (2006) 2355
7. a) J. C. Geng, L. W. Liu, S. L. Xiao, G. H. Cui, *Transition Met. Chem.* **38** (2013) 143; b) X.-Y. Yu, J. Lu, J.-H. Yu, X. Zhang, J.-Q. Xu, T.-G. Wang, *Z. Anorg. Allg. Chem.* **633** (2007) 490
8. a) Sh.-Y. Yang, L.-Sh. Long, R.-B. Huang, L. S. Zheng, S. W. Ng, *Acta Crystallogr., E* **59** (2003); m731; b) Sh.-Y. Yang, L.-Sh. Long, R.-B. Huang, L. S. Zheng, S. W. Ng, *Acta Crystallogr., E* **59** (2003) m921; c) D. Poleti, Lj. Karanović, *J. Serb. Chem. Soc.* **70** (2005) 1441
9. F. R. Allen, *Acta Crystallogr., Sect. B: Struct. Sci.* **58** (2002) 380

10. a) J. Rogan, D. Poleti, *Thermochim. Acta* **413** (2004) 227; b) J. Rogan, D. Poleti, Lj. Karanović, Z. Anorg. Allg. Chem. **632** (2006) 133; c) J. Rogan, D. Poleti, Lj. Karanović, Z. Jagličić, *J. Mol. Struct.* **985** (2011) 371
11. A. D. Bond, *CrystEngComm* **9** (2007) 833
12. S. Mohamed, D. A. Tocher, M. Vickers, P. G. Karamertzanis, S. L. Price, *Cryst. Growth Des.* **9** (2009) 2881
13. S. H. Dale, M. Elsegood, M. Hemmings, A. L. Wilkinson, *CrystEngComm* **6** (2004) 207
14. P. Vishweshwar, J. A. McMahon, J. A. Bis, M. J. Zaworotko, *J. Pharm. Sci.* **95** (2006) 499
15. A. Lemmerer, J. Bernstein, V. Kahlenberg, *J. Chem. Crystallogr.* **41** (2011) 991
16. S. Horiuchi, Y. Tokura, *Nat. Mater.* **7** (2008) 357
17. P. Rodriguez-Cuamatzi, O. I. Arillo-Flores, M. I. Bernal-Uruchurtu, H. Hopfl, *Supramol. Chem.* **19** (2007) 559
18. K. Biradha, M. J. Zaworotko, *Cryst. Eng.* **1** (1998) 67
19. R. Thomas, G. U. Kulkarni, *J. Mol. Struct.* **873** (2008) 160
20. D. Mrvoš-Sermek, Z. Popović, D. Matković-Čalogović, *Acta Crystallogr., C* **52** (1996) 2538
21. P. Novak, S. Sekušak, D. Vikić-Topić, Z. Popović, *J. Chem. Soc., Faraday Trans.* **94** (1998) 1051
22. A. Altomare, M. C. Burla, M. Camalli, G. L. Cascarano, C. Giacovazzo, A. Guagliardi, A. G. G. Moliterni, G. Polidori, R. Spagna, *J. Appl. Crystallogr.* **32** (1999) 115
23. G. M. Sheldrick, *Acta Crystallogr., Sect. A: Found. Crystallogr.* **64** (2008) 112
24. L. J. Farrugia, *J. Appl. Crystallogr.* **32** (1999) 837
25. K. Ha, *Z. Kristallogr. New Cryst. Struct.* **225** (2010) 653
26. S. W. Ng, *Acta Crystallogr., Sect. C: Cryst. Struct. Commun.* (1999) IUC9900098
27. A. Langkilde, D. Madsen, S. Larsen, *Acta Crystallogr., B* **60** (2004) 502
28. a) J. E. Johnson, R. A. Jacobson, *Acta Crystallogr., B* (1973) 1669; b) G. J. Pyrka, A. A. Pinkerton, *Acta Crystallogr., C* **48** (1992) 91; c) H. Schodel, C. Nather, H. Bock, F. Butenschon, *Acta Crystallogr., B* **52** (1996) 842
29. S. F. Haddad, B. F. Ali, R. H. Al-Far, *Polyhedron* **30** (2011) 1061
30. S. H. Dale, M. R. J. Elsegood, *Acta Crystallogr., E* **59** (2003) o1087
31. H. Küppers, *Z. Kristallogr.* **192** (1990) 97
32. X. Wang, Ch. Qin, E. Wang, L. Xu, *J. Mol. Struct.* **737** (2005) 49
33. J. B. Benedict, T. Bullard, W. Kaminsky, B. Kahr, *Acta Crystallogr., C* **60** (2004) m551
34. H. Küppers, *Z. Kristallogr.* **146** (1977) 269
35. a) S. G. Baca, Y. Simonov, M. Gdaniec, N. Gerbelev, I. G. Filippova, G. A. Timco, *Inorg. Chem. Commun.* **6** (2003) 685; b) S. G. Baca, I. G. Filippova, C. Ambrus, M. Gdaniec, Y. A. Simonov, N. Gerbelev, O. A. Gherco, S. Decurtins, *Eur. J. Inorg. Chem.* (2005) 3118
36. D. Poleti, J. Rogan, *Acta Crystallogr., C* **69** (2013) 841
37. D. C. Luehrs, B. C. Cornilsen, C. B. Lover, T. L. Niels, *Inorg. Chim. Acta* **145** (1988) 81
38. E. Castellucci, L. Angeloni, N. Neto, G. Sbrana, *Chem. Phys.* **43** (1979) 365
39. J. Coates, in: *Encyclopedia of Analytical Chemistry*, R. A. Meyers, Ed., Wiley, Chichester, 2000, p. 10815
40. a) A. Girlando, R. Bozio, C. Pecile, J. B. Torrance, *Phys. Rev. B* **26** (1982) 2306; b) P. García, S. Dahaoui, C. Katan, M. Souhassoua, C. Lecomte, *Faraday Discuss.* **135** (2007) 217; c) A. Nagahori, N. Kubota, C. Itoha, *Eur. Phys. J., B* **86** (2013) 109
41. S. L. Johnson, K. A. Rumon, *J. Phys. Chem.* **69** (1965) 74.

Copyright of Journal of the Serbian Chemical Society is the property of National Library of Serbia and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.