

Dehydrogenative Coupling of Monohydrosilanes Mediated by (2-Phosphinoethyl)silylrhodium(I) Complex


Masaaki Okazaki*, Shin Ohshitanai, Hiromi Tobita,* and Hiroshi Ogino
Department of Chemistry, Graduate School of Science, Tohoku University, Sendai 980-8578

(Received May 25, 2001; CL-010491)


Treatment of HSiMe_2Ph with a square-planar rhodium(I) complex $\text{Rh}[(\kappa^2\text{-Si},P)\text{-Me}_2\text{SiCH}_2\text{CH}_2\text{PPh}_2](\text{PMe}_3)_2$ gave rise to selective dehydrogenative coupling of hydrosilanes to afford $(\text{SiMe}_2\text{Ph})_2$ and $\text{RhH}_2[(\kappa^2\text{-Si},P)\text{-Me}_2\text{SiCH}_2\text{CH}_2\text{PPh}_2](\text{PMe}_3)_2$.

The empirical *trans*-influencing ability of silyl ligands has been estimated to exceed both hydrido and alkyl ligands.¹ Considering the strong σ -donor and the high *trans*-influencing ability of silyl ligands, the silyl ligands are expected to exhibit exceptionally strong *trans*-effect. Therefore, silyl ancillary ligands would greatly accelerate the generation of the reactive unsaturated metal center. Nevertheless, to our knowledge, little has been known about the influence of silyl ancillary ligands on the reactivity of transition-metal complexes.² This comes mainly from the facile cleavage of the metal–silicon bonds via reductive elimination, nucleophilic attack at the silyl silicon atom, insertion, and σ -bond metathesis.³ To suppress the elimination of silyl ligands, five-membered (2-phosphinoethyl)silyl chelate ligands have been employed.^{4,5} In this communication, we report the synthesis, structure, and reactivity of $\text{Rh}[(\kappa^2\text{-Si},P)\text{-Me}_2\text{SiCH}_2\text{CH}_2\text{PPh}_2](\text{PMe}_3)_2$.

Reaction of $\text{RhCl}(\text{PMe}_3)_3$ with $\text{HMe}_2\text{SiCH}_2\text{CH}_2\text{PPh}_2$ in toluene at 50 °C gave $\text{RhCl}(\text{H})[(\kappa^2\text{-Si},P)\text{-Me}_2\text{SiCH}_2\text{CH}_2\text{PPh}_2](\text{PMe}_3)_2$ (**1**)⁶ in 65% yield (eq 1). The ^1H , ^{29}Si , and ^{31}P NMR data support the octahedral *mer*-geometry of complex **1**. The iridium analogue $\text{IrCl}(\text{H})[(\kappa^2\text{-Si},P)\text{-Me}_2\text{SiCH}_2\text{CH}_2\text{PPh}_2](\text{PMe}_3)_2$ has been synthesized by the thermal reaction of $[\text{Ir}(\text{CO})(\text{PMe}_3)_4]\text{Cl}$ with $\text{HMe}_2\text{SiCH}_2\text{CH}_2\text{PPh}_2$.^{5c}



Treatment of **1** with 1 equiv of MeLi at room temperature led to the almost quantitative formation of a coordinatively unsaturated silylrhodium(I) complex $\text{Rh}[(\kappa^2\text{-Si},P)\text{-Me}_2\text{SiCH}_2\text{CH}_2\text{PPh}_2](\text{PMe}_3)_2$ (**2**)⁷ (eq 2). Workup of the resulting solution and recrystallization from toluene at –75 °C gave orange crystals of **2** in 35% isolated yield. The molecular structure and selected bond lengths and angles of **2** are shown in Figure 1.⁸ Complex **2** is among few silyl rhodium(I) complexes⁹ and adopts a slightly distorted square planar geometry. The angles P1–Rh–P2 and P3–Rh–Si are 167.65(3) ° and 176.58(2) °, respectively. The Rh–Si bond length



is 2.3937(8) Å that is relatively longer than those of the previously reported silylrhodium(I) complexes $(\text{Ph}_3\text{Si})\text{Rh}(\text{PMe}_3)_3$ [2.317(1) Å]^{9a} and $(\text{PhMe}_2\text{Si})\text{Rh}(\text{PMe}_3)_3$ [2.3804(10) Å].^{9c} The change of the Rh–Si bond lengths is attributable to the degree of the $\text{M}(\text{d}\pi)\text{--SiX}(\sigma^*)$ interaction.¹⁰ When more electron-withdrawing groups are bonded to silicon, this interaction becomes stronger and, as a result, the Rh–Si bond becomes shorter.³ The silyl ligand in **2** bears three electron-donating alkyl groups on the silicon atom which makes the Rh–Si bond of **2** the longest among them. In agreement with the strong *trans*-influence of the silyl ligand, the length of the Rh–P3 bond is longer by 0.05 Å than that of the Rh–P2 bond.






Figure 1. Molecular structure of **2**. Selected bond lengths (Å) and angles (deg): Rh–P1 2.2672(7), Rh–P2 2.2823(7), Rh–P3 2.3339(7), Rh–Si 2.3937(8), P1–Rh–P2 167.65(3), P1–Rh–P3 95.96(3), P1–Rh–Si 81.24(3), P2–Rh–P3, 95.13(3), P2–Rh–Si 87.85(3), P3–Rh–Si 176.58(2).

The reaction 2 may proceed via formation of $\text{RhH}(\text{Me})[(\kappa^2\text{-Si},P)\text{-Me}_2\text{SiCH}_2\text{CH}_2\text{PPh}_2](\text{PMe}_3)_2$, but it was not detected spectroscopically in the course of the reaction. We previously reported the synthesis of $\text{IrH}(\text{Me})[(\kappa^2\text{-Si},P)\text{-Me}_2\text{SiCH}_2\text{CH}_2\text{PPh}_2](\text{PMe}_3)_2$ by the reaction of $\text{IrCl}(\text{H})[(\kappa^2\text{-Si},P)\text{-Me}_2\text{SiCH}_2\text{CH}_2\text{PPh}_2](\text{PMe}_3)_2$ with MeLi.^{5c} Thermolysis of the hydrido(methyl)iridium complex in toluene at 55 °C led to the elimination of methane to give the unidentified product(s) with an Ir–H bond. No formation of the iridium analogue of **2** $\text{Ir}[(\kappa^2\text{-Si},P)\text{-Me}_2\text{SiCH}_2\text{CH}_2\text{PPh}_2](\text{PMe}_3)_2$ was observed. This striking difference of the thermal stability between Rh and Ir is consistent with the general trend: The iridium complexes with high oxidation state are more stable than the corresponding rhodium complexes.¹¹ Complex **2** represents the first example of a coordinatively unsaturated trialkylsilylrhodium(I) complex. Due to the strongly electron-releasing character of the trialkylsilyl moiety, **2** may be highly reactive toward various small molecules.

In fact, the reaction of **2** with 6 equiv of HSiMe_2Ph proceeded spontaneously at room temperature to give the hydrido(silyl)rhodium(III) complex **3**¹² quantitatively. After 20 h at room temperature, the mixture of **3** and HSiMe_2Ph was converted to dihydrido complex **4**¹³ and $(\text{SiMe}_2\text{Ph})_2$ in 91% yield each. The $^{31}\text{P}\{^1\text{H}\}$ NMR data established the *fac*-geometry of **3**. The RhH signal in the ^1H NMR spectrum appears at –10.30 ppm as pseudo double quartets




($J(\text{HP}_{\text{trans}}) = 123.0 \text{ Hz}$, $J(\text{HRh}) = J(\text{HP}_{\text{cis}}) = 16.0 \text{ Hz}$). Other NMR data also support the structure of **3**. The dihydrido complex **4** was independently synthesized by treatment of **1** with LiAlH₄ in THF and characterized based on the results of elemental analysis and spectroscopic data. Scheme 1 shows a plausible mechanism for the formation of **4** and (SiMe₂Ph)₂. The first step is the dissociation of PMe₃ from **3** and then oxidative addition of HSiMe₂Ph takes place to give a silylrhodium(V) intermediate RhH₂(SiMe₂Ph)₂[(κ^2 -Si,P)-Me₂SiCH₂CH₂PPh₂](PMe₃) (**A**). A similar silylrhodium(V) complex to **A** has been reported by Nagashima et al.¹⁴ Complex **A** undergoes reductive elimination of disilane and subsequent ligation of PMe₃ to Rh gives **4**.



Scheme 1. Formation mechanism of **4** and disilane.

Osakada et al. reported the reaction of RhCl(PMe₃)₃ with HSiAr₃.¹⁵ The reaction proceeded at room temperature to give *mer*-[RhCl(H)(SiAr₃)(PMe₃)₃] (**5**) (eq 4). In the reaction, the Si–Si reductive elimination products such as RhCl(H)₂(PMe₃)₃ and (SiAr₃)₂ were not observed.¹⁶ The dramatic difference of reactivity between **3** and **5** is of great interest. The difference is attributable to the exceptionally strong *trans*-effect of the silyl ligand in **3**. Further reaction of **3** and **5** requires the dissociation of the phosphine ligand, which allows it to form bis(silyl)rhodium(V) intermediate. In complex **3**, the silyl moiety of (2-phosphinoethyl)silyl chelate ligand would accelerate the dissociation of the *trans*-PMe₃ ligand, while such an effect is not operative in complex **5**.



It has been known that the dehydrogenative coupling of monohydrosilanes requires the considerably severe conditions and is accompanied by the scrambling of substituents on the silicon atoms.¹⁷ It should be noted that the reaction **3** proceeds at room temperature to give the silicon–silicon coupling product exclusively.

This work was supported by Grants-in-Aid for Scientific Research (Nos. 11740367, 12640533, and 13029012) from the Ministry of Education, Culture, Sports, Science and Technology of Japan. The authors wish to thank NISSAN SCIENCE FOUNDATION and Inoue Foundation for Science for financial supports.

Dedicated to Prof. Hideki Sakurai on the occasion of his 70th birthday.

References and Notes

- a) J. Chatt, C. Eaborn, and S. Ibekwe, *Chem. Commun.*, **1966**, 700. b) R. N. Haszeldine, R. V. Parish, and J. H. Setchfield, *J. Organomet. Chem.*, **57**, 279 (1973).
- H. Tobita, K. Hasegawa, J. J. G. Minglana, L.-S. Luh, M. Okazaki, and H. Ogino, *Organometallics*, **18**, 2058 (1999).
- T. D. Tilley, "The Chemistry of Organic Silicon Compounds," ed. by S. Patai and Z. Rappoport, Wiley, New York (1989), p 1415.
- a) M. J. Auburn, R. D. Holmes-Smith, and S. R. Stobart, *J. Am. Chem. Soc.*, **106**, 1314 (1984). b) M. J. Auburn, S. L. Grundy, S. R. Stobart, and M. J. Zaworotko, *J. Am. Chem. Soc.*, **107**, 266 (1985). c) M. J. Auburn and S. R. Stobart, *Inorg. Chem.*, **24**, 318 (1985).
- a) M. Okazaki, Y. Kawano, H. Tobita, S. Inomata, and H. Ogino, *Chem. Lett.*, **1995**, 1005. b) M. Okazaki, H. Tobita, and H. Ogino, *Chem. Lett.*, **1996**, 477. c) M. Okazaki, H. Tobita, and H. Ogino, *Organometallics*, **15**, 2790 (1996). d) M. Okazaki, H. Tobita, and H. Ogino, *Chem. Lett.*, **1997**, 437. e) M. Okazaki, H. Tobita, and H. Ogino, *J. Chem. Soc., Dalton Trans.*, **1997**, 3531. f) M. Okazaki, H. Tobita, and H. Ogino, *Chem. Lett.*, **1998**, 69. g) M. Okazaki, H. Tobita, Y. Kawano, S. Inomata, and H. Ogino, *J. Organomet. Chem.*, **553**, 1 (1998).
- Data for **1**: ¹H NMR (300 MHz, C₆D₆) δ -9.32 (pseudo dq, ²J(HP_{trans}) = 158.1 Hz, ²J(HP_{cis}) = ¹J(HRh) = 15.3 Hz, 1H, RhH), -7.6 (ddd, ²J(PP_{trans}) = 364.3 Hz, ¹J(PRh) = 107.5 Hz, ²J(PP_{cis}) = 31.5 Hz, PMe₃ (*trans* to PPh₂)), 55.5 (ddd, ²J(PP_{trans}) = 364.3 Hz, ¹J(PRh) = 109.7 Hz, ²J(PP_{cis}) = 26.2 Hz, PPh₂). ²⁹Si{¹H} NMR (59.6 MHz, C₆D₆) δ 40.4 (dddd, ¹J(SiRh) = 26.1 Hz, ²J(SiP_{cis}) = 10.5, 8.2, 6.9 Hz). Anal. Calcd for C₂₂H₃₉SiP₃ClRh: C 46.94, H 6.98%. Found: C 47.21, H 6.99%.
- Data for **2**: ¹H NMR (300 MHz, toluene-*d*₈, -40 °C) δ 0.80 (d, ²J(HP) = 4.7 Hz, 9H, PMe₃), 0.86 (s, 6H, SiMe₃), 1.32 (d, ²J(HP) = 5.7 Hz, 9H, PMe₃), 2.43 (m, 2H, PCH₂), 6.95–7.07, 7.95 (m, 10H, PPh₂). ³¹P{¹H} NMR (121.5 MHz, toluene-*d*₈, -40 °C) δ -21.1 (pseudo dt, ¹J(PRh) = 112.9 Hz, ²J(PP_{cis}) = 35.9 Hz, PMe₃ (*trans* to SiMe₃)), -9.0 (ddd, ²J(PP_{trans}) = 296.0 Hz, ¹J(PRh) = 148.0 Hz, ²J(PP_{cis}) = 36.6 Hz, PMe₃ (*trans* to PPh₂)), 70.5 (ddd, ²J(PP_{trans}) = 296.0 Hz, ¹J(PRh) = 157.2 Hz, ²J(PP_{cis}) = 35.1 Hz, PPh₂). Anal. Calcd for C₂₂H₃₈SiP₃Rh: C 50.19, H 7.28%. Found: C 49.77, H 7.15%.
- Crystallographic data for **2**: C₂₂H₃₈SiP₃Rh, *M_r* = 526.45, *triclinic*, space group P1 (variant No.2), *a* = 11.4349(6) Å, *b* = 12.9113(9) Å, *c* = 9.5714 Å, *α* = 100.379(5)°, *β* = 104.093(2)°, *γ* = 106.562(3)°, *V* = 1265.5(1) Å³, *T* = -123.0 °C, *Z* = 2, *D_c* = 1.38 g cm⁻³, *μ*(Mo Kα) = 9.61 cm⁻¹, *R* = 0.068, *R_w* = 0.189 for 5596 unique reflections.
- a) D. L. Thorn and R. L. Harlow, *Inorg. Chem.*, **29**, 2017 (1990). b) P. Hofmann, C. Meier, W. Hiller, M. Heckel, J. Riede, and M. U. Schmidt, *J. Organomet. Chem.*, **490**, 51 (1995). c) M. Aizenberg, J. Ott, C. L. Eisevier, and D. Milstein, *J. Organomet. Chem.*, **551**, 81 (1998).
- K. Hübler, P. A. Hunt, S. M. Maddock, C. E. F. Rickard, W. R. Roper, D. M. Salter, P. Schwerdtfeger, and L. J. Wright, *Organometallics*, **16**, 5076 (1997).
- "Advanced Inorganic Chemistry," ed. by F. A. Cotton and G. Wilkinson, Wiley, New York (1988), p 777.
- Isolation of **3** was unsuccessful due to its thermal instability. Data for **3**: ¹H NMR (300 MHz, toluene-*d*₈, -20 °C) δ -10.30 (pseudo dq, ¹J(HP_{trans}) = 123.0 Hz, ¹J(HRh) = ²J(HP_{cis}) = 16.0 Hz, 1H, RhH), -7.6 (ddd, ¹J(PRh) = 80.0 Hz, ²J(PP_{cis}) = 32.0 Hz, ²J(PP_{trans}) = 28.6 Hz, PMe₃), -18.2 (ddd, ¹J(PRh) = 103.0 Hz, ²J(PP_{cis}) = 28.6 Hz, ²J(PP_{trans}) = 24.8 Hz, PMe₃), 60.1 (ddd, ¹J(PRh) = 82.0 Hz, ²J(PP_{cis}) = 32.0 Hz, ²J(PP_{trans}) = 24.8 Hz, PPh₂).
Data for **4**: ¹H NMR (300 MHz, toluene-*d*₈) δ -10.57 (pseudo dq, ²J(HP_{trans}) = 135.0 Hz, ²J(HP_{cis}) = ¹J(HRh) = 18.0 Hz, 1H, RhH), -9.27 (pseudo dq, ²J(HP_{trans}) = 135.0 Hz, ²J(HP_{cis}) = ¹J(HRh) = 21.0 Hz, 1H, RhH), 0.61, 0.83 (s, 3H, SiMe), ³¹P{¹H} NMR (121.5 MHz, toluene-*d*₈) δ -20.0 (ddd, ¹J(PRh) = 87.7 Hz, ²J(PP) = 28.6 Hz, ²J(PP) = 19.1 Hz, PMe₃), -11.3 (dt, ¹J(PRh) = 101.1 Hz, ²J(PP) = 26.7 Hz, PMe₃), 68.2 (dt, ¹J(PRh) = 101.1 Hz, ²J(PP) = 21.9 Hz, PPh₂). Anal. Calcd for C₂₂H₄₀SiP₃Rh: C 50.00, H 7.63%. Found: C 49.29, H 7.44%.
- H. Nagashima, K. Tatebe, T. Ishibashi, J. Sakakibara, and K. Itoh, *Organometallics*, **14**, 2868 (1995).
- a) K. Osakada, T. Koizumi, S. Sarai, and T. Yamamoto, *Organometallics*, **16**, 3973 (1997). b) K. Osakada, T. Koizumi, S. Sarai, and T. Yamamoto, *Organometallics*, **17**, 1868 (1998).
- The reaction of RhCl(PMe₃)₃ with HSiMe₂Ph gave the Si–H oxidative addition product quantitatively. Even at 70 °C, no formation of (SiMe₂Ph)₂ was observed.
- a) J. Y. Corey, "Advances in Silicon Chemistry," ed. by G. Larson, JAI Press, Inc., Greenwich, (1991), Vol. 1, p 327. b) M. D. Curtis and P. S. Epstein, *Adv. Organomet. Chem.*, **19**, 213 (1981).