


This article was downloaded by: [University of Stellenbosch]

On: 21 February 2013, At: 04:49

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954

Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

AN EFFICIENT PROCEDURE FOR THE PREPARATION OF MONO, AND DI-BIS-INDOLYL METHANES CATALYZED BY MOLIBDOPHOSPHORIC ACID

Mohammad Ali Zolfigol^a, Peyman Salehi^b & Morteza Shiri^a

^a Chemistry Department, College of Science, Bu-Ali Sina University, Hamadan, I. R. Iran

^b Department of Phytochemistry, Aromatic and Medicinal Plants and Drug Research Institute, Shahid Beheshti University, Tehran, Iran

Version of record first published: 16 Aug 2010.

To cite this article: Mohammad Ali Zolfigol, Peyman Salehi & Morteza Shiri (2004): AN EFFICIENT PROCEDURE FOR THE PREPARATION OF MONO, AND DI-BIS-INDOLYL METHANES CATALYZED BY MOLIBDOPHOSPHORIC ACID, *Phosphorus, Sulfur, and Silicon and the Related Elements*, 179:11, 2273-2277

To link to this article: <http://dx.doi.org/10.1080/10426500490484904>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan,

sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

AN EFFICIENT PROCEDURE FOR THE PREPARATION OF MONO, AND DI-BIS-INDOLYL METHANES CATALYZED BY MOLIBDATOPHOSPHORIC ACID

Mohammad Ali Zolfigol,^a Peyman Salehi,^b and Morteza Shiri^a
Chemistry Department, College of Science, Bu-Ali Sina
University, Hamadan, I. R. Iran,^a and Department of
Phytochemistry, Aromatic and Medicinal Plants and Drug
Research Institute, Shahid Beheshti University, Evin,
Tehran, Iran^b

(Received May 4, 2004; accepted May 4, 2004)

H₃PMo₁₂O₄₀ · xH₂O was found to be an effective catalyst for the preparation of bis-indolyl derivatives from indole and aromatic, aliphatic, heterocyclic aldehydes or ketones in ethanol at room temperature.

Keywords: Aldehyde; bis-indolyl methanes; heteropolyacids; indole; ketone


Acids are widely used as catalysts in industry for producing more than 1×10^8 mt/year of products. The most commonly used are HF, H₂SO₄, HClO₄, and H₃PO₄ (in liquid form or supported on Keiselguhr). Solid acids have many advantages such as simplicity in handling, decreased reactor and plant corrosion problems, and more environmentally safe disposal. Also, wastes and by-products can be minimized or avoided by developing cleaner synthesis routes. On the other hand, reduction in the amount of liquid acid needed and/or simplification in handling procedures is required for risk reduction, economic advantage, and environment protection.¹ Uses of heteropolyacids (HPAs) as catalysts for fine organic synthetic processes have been developed that are very important for industries related to fine chemicals, such as flavors, pharmaceuticals, and food industries.² Heteropolyacids are cheap, reusable, heterogeneous, and easily available catalysts.³ In recent years,

Address correspondence to Mohammad Ali Zolfigol, Chemistry Department, College of Science, Bu-Ali Sina University, Hamadan 65174, P.O. Box 4135, I. R. Iran. E-mail: Zolfi@basu.ac.ir

heteropolyacids and their salts have received considerable attention as powerful reaction mediums for effecting various organic transformations³ such as selective bromination of aromatic compounds, silylation of alcohols, oxidation of aromatic amines to nitro compounds, thioacetalization and transthoacetalization reactions, oxidation of hydroxy groups to their corresponding carbonyl functions, *gem*-acylation,⁴ and Friedel-Crafts⁵ reactions. The use of Lewis acids in the synthesis of bis-indolyl methanes is also well known.⁶ Lewis acids such as AlCl_3 and LiClO_4 , I_2 ⁸ have been used for this purpose. However, the reaction can also be assisted by IR irradiation using a bentonitic clay as catalyst under solvent-free conditions.⁹ The above-mentioned applications of Lewis acids suffer from two limitations. First, more than stoichiometric amounts of the Lewis acid are needed in many cases. Second, these Lewis acids are moisture sensitive and easily decomposed and deactivated in the presence of even a small amount of water. Recently, lanthanide and indium triflates were reported as effective catalysts in the reaction of indoles with aldehydes or ketones.¹⁰ While these salts are active in aqueous solution, longer reaction times (12–36 h) are required to obtain the desired product in good yield. Furthermore, lanthanide triflates are highly expensive. During the course of our work towards the application of solid acids in organic reactions¹¹ we found that $\text{H}_3\text{PMo}_{12}\text{O}_{40} \cdot x\text{H}_2\text{O}$ is an efficient catalyst for the condensation of indole and aldehyde or ketones. Here we wish to report a rapid and convenient procedure for the preparation of bis-indolyl methanes from indole and aldehydes or ketones in the presence of $\text{H}_3\text{PMo}_{12}\text{O}_{40} \cdot x\text{H}_2\text{O}$ in ethanol (Scheme 1 and Table I).

The reaction of benzaldehyde and indole in the presence molybdatophosphoric acid (5% eq) was complete after 20 min in ethanol as solvent at room temperature (entry 1). Both aromatic and heterocyclic aldehydes reacted similarly to give the corresponding bis-indolyl derivatives in high yields and short reaction times (entries 1–7 and 10). Aliphatic aldehyde and ketones reacted smoothly providing the corresponding products in good-to-moderate yields (entries 8, 9, 11). The reaction of 3-nitrobenzaldehyde was complete after 2 h but 4-nitrobenzaldehyde produced the corresponding bis-indolyl methane's after 6 h. Interestingly, when we used terephthalaldehyde, *p*-bis-indolylmethane benzaldehyde was produced in excellent yield (Scheme 1). Also *p*-di(bis-indolylmethane)benzene was obtained in dimethyl sulfoxide (DMSO) as solvent at 120°C after 1 h in high yield.

In conclusion, the cheapness and ready availability of the catalyst, easy and clean workup, and good yields make this method attractive for organic chemists.



SCHEME 1

EXPERIMENTAL

General

Chemicals were purchased from Fluka, Merck, and Aldrich chemical companies. Yields refer to isolated pure products. Products were characterized by comparison of their spectral (IR, ¹H NMR), thin layer chromatography (TLC), and physical data with authentic samples.












General Procedure for Mono Bis-Indolylmethanes

H₃PMo₁₂O₄₀·xH₂O (0.05 mmol) was added to the stirring solution of indole (2 mmol), aldehyde, or ketone (1 mmol) in EtOH (5 ml) at room temperature. The reaction was followed by TLC. On completion H₂O (20 mL) was added to the reaction mixture. The products were precipitated and filtered. Then the residue was purified by column chromatography on silica gel (*n*-hexane/ethyl acetate: 8:2).

Typical Procedure for 1,4-Phenylene-di(bis-indolylmethane)

The mixture of indole (4 mmol), terephthalaldehyde (1 mmol), and H₃PMo₁₂O₄₀·xH₂O (0.10 mmol) in DMSO (10 ml) was heated to 120°C. Reaction was complete after 1 h (monitored by TLC). H₂O (30 ml) was

TABLE I Preparation of Bis-Indolyl Methanes from Indole and Aldehydes or Ketones in the Presence of $\text{H}_3\text{PMo}_{12}\text{O}_{40}$ as a Catalyst in Ethanol at Room Temperature

Entry	Substrate	Time (h)	Isolated yield (%)
1		0.33	93
2		0.18	92
3		1	90
4		0.5	95
5		6	90
6		2	91
7		2	92
8		10	81
9		15	73
10		0.5	80
11		18	90

added to the reaction mixture. The red product precipitated and was isolated by filtration. Then the residue was washed with diethyl ether. Pure p-di(bis-indolylmethane) benzene was obtained.

REFERENCES

- [1] a) M. A. Zolfigol, D. Habibi, B. F. Mirjalili, and A. Bamoniri, *Tetrahedron Lett.*, **44**, 3345 (2003); b) M. A. Zolfigol, I. MohammadPoor-Baltork, D. Habibi, B. F. Mirjalili, and A. Bamoniri, *Tetrahedron Lett.*, **44**, 8165 (2003).
- [2] T. Okuhara, N. Mizuno, and M. Misono, *Adv. Catal.*, **41**, 113 (1996).
- [3] H. Firouzabadi, N. Iranpoor, K. Amani, and F. Nowrouzi, *J. Chem. Soc., Perkin Trans.*, **1**, 2601 (2002) and references cited therein.

- [4] H. Firouzbadi, N. Iranpoor, F. Nowrouzi, and K. Amani, *Tetrahedron Lett.*, **44**, 3951 (2003).
- [5] H. Firouzbadi, N. Iranpoor, and F. Nowrouzi, *Tetrahedron Lett.*, **44**, 5343 (2003).
- [6] a) A. Chatterjee, S. Manna, J. Benerji, C. Pascard, T. Prange, and J. Shoolery, *J. Chem. Soc., Perkin Trans.*, **1**, 553 (1980); b) J. Benerji, A. Chatterjee, S. Manna, C. Pascard, T. Prange, and J. Shoolery, *Heterocycles*, **15**, 325 (1981); c) W. Noland, M. Venkiteswaren, and C. Richards, *J. Org. Chem.*, **26**, 4241 (1961).
- [7] J. S. Yadav, B. V. Subba Reddy, Ch. V. S. R. Murthy, S. M. Kumar, and Ch. Madan, *Synthesis*, 783 (2001).
- [8] B. P. Bandgar and K. A. Shaikh, *Tetrahedron Lett.*, **44**, 1959 (2003).
- [9] G. Penieres-Carrillo, J. G. Garcia-Estrada, J. L. Gutierrez-Ramirez, and C. Alvarez-Toledano, *Green Chem.*, **5**, 337 (2003).
- [10] a) D. P. Chen, L. B. Yu, and P. G. Wang, *Tetrahedron Lett.*, **37**, 4467 (1996); b) S. J. Ji, M. F. Zhou, D. G. Gu, S. Y. Wang, and T. P. Loh, *Synlett.*, 2077 (2003).
- [11] a) M. A. Zolfigol, *Tetrahedron*, **57**, 9509 (2001); b) M. A. Zolfigol, T. Madrakian, E. Ghaemi, A. Afkhami, S. Azizian, and S. Afshar, *Green Chem.*, **4**, 611 (2002); c) P. Salehi, M. M. Khodaei, M. A. Zolfigol, and S. Sirouszadeh, *Bull. Chem. Soc. Jpn.*, **76**, 1863 (2003); d) F. Shirini, M. A. Zolfigol, B. Mallakpour, S. E. Mallakpour, I. Mohammadpoor-Bltork, and A. R. Hajipour, *Tetrahedron Lett.*, **43**, 1555 (2002); e) M. A. Zolfigol, F. Shirin, A. Ghorbani Choghamarani, and I. Mohammadpoor-Baltork, *Green Chem.*, **4**, 562 (2002); f) M. A. Zolfigol, G. Chehardoli, and S. E. Mallakpour, *Synth. Commun.*, **33**, 833 (2003).