

Nitrosyl-bridged Cyclopentadienyl-cobalt and -rhodium Cations

By SARA CLAMP, NEIL G. CONNELLY,* and JOHN D. PAYNE

(Department of Inorganic Chemistry, University of Bristol, Bristol BS8 1TS)


Summary $[M(CO)_2(\eta-C_5R_5)]$ ($M = Co$ or Rh , $R = H$ or Me) and $[NO]^+$ afford $[M(CO)(NO)(\eta-C_5R_5)]^+$ (**3**) and then $[M_2(CO)_2(\mu-NO)(\eta-C_5R_5)_2]^+$ (**4**) which decarbonylates to give $[M_2(\mu-CO)(\mu-NO)(\eta-C_5R_5)_2]^+$ (**5**); complex (**4**) undergoes reversible nitrosyl-bridge cleavage with alkenes, and (**5**) reacts with $[NO]^+$ to give $[\{M(\mu-NO)(\eta-C_5R_5) \}_2]^{2+}$

(**7**) which is reduced to $[\{M(\mu-NO)(\eta-C_5R_5) \}_2]^+$ (**6**; $M = Co$).

THE binuclear cyclopentadienylmetal carbonyls $[M_2(\mu-CO)(CO)_2(\eta-C_5R_5)_2]$ (**1**) and $[\{M(\mu-CO)(\eta-C_5R_5) \}_2]$ (**2**) ($M = Co$ or Rh , $R = H$ or Me) are of topical interest as precursors

(42%), $\tilde{\nu}(\text{NO})$ 1560 cm^{-1} , the paramagnetism of which is revealed by the 15-line e.s.r. spectrum [g_{av} 2.04, $A_{\text{iso}}(^{59}\text{Co})$ 42 G]. Complex (6) which can, in the case of the C_6Me_5 derivative, be made directly from (5) and NO gas in 90% yield, results from the reaction between (5) and $[\text{NO}]^+$. The formation of the complex $[\{\text{M}(\mu\text{-NO})(\eta\text{-C}_5\text{R}_5)_2\}_2][\text{PF}_6]_2$ (7; $\text{M} = \text{Co}$, $\text{R} = \text{Me}$), $\tilde{\nu}(\text{NO})$ 1623 cm^{-1} , is followed by spontaneous reduction to (6; $\text{M} = \text{Co}$, $\text{R} = \text{Me}$), $\tilde{\nu}(\text{NO})$ 1560 cm^{-1} , g_{av} 2.06, $A_{\text{iso}}(^{59}\text{Co})$ 39 G. Further evidence for the existence of the diamagnetic dication (7) is two-fold. First, cyclic voltammetry at a Pt electrode in CH_2Cl_2 shows that (6; $\text{M} = \text{Co}$, $\text{R} = \text{Me}$) undergoes reversible one-electron oxidation at 0.84 V (*vs.* a saturated calomel electrode); the high potential correlates with the ready reduction of (7; $\text{M} = \text{Co}$, $\text{R} = \text{Me}$). Secondly, although $[\text{Rh}(\text{CO})_2(\eta\text{-C}_5\text{H}_5)]$ and $[\text{NO}][\text{PF}_6]$ yield only the red brown complex (4; $\text{M} = \text{Rh}$, $\text{R} = \text{H}$) (40%), $\tilde{\nu}(\text{CO})$ 2074 and 2048, $\tilde{\nu}(\text{NO})$ 1636 cm^{-1} , which is slowly decarbonylated to the dark blue (5; $\text{M} = \text{Rh}$, $\text{R} = \text{H}$), $\tilde{\nu}(\text{CO})$ 1846, $\tilde{\nu}(\text{NO})$ 1524 cm^{-1} , $[\text{Rh}(\text{CO})_2(\eta\text{-C}_5\text{Me}_5)]$ affords the deep blue complex (7; $\text{M} = \text{Rh}$, $\text{R} = \text{Me}$) (55%), $\tilde{\nu}(\text{NO})$ 1610 cm^{-1} .

Although the one-electron oxidation of $[\{\text{Co}(\text{NO})(\eta\text{-C}_5\text{H}_5)_2\}]$ and of $[\text{Co}_2(\mu\text{-CO})(\mu\text{-NO})(\eta\text{-C}_5\text{H}_5)_2]$ has been detected by cyclic voltammetry,⁵ our results show the existence of the three-membered redox series in equation (1) and, by implication, the isoelectronic series in equations (2) and (3).



The synthetic potential of the binuclear complexes described is illustrated by the reactions of (**4**; M = Rh, R = H) with ligands (L = PPh₃, cyclo-octene, *etc.*) to give the complexes [Rh(CO)₂(η-C₅H₅)] and [Rh(NO)L(η-C₅H₅)]⁺ [L = cyclo-octene, 93%, khaki crystals, $\tilde{\nu}(\text{NO})$ 1845 cm⁻¹] (Scheme). Ethylene reacts similarly, but reversibly, in that partial evacuation of a CH₂Cl₂ solution of [Rh(CO)₂(η-C₅H₅)] and [Rh(NO)(C₂H₄)(η-C₅H₅)]⁺ leads to the regeneration of (**4**; M = Rh, R = H). The reversible formation of binuclear species, *via* nitrosyl-ligand bridge-terminal bond-

† Compounds for which yields are quoted have been isolated as solids and characterised by elemental analyses (C and H).

ing, is unprecedented and has clear implications for the reactions at dimetal centres.

We thank the S.R.C. for studentships (to S. C. and J. D. P.) and for funds to purchase electrochemical equip-

ment, and Johnson Matthey for the generous loan of rhodium trichloride.

(Received, 18th May 1981; Com. 600.)

¹ W. A. Herrmann, C. Bauer, J. Plank, W. Kalcher, D. Speth, and M. L. Ziegler, *Angew. Chem., Int. Ed. Engl.*, 1981, **20**, 193 and references therein.

² W. A. Herrmann, J. Plank, M. L. Ziegler, and P. Wülknitz, *Chem. Ber.*, 1981, **114**, 716.

³ L. M. Cirjak, R. E. Ginsburg, and L. F. Dahl, *J. Chem. Soc., Chem. Commun.*, 1979, 470.

⁴ N. M. Boag, M. Green, R. M. Mills, G. N. Pain, F. G. A. Stone, and P. Woodward, *J. Chem. Soc., Chem. Commun.*, 1980, 1171.

⁵ I. Bernal, J. D. Korp, G. M. Reisner, and W. A. Herrmann, *J. Organomet. Chem.*, 1977, **139**, 321.