

This article was downloaded by: [Temple University Libraries]

On: 23 September 2013, At: 05:39

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uopp20>

SYNTHESIS OF 3,6-DI-(tert-BUTYL)FLUORENE BY NICKEL(0) CATALYZED COUPLING OF ARYL HALIDES

Gitendra C. Paul^a & Joseph J. Gajewski^a

^a Department of Chemistry, Indiana University, Bloomington, IN, 47405

Published online: 09 Feb 2009.

To cite this article: Gitendra C. Paul & Joseph J. Gajewski (1998) SYNTHESIS OF 3,6-DI-(tert-BUTYL)FLUORENE BY NICKEL(0) CATALYZED COUPLING OF ARYL HALIDES, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 30:2, 222-225, DOI: [10.1080/00304949809355284](https://doi.org/10.1080/00304949809355284)

To link to this article: <http://dx.doi.org/10.1080/00304949809355284>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

- Abstr.*, **80**, 59772 (1974)]; J. Otto and K. Paluch, *Rocz. Chem.*, **47**, 967 (1973) and **46**, 1551 (1972) [*Chem. Abstr.*, **79**, 136679 (1973) and **78**, 57909 (1973)].
5. a) M. Hirano, S. Yakabe, J. H. Clark and T. Morimoto, *J. Chem. Soc., Perkin Trans. 1*, 2693 (1996); b) M. Hirano, K. Ukawa, S. Yakabe and T. Morimoto, *Org. Prep. Proced. Int.*, **29**, 480 (1997); c) M. Hirano, K. Komiya, S. Yakabe, J. H. Clark and T. Morimoto, *ibid.*, **28**, 705 (1996); d) M. Hirano, K. Komiya and T. Morimoto, *ibid.*, **27**, 703 (1995); e) M. Hirano, S. Yakabe, J. H. Clark, H. Kudo and T. Morimoto, *Synth. Commun.*, **26**, 1875 (1996); f) M. Hirano, Y. Ueno and T. Morimoto, *ibid.*, **25**, 3125 (1995); (g) *idem.*, *ibid.*, **25**, 3765 (1995); h) M. Hirano, H. Kudo and T. Morimoto, *Bull. Chem. Soc. Jpn*, **67**, 1492 (1994).
 6. For electrophilic aromatic halogenations, see; Ref. 1(c); J. March, *Advanced Organic Chemistry*, 4th ed., John Wiley & Sons, **1992**, p. 531; M. B. Smith, *Organic Synthesis*, McGraw-Hill, New York, **1994**, p.185.
 7. Recently developed aromatic brominations: K. Smith and D. Bahzad, *Chem. Commun.*, 467 (1996); V. Paul, A. Sudalai and K. V. Srinivasan, *Synth. Commun.*, **25**, 2401 (1995); M. C. Carreno, J. L. G. Ruano, G. Sanz, M. A. Toledo and A. Urbano, *J. Org. Chem.*, **60**, 5328 (1995); H. Eguchi, H. Kawaguchi, S. Yoshinaga, A. Nishida, T. Nishiguchi and S. Fujisaki, *Bull. Chem. Soc. Jpn* **67**, 1918 (1994); Y. Goldberg and H. Alper, *J. Org. Chem.*, **58**, 3072 (1993); B. C. Ranu, D. C. Sarkar and R. Chakraborty, *Synth. Commun.*, **22**, 1095 (1992); J. G. Lee, H. T. Cha, U. C. Yoon, Y. S. Suh, K. C. Kim and I. S. Park, *Bull. Korean Chem. Soc.*, **12**, 4 (1991); S. Kajigaeshi, M. Moriwaki, T. Tanaka, S. Fujisaki, T. Kakinami and T. Okamoto, *J. Chem. Soc., Perkin Trans 1*, 897 (1990).
 8. M. Siskin, G. Brons, A. R. Katritzky and R. Murugan, *Energy Fuels*, **4**, 482 (1990).
 9. R. A. Smith, *J. Am. Chem. Soc.*, **55**, 3718 (1933).

SYNTHESIS OF 3,6-DI-(*tert*-BUTYL)FLUORENE BY NICKEL(0) CATALYZED COUPLING OF ARYL HALIDES


Submitted by
(09/09/97)

Gitendra C. Paul* and Joseph J. Gajewski*

Department of Chemistry
Indiana University, Bloomington, IN 47405

In the course of our investigations of fluorenyl substituted *m*-xylylenes, substantial amounts of 3,6-di-(*tert*-butyl)fluorenone were required. An obvious route is *via* the corresponding 3,6-di-(*tert*-butyl)fluorene. Unfortunately electrophilic substitution of fluorene occurs at 2- and 7-positions rather than the 3- and 6-positions. This led Kajigaeshi and coworkers¹ to synthesize the title compound from


p,p'-di-(*tert*-butyl)diphenylmethane according to **Scheme 1**.



Scheme 1

While following this route, we encountered certain difficulties. The Ullmann reaction which requires high temperature, was not reproducible in our hands. There were also some inconsistencies in the Experimental Section, in particular for the iodination step. Here we provide full details as well as improvements in the synthesis of the fluorenone **4** including major modification at the coupling step.

The intramolecular coupling of the diiodide **2** was achieved using nickel(0) complex under mild conditions. The method was a combination of one originally developed by Semmelhack,² later modified by Kende³ [*in situ* generation of the nickel(0) complex] and Kumada⁴ [use of catalytic amount of the nickel(0) complex]. Thus *p,p'*-diphenylmethane⁵ was iodinated with a mixture of periodic acid and iodine according to a literature procedure⁶ to afford **2** in 83% yield. The diiodide was then coupled in the presence of *tetrakis*(triphenylphosphine)nickel(0) catalyst which was generated *in situ* from the air stable *bis*(triphenylphosphine)nickel(II) chloride (10 mol%), triphenylphosphine and zinc powder [to reduce nickel(II) to nickel(0)] in DMF. Fluorene **3** was obtained in 85% yield (**Scheme 2**).



Scheme 2

This coupling could also be achieved from *in situ* preparation of nickel(0) complex (*stoichiometric amount*) directly from nickel(II) chloride hexahydrate, triphenylphosphine and activated zinc.⁷ Thus these results show that in addition to the formation of 6-, 7- and other larger rings, a 5-membered ring can also be formed using nickel(0) catalyst.⁸ The 3,6-di-(*tert*-butyl)fluorene thus obtained was oxidized with a 1:1 mixture of PDC and *tert*-butylhydroperoxide⁹ at room temperature to give 3,6-di-(*tert*-butyl)fluorenone in good yield (80%).

EXPERIMENTAL SECTION

1H and ^{13}C NMR spectra were recorded on a Varian VXR-400 MHz instrument in $CDCl_3$. All chemical shifts are reported as δ values in parts per million downfield from TMS.

Synthesis of 2,2'-Diiodo-4,4'-di-(*tert*-butyl)diphenylmethane (2).— To a solution of *p,p'*-diphenylmethane (8.91 g, 31.78 mmol), periodic acid dihydrate (2.90 g, 12.73 mmol) and iodine (6.46 g, 25.46 mmol) in glacial acetic acid (34 mL) was added water (5.5 mL) and H₂SO₄ (95%, 1.2 mL). The mixture was stirred and heated to 70–78° for 24 h. A brown oily liquid deposited at the bottom of the flask. The mixture was poured into ice-water and extracted with ether. The ether layer was washed with NaHSO₃ solution, Na₂CO₃, then water and brine. After drying (MgSO₄) rotary evaporation gave a light yellow residue (16.1 g) which was then chromatographed through an alumina column (hexane:benzene = 2:1). This gave a white semisolid (14.3 g, >83%) which was used without further purification. ¹H NMR: δ 7.86 (d, *J* = 2.0 Hz, 2H), 7.27 (dd, *J* = 2.0, 8.0 Hz, 2H), 6.85 (d, *J* = 8.0 Hz, 2H), 4.06 (s, 2H), 1.29 (s, 18H).

3,6-Di-(*tert*-butyl)fluorene (3).— **Method A:**⁴ *bis*(Triphenylphosphine)nickel dichloride (0.141 g, 0.216 mmol), Zn powder (0.395 g, 6.05 mmol) and triphenylphosphine (0.453 g, 1.73 mmol) were placed in a 2-neck flask (100 mL) fitted with a stopcock adapter which was connected to a vacuum line. The flask was carefully evacuated and then filled with nitrogen. Degassed DMF (20 mL) was added and the mixture was heated using an oil bath (bath temp. 50–75°). After 1 h, a solution of diiodide **2** (1.15 g, 2.16 mmol) in 20 mL of DMF was added *via* a cannula and was allowed to stir at 70–75° for 24 h. It was then stirred at room temp. for an additional 24 h by which time the original green color had changed to reddish brown. The mixture was then treated with 1 N HCl, and extracted with diethyl ether. The ethereal layer was washed successively with water and brine and dried (MgSO₄). Rotary evaporation gave a solid (2.9 g) whose proton NMR revealed it to be the desired product (100% conversion) along with triphenylphosphine oxide and triphenylphosphine. Most of the phosphine oxide was removed by precipitation (hexane) and filtration. The filtrate was chromatographed through silica gel (hexane : benzene = 10 : 1) to afford the desired fluorene **3** (0.510 g, 85%).

Method B:⁷ Nickel(II) chloride hexahydrate (NiCl₂·6H₂O, 0.283 g, 1.19 mmol), triphenylphosphine (1.25 g, 4.76 mmol) and activated zinc (~0.1 g, 1.5 mmol) (zinc was activated by treating with 10% HCl followed by washing successively with water and acetone and then drying) were placed in a 2-neck flask containing stirrer and a stopcock adapter. The flask was evacuated and filled with nitrogen. After addition of DMF (8 mL), the mixture was heated in an oil bath as above. After 1 h, a solution of diiodide **2** (0.68 g, 1.18 mmol) in DMF (3 mL) was added. After stirring for 24 h at 70–75°, the reaction mixture was treated as described above to afford a white solid whose proton NMR revealed it to be fluorene **3** (>90% conversion) along with triphenylphosphine oxide and triphenylphosphine. This mixture was taken up in THF and treated with 30% hydrogen peroxide to convert the phosphine to the phosphine oxide. Addition of saturated sodium sulfite solution (4 mL) followed by extraction (ether- hexane mixture), drying, filtration and rotary evaporation gave a residue which was triturated with hexane and filtered to remove most of the phosphine oxide. The filtrate was chromatographed as above to afford fluorene **3** (0.250 g, 76%). ¹H NMR: δ 7.82 (d, *J* = 1.6 Hz, 4,5-H, 2H), 7.45 (d, *J* = 8.0 Hz, 1,8-H, 2H), 7.34 (dd, *J* = 1.6, 8.0 Hz, 2,7-H, 2H), 3.81 (s, 2H), 1.42 (s, 18H).

3,6-Di-(*tert*-butyl)fluorenone (4).- The procedure was the same as that used for fluorene.⁹ Thus from 3,6-di-(*tert*-butyl)fluorene **3** (2.36 g, 8.50 mmol), PDC (12.79 g, 34 mmol), celite (6 g), Bu'OOH (70%, 4.38 g, 34 mmol) and benzene (70 mL) there was obtained 2.3 g (>95% conversion) crude yellow product. Recrystallization from a mixture of hexane and benzene gave pure **4** (yellow crystals, ~2 g, 80%). The compound sublimes at atmospheric pressure below its mp., mp. (sealed tube) 202-203° (lit¹ mp. 208-209°). ¹H NMR: δ 7.57 (d, *J* = 8.0 Hz, 2H), 7.55 (d, *J* = 1.6 Hz, 2H), 7.30 (dd, *J* = 1.6, 8.0 Hz, 2H), 1.39 (s, 18H). ¹³C NMR: δ 193.54, 158.74, 144.72, 132.46, 125.84, 124.01, 117.31, 35.62, 31.21. MS (*m/z*): 292 (*M*⁺, 67), 277 (100). HRMS for C₂₁H₂₄O: found 292.18356, calcd. 292.18283. Compound **4** was also converted to precursors of *m*-xylylene diradicals.¹⁰

Acknowledgement.- We thank the Department of Energy for support of this work.

REFERENCES

1. S. Kajigaeshi, T. Kadowaki, A. Nishida and S. Fujisaki, *Bull. Chem. Soc. Jpn*, **59**, 97 (1986).
2. M. F. Semmelhack, P. Helquist, L. D. Jones, L. Keller, L. Mendelson, L. S. Ryono, J. G. Smith and R. D. Stauffer, *J. Am. Chem. Soc.*, **103**, 6460 (1981) and references cited therein.
3. A. S. Kende, L. S. Liebeskind and D. M. Braitsch, *Tetrahedron Lett.*, 3375 (1975).
4. M. Zembayashi, K. Tamao, J.-I. Yoshida and M. Kumada, *ibid.*, 4089 (1977).
5. M. Tashiro, T. Yamato and G. Fukata, *J. Org. Chem.*, **43**, 1413 (1978).
6. H. Suzuki, K. Nakamura and R. Goto, *Bull. Chem. Soc. Jpn*, **39**, 128 (1968).
7. M. Tiecco, L. Testaferri, M. Tingoli, D. Chianelli and M. Montanucci, *Synthesis*, 736 (1984).
8. To our knowledge no 5-membered ring has been synthesized using this procedure.²
9. N. Chidambaram and S. Chandrasekaran, *J. Org. Chem.*, **52**, 5048 (1987).
10. J. J. Gajewski and G. C. Paul, *Tetrahedron Lett.*, 00000 (1997)
