Covalent Alternatives for the Elusive Iminophosphonium Cation. Crystal Structures of PCI(NPrⁱ₂)₂NPh·AICI₃ and P(NPrⁱ₂)₂[NPh(SO₂CF₃)]O[§]

Neil Burford, *,†,^a Rupert E. v. H. Spence^a and John F. Richardson *,‡,^b

^a Department of Chemistry, Dalhousie University, Halifax, Nova Scotia B3H 4J3, Canada ^b Department of Chemistry, University of Louisville, Louisville, KY 40292, USA

Reactions between diaminoiminophosphinic chlorides and the chloride-abstracting agents AlCl₃ and Ag(SO₃CF₃) have been comprehensively examined as potential routes to the three-co-ordinate iminophosphonium cation. However, bis(dialkylamino)(phenylimino)phosphinic chlorides react to give the covalent phosphoryl systems PCl(NR₂)₂NPh-AlCl₃ and P(NR₂)₂[NPh(SO₂CF₃)]O respectively, in quantitative yield. The AlCl₃ complexes are also prepared quantitatively from the reaction of P(NPri₂)₂⁺AlCl₄⁻ with PhN₃ (Staudinger reaction). All compounds have been comprehensively characterised and the structures of the isopropyl derivatives confirmed by X-ray crystallography. Crystal data: PCl(NPri₂)₂NPh-AlCl₃, space group *P*2₁, *a* = 8.719(2), *b* = 15.553(2), *c* = 9.509(2) Å, $\beta = 102.73(2)^{\circ}$, *Z* = 2, *R* = 0.029; P(NPri₂)₂[NPh(SO₂CF₃)]O, space group *P*2₁/*n*, *a* = 10.594(3), *b* = 13.222(2), *c* = 17.084(3) Å, $\beta = 97.71(2)^{\circ}$, *Z* = 4, *R* = 0.048. The results indicate that the three-coordinate diaminoiminophosphonium cation is thermodynamically unstable with respect to the observed alternative covalent structures, and that steric shielding is responsible for the stability of the analogous diaminomethylenephosphonium cation.

We have begun a programme into the synthesis of low coordinate phosphonium centres which are analogues of the alkenic, alkynic and allenic environments for carbon, and therefore represent important keys in the development of pnicogen chemistry. The first structurally characterised example of such a centre is the three-co-ordinate methylenephosphonium centre **2b** $[E = C(SiMe_3)_2, Y = NPr_2^i]$, recently reported by Bertrand and co-workers.¹ In 1982 Sanchez and co-workers^{2a} recognised the chlorophosphoryl unit 1 (E = NR or S) as a potential precursor to the alkenic phosphonium centre and reported a solution identification of imino-derivatives (E = NR) of 2a. However, more recent studies by Chernega et al.³ and this laboratory^{4,5} reveal alternative structural arrangements for the cationic imino-³ and chalcogeno-^{4,5} phosphoryl unit. Therefore, we have now performed a comprehensive investigation into the PCl(NR₂)₂NPh-AlCl₃ and -Ag(SO₃CF₃) systems ($\mathbf{R} = \mathbf{E} \mathbf{t}$ or \mathbf{Pr}^i). The spectroscopic and structural data conclusively demonstrate the covalent nature of these systems, and provide a better understanding of the stability of the rare three-co-ordinate phosphonium centre 2.



Experimental

General Procedures.-Phenyl azide (20%, hexane solution,

† For correspondence regarding chemistry.

Pfaltz and Bauer) and Ag(SO₃CF₃) (Aldrich) were obtained commercially and used without purification. Anhydrous AlCl₃ (Aldrich) was sublimed under vacuum before use. Dichloromethane and hexane were dried over P_2O_5 and CaH_2 and stored in evacuated bulbs, CD2Cl2 and CDCl3 were dried over P_2O_5 and CD_2Cl_2 was stored over $[P(NEt_2)_2Se]_2[AlCl_4]_2$.⁴ The compounds $PCl(NEt_2)_2^6$ and $PCl(NPr_2)_2^7$ were prepared by literature methods and characterised by NMR spectroscopy, $PCl(NEt_2)_2NPh$ by reaction of $PCl(NEt_2)_2$ with PhN_3 in hexane and characterised similarly. Solids were manipulated in a Vacuum/Atmosphere nitrogen-filled glove-box. Glass equipment was flame dried before use. Reactions were performed in an evacuated (10-3 Torr, ca. 0.133 Pa) dual-compartment vessel, a modification of that described earlier.⁸ Melting points were recorded on a Fisher-Johns apparatus and are uncorrected. Elemental analyses were performed by Beller Laboratories, Göttingen. Infrared spectra were recorded as Nujol mulls on CsI plates using a Perkin-Elmer 283B spectrophotometer, NMR spectra on a Nicolet NT-360 spectrometer in 5 and 10 mm flame-sealed Pyrex tubes. Chemical shifts are reported in ppm relative to external standards, 85% H₃PO₄ for ³¹P and $[Al(H_2O)_6]^{3+}$ for ²⁷Al, and relative to the solvent signal for ¹H and ¹³C. *i*, *o*, *m* and *p* refer to the *ipso*, *ortho*, *meta* and *para* carbon atoms respectively. Mass spectra were obtained using a CEC model 21-104 mass spectrometer.

Preparation of PCl(NPrⁱ₂)₂NPh.—A solution of PhN₃ (1.73 g, 14.6 mmol) in CH₂Cl₂ (10 cm³) was added to a solution of PCl(NPrⁱ₂)₂ (3.52 g, 13.2 mmol) in CH₂Cl₂ (11 cm³) and the reaction mixture was heated to 95 °C for 48 h. The volatiles were removed to give a solid, which was purified by double vacuum sublimation at 115 °C and recrystallisation from warm hexane to give a large single colourless crystal, characterised as bis(diisopropylamino)(phenylimino)phosphinic chloride, yield 0.48 g, 1.3 mmol (10%), m.p. 86 °C. Mass spectrum [*m*/*z* (relative intensity)]: 357 (26), 322 (22), 314 (68), 166 (74) and 100 (100). NMR: ³¹P, δ -8; ¹³C, δ 147.4 (*i*), 128.5 (*m*), 122.4 (*J*_{PC} = 20.1 Hz) (*o*), 118 (*p*), 47.5, 22.5 and 22.2. IR: 1600s, 1505s, 1200s, 1175s, 1160(sh), 1125s, 1095m, 1070w, 1015m, 990s,

1615

[‡] For correspondence regarding crystallography.

[§] Bis(diisopropylamino)(phenylimino)phosphinic chloride-aluminium trichloride (1/1) and bis(diisopropylamino)[phenyl(trifluoromethyl-sulphonyl)amino]phosphine oxide.

Supplementary data available: see Instructions for Authors, J. Chem. Soc., Dalton Trans., 1991, Issue 1, pp. xviii–xxii.

970(sh), 885m, 850w, 750s, 690s, 665m, 615m, 560(sh), 540s, 505(sh), 450w, 410w and $385w \text{ cm}^{-1}$.

A second crop of crystals (2.20 g; total yield 2.68 g, 7.46 mmol, 57%) was obtained from the hexane solution. The compound is air sensitive, and hydrolyses slowly over several months in the solid state.

Preparation and Isolation of $PCl(NR_2)_2NPh \cdot AlCl_3$ (R = Et or Pr^{i}) Complexes.—A solution of $P(NR_2)_2^+AlCl_4^-$ was prepared ⁹ in situ by the reaction of PCl(NR₂)₂ ($\bar{R} = Et$, 1.31 g, 6.22 mmol; Pr^{i} , 2.42 g, 9.08 mmol) with AlCl₃ (R = Et, 0.83 g, 6.22 mmol; Pr^i , 1.21 g, 9.08 mmol) in CH_2Cl_2 (R = Et, 38 g; Prⁱ, 46 g). Upon solvation of AlCl₃ the solution was cooled to -78 °C and a solution of phenyl azide (R = Et, 0.74 g, 6.22 mmol; Pr^{i} , 1.08 g, 9.08 mmol) in $CH_{2}Cl_{2}$ (R = Et, 13 g; Pr^{i} , 23 g) was added in small aliquots. Evolution of N₂ began as the solution warmed to room temperature and the reaction was complete after stirring for several hours at room temperature. The coloured solutions (R = Et, yellow; Pr^i , orange) were degassed, two thirds of the CH₂Cl₂ was removed and hexane (5 cm³) was added. Slow removal of solvent yielded a crystalline solid, which was recrystallised by slow cooling of the warm (55 °C) solvent mixture, to give large transparent crystals. The crystals were characterised as bis(dialkylamino)(phenylimino)phosphinic chloride-trichloroaluminium.

R = Et: yield 1.11 g, 2.55 mmol (41%), m.p. 163–165 °C. Mass spectrum [*m*/*z* (relative intensity)]: 301 (27), 266 (12), 229 (19), 194 (13) and 138 (100). NMR: ³¹P, δ 48.5; ¹³C, δ 141.2 (*i*), 129.7 (*m*), 129.3 (*o*), 127.1 (*p*), 40.5 and 12.1; ²⁷Al, δ 100 (linewidth at half height $\Delta v_{\frac{1}{2}}$ = 132 Hz). IR: 1595m, 1490(sh), 1340(sh), 1285w, 1200s, 1150s, 1115w, 1095w, 1075w, 1040s, 1020s, 975w, 950w, 920w, 910w, 850s, 810m, 795m, 765m, 710w, 695m, 680m, 625m, 565s, 525(sh), 500s, 490s, 460(sh), 435w and 400s cm⁻¹ (Found: C, 38.70; H, 5.90; N, 9.55; P, 7.00. Calc. for C₁₄H₂₅AlCl₄N₃P: C, 38.65; H, 5.80; N, 9.65; P, 7.10%).

The isolated yields are not optimised. Phosphorus-31 NMR studies of these reaction mixtures and those of the free base iminophosphinic chlorides with AlCl₃ [PCl(NR₂)₂NPh: R = Et, 0.11 g, 0.37 mmol; Prⁱ, 0.11 g, 0.30 mmol. CH₂Cl₂: R = Et, 4 g; Prⁱ, 3.1 g. AlCl₃: R = Et, 0.05 g, 0.37 mmol; Prⁱ, 0.04 g, 0.30 mmol] showed essentially quantitative formation of the PCl(NR₂)₂NPh-AlCl₃ complexes. The ²⁷Al NMR spectra show no indication of the presence of AlCl₄⁻ (δ 102, $\Delta v_{\pm} < 30$ Hz).⁴ The spectra were unchanged when solutions of PCl(NR₂)₂NPh-AlCl₃ were heated to 60 °C for 24 h.

Exposure of PCl(NR₂)₂NPh-AlCl₃ Complexes to Air.— Ground crystals of PCl(NR₂)₂NPh-AlCl₃ (R = Et or Prⁱ) were exposed to the air for 10 min. Dissolution of the material in CDCl₃ (used as received) left a white precipitate. The solution was examined by NMR spectroscopy, and the only phosphoruscontaining species present was tentatively characterised as bis-(dialkylamino)(phenylimino)phosphinic hydroxide P(NR₂)₂-(NPh)OH. NMR: ³¹P, R = Et, δ 39; Prⁱ, 35; ¹³C, R = Et, δ 136.4 (*i*), 129.6 (*m*), 125.1 (*p*), 120.6 (*o*), 41.3 and 13.1; Prⁱ, 136.5 (*i*), 129.2 (*m*), 125.2 (*p*), 122.4 (*o*) (J_{PC} = 6.8 Hz), 50.5 and 23.1; ¹H, R = Et, δ 9.57 (1 H) (concentration dependent, range 9.5–11.4), 7.3–7.1 (5 H, aromatic), 3.4–3.2 (8 H, dq) and 1.3 (12 H, t, J_{HH} = 7.2); Prⁱ, 11.45 (1 H, d, J_{HP} = 14) (concentration dependent), 7.5–7.0 (5 H, aromatic), 4.1–4.0 (4 H, m), 1.59

J. CHEM. SOC. DALTON TRANS. 1991

(12 H, d, $J_{\rm HH}$ = 6.9) and 1.47 (12 H, d, $J_{\rm HH}$ = 6.8 Hz). These compounds could not be isolated. A small amount of aluminium was present in the solution as AlCl₄⁻: ²⁷Al, δ 102 ($\Delta v_{\frac{1}{2}} < 30$ Hz).

Reaction of $PCl(NPr^{i}_{2})_{2}NPh$ with $Ag(SO_{3}CF_{3})$. Preparation and Isolation of P(NPrⁱ₂)₂[NPh(SO₂CF₃)]O.—A solution of PCl(NPrⁱ₂)₂NPh (0.743 g, 2.08 mmol) in CH₂Cl₂ (20 cm³) was added to Ag(SO₃CF₃) (0.534 g, 2.08 mmol). Reaction occurred immediately and the solution was stirred overnight. The CH_2Cl_2 was removed, hexane (20 cm³) was added, and the resulting pale yellow solution was decanted from the white precipitate. Slow removal of the solvent yielded an off-white air-stable solid, which was recrystallised six times from hexane and characterised as bis(diisopropylamino)[phenyl(trifluoromethylsulphonyl)amino]phosphine oxide, yield 0.343 g, 0.73 mmol (35%), m.p. 117–118 °C. Mass spectrum [m/z] (relative intensity)]: 471 (1) and 247 (100). NMR: ³¹P, δ 13; ¹³C, δ 136.4 (*i*), 131.9 (*m*), 129.9 (*p*), 129.6 (*o*), 120.3 (${}^{1}J_{CF} = 325$ Hz), 48.7, 23.6 and 23.0. IR: 1600w, 1255s, 1225w, 1200s, 1170w, 1130m, 1070w, 1020s, 970s, 940s, 915m, 880s, 870(sh), 705s, 680s, 620m, 605s, 580m, 565(sh), 550s, 525s, 445m, 405w and 385w $\rm cm^{-1}$ (Found: C, 48.45; H, 7.15; N, 9.00. Calc. for C₁₉H₃₃F₃N₃O₃PS: C, 48.40; H, 7.05; N, 8.90%).

NMR studies on the reaction mixture show the reaction to be quantitative. Similarly, reactions of $PCl(NEt_2)_2NPh$ with excess of $Ag(SO_3CF_3)$ gave a single product (³¹P, δ 14).

Reaction of $P(NPr_{2}^{i})_{2}^{+}SO_{3}CF_{3}^{-}$ with Phenyl Azide.—The azide (2.08 mmol) in $CH_{2}Cl_{2}$ (10 cm³) was added to a stirred solution of $P(NPr_{2}^{i})_{2}^{+}SO_{3}CF_{3}^{-}$ [prepared in situ, 2.09 mmol Ag(SO_{3}CF_{3}), 2.08 mmol $PCl(NPr_{2}^{i})_{2}$] in $CH_{2}Cl_{2}$ (20 cm³). A ³¹P NMR spectrum of the reaction mixture revealed $P(NPr_{2}^{i})_{2}^{-}$ [NPh(SO₂CF₃)]O to be the principal (>80%) component; $P(NPr_{2}^{i})_{2}^{+}$ and $PCl(NPr_{2}^{i})_{2}$ were the only other phosphoruscontaining species present.

X-Ray Data Collection, Solution and Refinement.-Suitable crystals of P(NPrⁱ₂)₂[NPh(SO₂CF₃)]O (obtained by slow removal of solvent from hexane solutions) were mounted on glass fibres and affixed with epoxy glue. Crystals of PCl- $(NPr_{2}^{i})_{2}NPh AlCl_{3}$ (grown by slow solvent removal under vacuum from a hexane-CH2Cl2 solution) were mounted in thin-walled Pyrex capillaries under an atmosphere of N₂ in a dry-box. Unit-cell parameters were obtained from the setting angles of 25 accurately centred reflections having $15 < \theta < 18^{\circ}$. The choice of space groups was based on systematically absent reflections and confirmed by the program TRACER¹⁰ and the successful solution and refinement of the structures. The choice of the alternative centric space group $P2_1/m$ for PCl(NPrⁱ₂)₂-NPh. AlCl₃ was eliminated due to the lack of molecular symmetry that would be required. All pertinent crystallographic data are summarised in Table 1.

Data were collected at room temperature on an Enraf-Nonius CAD-4 diffractometer (MoK α , $\lambda = 0.709$ 30 Å, graphite monochromator) using the ω -2 θ scan technique. The stability of the crystals was monitored every 60 min using three standard reflections and no significant decay was observed. Data were corrected for Lorentz and polarisation effects and an empirical absorption correction was applied based on a series of ψ scans. Transmission coefficients are listed in Table 1. Scattering factors were taken from Cromer and Waber¹¹ and anomalous dispersion effects were included in F_{e} .¹²

The structures were solved using direct methods¹³ to find the heavier elements and the remaining non-hydrogen atoms were located from a series of Fourier difference maps. Hydrogen atoms were included in ideal positions as fixed contributions $(B_{iso} = 1.2B \text{ of the bonded atom})$ to the structure factors. The final choice of enantiomer for PCl(NPrⁱ₂)₂NPh-AlCl₃ was based on a slightly lower *R* factor (0.0295 vs. 0.0297). The model was refined using full-matrix least-squares techniques based on *F*,

Table 1 Crystal data collection and refinement parameters

Formula		CUENOR				
r officia M	$C_{18}\Pi_{33}AICI_4IN_3F$	$C_{19}\Pi_{33}\Gamma_{3}\Pi_{3}O_{3}\Gamma_{3}$				
M Succession	491.23	4/1.55				
Space group	$\frac{PZ_1}{2}$	PZ_1/n				
Z E(000)	2	4				
F(000)	510	1000				
a/A	8.719(2)	10.594(3)				
b/A	15.553(2)	13.222(2)				
c/A	9.509(2)	17.084(3)				
β/°	102.73(2)	97.71(2)				
U/\dot{A}^3	1257.8	2731.4				
Dimensions/mm	$0.36\times0.45\times0.55$	$0.33 \times 0.38 \times 0.50$				
$D_{\rm c}/{\rm g~cm^{-3}}$	1.30	1.32				
μ/cm^{-1}	5.8	2.4				
Transmission coefficients	0.972-0.999	0.922-1.000				
Range of h, k, l	$+11, -20, \pm 12$	$+12, +15, \pm 20$				
$2\theta_{max}/^{\circ}$	55	50				
Reflections measured	3357	4447				
Unique	3001	4204				
Observed	2643	2585				
$[I > 3\sigma(I)]$ parameters	244	271				
Goodness of fit ^a	1.11	1.00				
R^{b}	0.029	0.048				
R' ^c	0.030	0.046				
Maximum Δ/σ	0.07	0.03				
Largest feature in difference						
map/e Å ⁻³	0.26	0.26				
^a $[\Sigma w(F_0 - F_c)^2/(N_0 - N_v)]^{\frac{1}{2}}$ where $N_0, N_v =$ numbers of reflections						

" $[\sum w(|F_o| - |F_c|)^2/(N_o - N_v)]^2$ where $N_o, N_v =$ numbers of reflections and parameters. ^b $\Sigma ||F_o| - |F_c||/\Sigma |F_o|$. ^c $R' = [\Sigma w(|F_o| - |F_c|)^2/\Sigma (wF_o)^2]^{\frac{1}{2}}$.

minimising the function $\Sigma w(|F_o| - |F_e|)^2$, where the weight was defined as $w = [\sigma(F)^2 + (0.005F)^2 + 0.4]^{-1}$ in all structures.¹⁴ Positional parameters for non-hydrogen atoms are given in Tables 2 and 3, selected bond lengths and angles in Table 4.

Additional material available from the Cambridge Crystallographic Data Centre comprises H-atom coordinates, thermal parameters and remaining bond lengths and angles.

Results and Discussion

The Diamino(phenylimino)phosphinic Chloride-Trichloroaluminium System.—Phosphorus-31 NMR studies of reactions between bis(dialkylamino)phosphenium tetrachloroaluminate $[P(NR_2)_2^+ AlCl_4^-]$ and phenyl azide, and reactions of bis-(dialkylamino)(phenylimino)phosphinic chloride $[PCl(NR_2)_2^-$ NPh] with AlCl₃, show the formation of the covalent Lewis acid-base complex $PCl(NR_2)_2NPh\cdotAlCl_3$ 3 (R = Et or Pr^i , E = NPh), in essentially quantitative yield. The complexes have been isolated as analytically pure crystalline solids. In solution the ³¹P NMR chemical shifts of the complexes (R = Et, δ 48.5; Pr^i , 51) are deshielded in comparison with the corresponding free bases (R = Et, δ 12; Pr^i , -8). A broad (R = Et, 132; Pr^i , 122 Hz) signal is observed at δ 100 in the ²⁷Al NMR spectra,



characteristic of an iminophosphoryl unit bound to $AlCl_3$.⁵ In the solid state, a strong to medium IR band is observed for both compounds in the region of 400 cm⁻¹, and can be assigned to the iminophosphoryl–aluminium linkage. The mass spectra do not exhibit a molecular ion, but the base peak corresponds to the free base PCl(NR₂)₂NPh, which is typical for phosphoryl–AlCl₃ complexes.^{4b,5,15} The structure of the complex

PCl(NPrⁱ₂)₂NPh-AlCl₃ has been confirmed by X-ray crystallography and a view is shown in Fig. 1. In general, the bond lengths and angles are as might be expected for the units involved.^{16a} Interestingly, the *anti* orientation of the phenyl group with respect to the chlorine atom is in contrast to the *gauche* orientation observed for the derivative PCl(NEt₂)₂NPh-AlCl₃.^{16b} The conformations are likely under the control of steric factors.

The quantitative formation of the iminophosphoryl-AlCl₃ complexes is in contrast to the results of Sanchez and coworkers,^{2a} who speculated on the formation of the three-coordinate iminophosphonium cation 2a (Y = NEt₂, ³¹P NMR δ 37.1) together with the complex PCl(NEt₂)₂NPh·AlCl₃ (δ 48.5), in solution. In addition, they suggested that the cation 2a (observed ³¹P signal at δ 37.1) is produced quantitatively as a thermolysis product (50 °C) of the PCl(NEt₂)₂NPh·AlCl₃ complex.^{2a} However, we have discovered that the iminophosphoryl-AlCl₃ complexes are thermally stable above 60 °C. More importantly, they are extremely air sensitive (not noted in ref. 2a), and minimal air exposure results in hydrolysis to give $P(NR_2)_2(NPh)OH$ which has been spectroscopically characterised (³¹P NMR: R = Et, δ 39; Prⁱ, 35). The ¹³C NMR spectrum of P(NEt₂)₂(NPh)OH is identical to that reported for cation 2a (R = Et), as is the ¹H NMR spectrum except for a concentration-dependent signal in the range typical of acidic protons (signal not reported in ref. 2). Hydrolysis reaction mixtures display a sharp signal at δ 102 characteristic of AlCl₄, indicative of the presence of an ionic system. However, AlCl₄ can be detected at very low concentrations,¹⁷ and it is likely that much of the aluminium is contained in the hydrolysis precipitate. The iminophosphinic chlorides PCl(NR₂)₂NPh $(R = Et \text{ or } Pr^{i})$ are themselves hydrolytically unstable, and we observe the ${}^{31}P$ NMR signal at δ 39 as an intermediate in the full hydrolysis of PCl(NEt₂)₂NPh to P(NEt₂)₂(NHPh)O,¹⁸ in the absence of AlCl₃.

The Diamino(phenylimino)phosphinic Chloride-Silver Trifluoromethanesulphonate System.—The compound PCl(NR₂)₂-NPh (R = Et or Prⁱ) reacts with Ag(SO₃CF₃) in CH₂Cl₂ to give AgCl and P(NR₂)₂[NPh(SO₂CF₃)]O in essentially quantitative yield (³¹P NMR spectroscopy). The structure of P(NPrⁱ₂)₂[NPh(SO₂CF₃)]O has been determined by X-ray crystallography,* Fig. 2. Reaction of P(NPrⁱ₂)₂+SO₃CF₃⁻ with PhN₃ produced the same compound. The ³¹P NMR chemical shifts of the new triaminophosphoryl centres (R = Et, δ 14; Prⁱ, 13) are in the range typical for triaminophosphine oxides ¹⁹ [cf. P(NEt₂)₂(NHPh)O, δ 15]. Furthermore, the IR spectrum (R = Prⁱ) exhibits a strong peak at 1255 cm⁻¹, characteristic²⁰ of the P=O stretching mode and the mass spectrum [m/z 471, parent ion; 247, P(NPrⁱ₂)₂O⁺] is consistent with the structure.

The Three-co-ordinate Phosphonium Electronic Environment. —The compounds $PCl(NR_2)_2NPh\cdotAlCl_3$ and $P(NR_2)_2[NPh\cdot(SO_2CF_3)]O$ represent two covalent structural alternatives to the alkenic three-co-ordinate iminophosphonium cation 2a. The structure observed for $P(NPr_2)_2[NPh(SO_2CF_3)]O$ 5a provides a direct comparison with bis(diisopropylamino)[bis-(trimethylsilyl)methylene]phosphonium 2b triflate $[(Pr_2N)_2P=C(SiMe_3)_2]^+[SO_3CF_3]^{-1}$ Thermodynamic preference for the covalent phosphine oxide alternative 5 (the P=O bond has a very high bond energy, 544 kJ mol⁻¹)²¹ is manifest in the imino-

^{*} A feature worthy of note is the P–N(S) bond length [P–N(1) 1.785(3) Å], considerably longer than the other P–N(amino) bonds [P–N(2) 1.649(3) and P–N(3) 1.639(3) Å], and close to a standard P–N single bond (1.78 Å).^{16a} The S–N bond [1.628(3) Å] is relatively short demonstrating a preference for N(1) to π -bond with the centre of higher oxidation state and lesser π involvement (sulphur is bound to two oxygen centres, while phosphorus has two nitrogen and one oxygen centre).

Atom	x	у	Ζ	Atom	x	у	Z
Cl(1)	0.528 26(8)	0.396*	0.683 79(9)	C(6)	1.007 1(4)	0.479 4(3)	0.951 3(4)
Cl(2)	0.866 1(1)	0.693 02(6)	0.656 4(1)	C(7)	0.814 1(4)	0.461 3(2)	0.398 5(3)
Cl(3)	0.507 7(1)	0.597 79(7)	0.539 2(1)	C(8)	0.709 4(5)	0.499 7(3)	0.265 0(4)
Cl(4)	0.660 5(1)	0.62142(7)	0.910 31(9)	C(9)	0.967 1(4)	0.426 6(3)	0.369 7(4)
P	0.749 70(8)	0.399 39(6)	0.648 32(8)	C(10)	0.625 7(4)	0.327 5(2)	0.395 5(4)
Al	0.714 2(1)	0.596 48(7)	0.706 9(1)	C(11)	0.463 7(4)	0.362 0(3)	0.321 5(5)
N(1)	0.819 4(3)	0.490 1(2)	0.717 7(3)	C(12)	0.699 5(5)	0.2772(3)	0.289 0(4)
N(2)	0.7320(3)	0.396 8(2)	0.474 8(2)	C(13)	0.950 0(4)	0.264 7(2)	0.658 9(3)
N(3)	0.8352(3)	0.3139(2)	0.726 4(3)	C(14)	0.9121(5)	0.168 9(2)	0.643 2(4)
CÌÌ	0.979 1(3)	0.4940(2)	0.804 3(3)	C(15)	1.121 1(4)	0.2801(3)	0.730 6(5)
C(2)	1.102 4(4)	0.5176(2)	0.741 0(4)	C(16)	0.794 5(5)	0.284 6(3)	0.865 2(4)
C(3)	1.253 3(4)	0.5244(2)	0.826 5(4)	C(17)	0.935 3(6)	0.258 1(3)	0.980 9(4)
C(4)	1.281 0(4)	0.507 1(3)	0.971 3(5)	C(18)	0.665 4(5)	0.215 6(3)	0.840 3(5)
Cisi	1.159 5(5)	0.485 5(3)	1.033 4(4)		(-)	- (-)	

 Table 2
 Positional parameters and their estimated standard deviations (e.s.d.s)

* The y coordinate of Cl(1) was fixed to define the origin.

Table 3 Positional parameters and their e.s.d.s for $P(NPr_2)_2[NPh(SO_2CF_3)]O$

Atom	x	У	Ζ	Atom	x	у	Ζ
S	0.384 5(1)	0.132 69(8)	0.288 92(6)	C(5)	0.438 2(4)	0.378 6(3)	0.478 1(2)
Р	0.558 8(1)	0.274 56(8)	0.222 76(6)	C(6)	0.478 1(4)	0.319 2(3)	0.419 4(2)
F(1)	0.593 6(3)	0.087 5(2)	0.383 8(2)	C(7)	0.524 6(4)	0.236 9(3)	0.064 3(2)
F(2)	0.428 4(3)	0.101 9(2)	0.443 1(1)	C(8)	0.647 1(5)	0.287 7(4)	0.048 6(3)
F(3)	0.449 8(3)	-0.0275(2)	0.370 8(2)	C(9)	0.536 6(5)	0.121 4(4)	0.066 2(2)
O(1)	0.657 0(2)	0.195 9(2)	0.230 4(2)	C(10)	0.341 5(4)	0.320 8(3)	0.119 6(2)
O(2)	0.256 7(3)	0.135 9(3)	0.305 6(2)	C(11)	0.330 9(5)	0.407 4(4)	0.059 4(3)
O(3)	0.419 3(3)	0.075 3(2)	0.225 0(2)	C(12)	0.241 5(4)	0.238 5(4)	0.093 3(3)
N(1)	0.442 7(3)	0.246 4(2)	0.286 2(2)	C(13)	0.544 9(4)	0.481 3(3)	0.234 6(2)
N(2)	0.471 8(3)	0.278 2(3)	0.135 3(2)	C(14)	0.598 3(5)	0.543 2(3)	0.170 2(3)
N(3)	0.615 8(3)	0.385 1(2)	0.253 1(2)	C(15)	0.539 5(5)	0.547 1(3)	0.307 9(2)
C(1)	0.403 1(4)	0.313 4(3)	0.346 2(2)	C(16)	0.752 9(4)	0.398 0(3)	0.288 2(2)
C(2)	0.289 5(4)	0.365 1(3)	0.332 0(2)	C(17)	0.782 4(4)	0.346 7(4)	0.368 7(3)
C(3)	0.251 1(4)	0.424 8(4)	0.391 3(3)	C(18)	0.847 2(4)	0.365 8(4)	0.233 3(3)
C(4)	0.326 6(5)	0.431 4(3)	0.464 2(2)	C(19)	0.471 5(5)	0.071 3(3)	0.377 5(3)

Table 4 Selected bond lengths (Å) and angles (°)

PCl(NPr ⁱ ₂) ₂ NI	$PCl(NPr_{2}^{i})_{2}NPh AlCl_{3}$ $P(NPr_{2}^{i})_{2}[NPh(SO_{2}CF_{2})_{3}]$		[NPh(SO ₂ CF ₃)]O	PCl(NPr ⁱ ₂) ₂ NPh•AlCl ₃		$P(NPr_{2}^{i})_{2}[NPh(SO_{2}CF_{3})]O$	
P(1) - N(1)	1.618(3)		1.785(3)	N(1)-P(1)-Cl(1)	104.0(1)	N(1)-P(1)-O(1)	109.7(2)
P(1)-Cl(1)	2.033(2)	P-O(1)	1.464(3)	N(1)-P(1)-N(2)	112.3(1)		102.1(1)
P(1)-N(2)	1.624(2)		1.649(3)	N(1)-P(1)-N(3)	115.7(1)		104.5(2)
P(1) - N(3)	1.622(3)		1.639(3)	Cl(1)-P(1)-N(2)	106.69(9)	O(1)-P(1)-N(2)	113.9(2)
N(2)-C(7)	1.508(4)		1.504(6)	Cl(1)-P(1)-N(3)	105.0(2)	O(1)-P(1)-N(3)	112.2(2)
N(2)-C(10)	1.511(5)		1.481(6)	N(2)-P(1)-N(3)	112.1(1)		113.5(2)
N(3)C(13)	1.511(4)		1.490(5)	P(1)-N(2)-C(7)	121.6(2)		119.9(2)
N(3)C(16)	1.511(5)		1.505(5)	P(1)-N(2)-C(10)	115.8(2)		124.7(2)
C(7)-C(8)	1.514(5)		1.517(7)	C(7)-N(2)-C(10)	122.6(2)		115.5(3)
C(7)-C(9)	1.519(5)		1.533(6)	P(1)-N(3)-C(13)	120.0(3)		122.8(2)
C(10)-C(11)	1.530(5)		1.535(6)	P(1)-N(3)-C(16)	118.4(2)		121.5(3)
C(10)-C(12)	1.530(6)		1.543(6)	C(13)-N(3)-C(16)	121.6(3)		114.8(3)
C(13)-C(14)	1.527(5)		1.538(6)	P(1)-N(1)-C(1)	119.6(2)		126.0(2)
C(13)-C(15)	1.517(5)		1.533(6)	P(1)-N(1)-Al	127.5(1)		120.3(2)
C(16)–C(17)	1.514(5)		1.528(7)	Cl(2)-Al-N(1)	108.21(9)	O(2)-S-N(1)	110.6(2)
C(16)-C(18)	1.535(6)		1.519(6)	Cl(3)-Al-N(1)	111.78(9)	O(3)-S-N(1)	109.5(2)
N(1)-C(1)	1.455(3)		1.458(5)	Cl(4)-Al-N(1)	108.26(9)	C(19)-S-N(1)	106.2(2)
N(1)-Al	1.884(3)	N(1)-S	1.628(3)	Cl(2)-Al-Cl(3)	106.90(6)	O(2)-S-O(3)	121.6(2)
Cl(2)-Al	2.126(1)	S-O(2)	1.422(3)	Cl(2)-Al-Cl(4)	110.49(6)	O(2)-S-C(19)	103.4(2)
Cl(3)-Al	2.127(1)	S-O(3)	1.418(3)	Cl(3)-Al-Cl(4)	111.18(6)	O(3)-S-C(19)	104.1(2)
Cl(4)-Al	2.124(1)	S-C(19)	1.851(4)				

derivative. Therefore, we believe that isolation of the first stable three-co-ordinate methylenephosphonium centre, **2b**, is made possible by steric protection ('kinetic stabilisation') of the P=C bond, which is enhanced by the unusually large twist angle of $60^{\circ.1}$ The steric shield offered by the phenyl substituent on the nitrogen centre in **2a** is not as uniform or as extensive as two SiMe₃ units, and evidently allows interaction between the cation and the anion for oxygen transfer. Thermodynamic control of the products from the imino-system is demonstrated



by the fact that $P(NPr_{2}^{i})_{2}[NPh(SO_{2}CF_{3})]O$ was obtained from $P(NPr_{2}^{i})_{2}^{+}$ SO₃CF₃⁻, by means of a Staudinger reaction (the triflate anion is present prior to formation of the phosphonium

1619



Fig. 1 Crystallographic view of PCl(NPrⁱ₂)₂NPh·AlCl₃



Fig. 2 Crystallographic view of P(NPrⁱ₂)₂[NPh(SO₂CF₃)]O

cation), and from the iminophosphoryl centre [triflate anion is released during reaction with $Ag(SO_3CF_3)$].

Alternative Ionic Structures from the Phosphoryl Units.— Retention of a covalent phosphoryl environment for PCl- $(NR_2)_2NPh$ -AlCl₃ is consistent with the derivative PCl- $(NPr_2)_2S$ -AlCl₃.⁴ However, other (when R = Me or Et) thio- and seleno-phosphoryl chloride analogues adopt novel dimeric diphosphonium heterocyclic structures 4 in the solid state.⁴ Clearly, this delicate thermodynamic ionic/covalent balance is again critically influenced by the degree of steric shielding at the diaminophosphoryl centre. Furthermore, it is interesting that the bis(dimethylamino)(methylimino)- and bis-(diethylamino)(ethylimino)-phosphinic halides PX(NR_2)_2NR (R = Me or Et; X = Cl or Br) undergo an autoionisation to the analogous diphosphonium heterocyclic phosphetidines,³ while a covalent monomeric phosphoryl structure is observed for the more sterically protected PCl(NPr₁)_2NPh.²²

Conclusion

The three-co-ordinate diaminoiminophosphonium cation is thermodynamically unstable: the tetrachloroaluminate salt

prefers the thermally stable diaminoiminophosphinic chloridetrichloroaluminium Lewis complex covalent alternative, and the triflate salt adopts the triaminophosphine oxide covalent alternative, despite the presence of diisopropylamino-substituents. Therefore, steric shielding at carbon is believed to be responsible for the observed stability of the analogous diaminomethylenephosphonium cation.¹

Acknowledgements

We thank the Natural Sciences and Engineering Research Council of Canada and the donors of the Petroleum Research Fund, administered by the American Chemical Society, for financial support and the Atlantic Region Magnetic Resonance Centre for obtaining the NMR spectra.

References

- A. Igau, A. Baceiredo, H. Grutzmacher, H. Pritzkow and G. Bertrand, J. Am. Chem. Soc., 1989, 111, 6853.
 (a) M. R. Marre, M. Sanchez and R. Wolf, Phosphorus Sulfur, 1982,
- 2 (a) M. R. Marre, M. Sanchez and R. Wolf, *Phosphorus Sulfur*, 1982, 13, 327; M. Sanchez, M. R. Marre, J. F. Brazier, J. Bellan and R. Wolf, *Phosphorus Sulfur*, 1983, 14, 331; (b) see also, M. R. Marre, M. Sanchez and R. Wolf, *J. Chem. Soc., Chem. Commun.*, 1984, 566; M. R. Mazieres, M. Sanchez, J. Bellan and R. Wolf, *Phosphorus Sulfur*, 1986, 26, 97.
- A. N. Chernega, M. Yu. Antipin, Yu. T. Struchkov, I. E. Boldeskul,
 A. P. Marchenko and A. M. Pinchuk, *Zh. Strukt. Khim.*, 1987, 28, 135;
 A. P. Marchenko, V. V. Miroshnichenko, V. A. Kovenya, A. M. Pinchuk, A. N. Chernega, M. Yu. Antipin and Yu. T. Struchkov, *Zh. Obshch. Khim.*, 1988, 58, 1758.
- 4 N. Burford, R. E. v. H. Spence and R. D. Rogers, (a) J. Am. Chem. Soc., 1989, 111, 5006; (b) J. Chem. Soc., Dalton Trans., 1990, 3611.
- 5 N. Burford, R. E. v. H. Spence, J. M. Whalen, R. D. Rogers and J. F. Richardson, *Organometallics*, 1990, 9, 2854.
- 6 A. B. Burg and P. J. Slota, jun., J. Am. Chem. Soc., 1958, 80, 1107.
- 7 A. H. Cowley, R. A. Kemp, J. G. Lasch, N. C. Norman, C. A. Stewart, B. R. Whittlesey and T. C. Wright, *Inorg. Chem.*, 1986, **25**, 740; R. B. King and P. M. Sundaram, *J. Org. Chem.*, 1984, **49**, 1784; M. J. S. Gynane, A. Hudson, M. F. Lappert, P. P. Power and H. Goldwhite, *J. Chem. Soc.*, *Dalton Trans.*, 1980, 2428.
- 8 A. L. Wayda and J. L. Dye, J. Chem. Educ., 1985, 62, 356.
- 9 A. H. Cowley and R. A. Kemp, Chem. Rev., 1985, 85, 367.
- 10 S. L. Lawson and R. A. Jacobsen, Ames Laboratory Report IS-1141, USABC, Iowa State University, Ames, IA, 1965.
- 11 D. T. Cromer and J. T. Waber, *International Tables for X-Ray Crystallography*, Kynoch Press, Birmingham, 1974, vol. 4, Table 2.2B (Present distributor: Kluwer Academic Publishers, Dordrecht and Boston).
- 12 D. T. Cromer, International Tables for X-Ray Crystallography, Kynoch Press, Birmingham, 1974, vol. 4, Table 2.3.1.
- 13 SDP/VAX, B. A. Frenz, in *Computing in Crystallography*, eds. H. Schenk, R. Olthof-Hazekamp, H. van Koningsveld and G. C. Bassi, Delft University Press, Delft, 1978, pp. 64–71.
- 14 R. C. G. Killean and J. L. Lawrence, *Acta Crystallogr., Sect. B*, 1973, **25**, 1750.
- 15 N. Burford, B. W. Royan, R. E. v. H. Spence and R. D. Rogers, J. Chem. Soc., Dalton Trans., 1990, 2111.
- 16 (a) D. E. C. Corbridge, The Structural Chemistry of Phosphorus, Elsevier, Amsterdam, 1974; (b) M. R. Mazieres, T. C. Kim, R. Wolf and M. Sanchez, Z. Kristallogr., 1988, 184, 147.
- 17 J. J. Delpuech, in NMR of Newly Accessible Nuclei, ed. P. Laszlo, Academic Press, New York, 1983, vol. 2, p. 153; J. W. Akitt, in Multinuclear NMR, ed. J. Mason, Plenum, New York, 1987, p. 259.
- 18 V. Gutmann, Ch. Kemenater and K. Utvary, *Monatsch-Chem.*, 1965, 96, 836.
- 19 K. R. Dixon, in *Multinuclear NMR*, ed. J. Mason, Plenum, New York, 1987, p. 369.
- 20 L. J. Bellamy, The Infra-red Spectra of Complex Molecules, Chapman and Hall, London, 1975.
- 21 J. E. Huheey, *Inorganic Chemistry*, 3rd edn., Harper & Row, New York, 1983.
- 22 N. Burford, R. E. v. H. Spence and J. F. Richardson, unpublished work.

Received 6th December 1990; Paper 0/05486K