

Articles

Synthesis, Emission, and Molecular Orbital Studies of Luminescent Hafnium Thiolate Complexes. Crystal Structures of $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Hf}(\text{SR})_2$ ($\text{R} = \text{}^n\text{Bu}$, C_6H_5 , $\text{C}_6\text{H}_4\text{OMe-}p$)

Vivian Wing-Wah Yam,* Gui-Zhong Qi, and Kung-Kai Cheung

Department of Chemistry, The University of Hong Kong, Pokfulam Road, Hong Kong, P.R. China

Received July 14, 1998

A series of luminescent hafnium thiolate complexes $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Hf}(\text{SR})_2$ ($\text{R} = \text{}^n\text{Bu}$, C_6H_5 , $\text{C}_6\text{H}_4\text{OMe-}p$, $\text{C}_6\text{H}_4\text{}^i\text{Bu-}p$) have been synthesized by the reaction of $(\eta^5\text{-C}_5\text{Me}_5)_2\text{HfCl}_2$ with LiSR and shown to exhibit rich luminescence behavior. The crystal structures of $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Hf}(\text{S}^n\text{Bu})_2$, $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Hf}(\text{SC}_6\text{H}_5)_2$, and $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Hf}(\text{SC}_6\text{H}_4\text{OMe-}p)_2$ have been determined. Fenske–Hall molecular orbital calculations on the hafnium thiolate complexes reveal a thiolate-based HOMO and a LUMO that is mainly composed of hafnium d character.

Introduction

In contrast to the metal-to-ligand charge-transfer (MLCT) excited states, which have been the dominant subjects of study in inorganic photochemistry over the past two decades,¹ complexes that show ligand-to-metal charge-transfer (LMCT) emissive states are very scarce.² We have explored the class of d^0 zirconium chalcogenolates,³ as part of our efforts in the study of metal chalcogenides and chalcogenolates,⁴ because the chalcogenolates, being good σ -donor type ligands, together with the good electron-accepting ability of d^0 metal centers, may give rise to low-lying emissive LMCT

excited states. To extend our research efforts in luminescent metal chalcogenolates, herein, we report the synthesis and luminescence studies of a series of hafnium thiolates, $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Hf}(\text{SR})_2$ ($\text{R} = \text{}^n\text{Bu}$, C_6H_5 , $\text{C}_6\text{H}_4\text{OMe-}p$, $\text{C}_6\text{H}_4\text{}^i\text{Bu-}p$). The X-ray crystal structures of $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Hf}(\text{S}^n\text{Bu})_2$, $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Hf}(\text{SC}_6\text{H}_5)_2$, and $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Hf}(\text{SC}_6\text{H}_4\text{OMe-}p)_2$ have been determined. The electronic structures and the nature of the excited states of these luminescent complexes have been probed by Fenske–Hall molecular orbital calculations.

Experimental Section

Materials and Reagents. Bis(pentamethylcyclopentadienyl)hafnium dichloride was purchased from Strem Chemicals Inc. 1-Butanethiol, thiophenol, 4-methoxythiophenol, and 4-*tert*-butylthiophenol were obtained from Lancaster Synthesis Ltd. *n*-Butyllithium was purchased from Aldrich Chemical Co. Analytical grade toluene, dimethoxyethane, and hexane were dried over sodium and distilled over sodium benzophenone ketyl using standard procedures before use.

Synthesis of Hafnium Thiolate Complexes. All reactions and manipulations were carried out under strictly anaerobic and anhydrous conditions using standard Schlenk technique.

$(\eta^5\text{-C}_5\text{Me}_5)_2\text{Hf}(\text{S}^n\text{Bu})_2$ (1). To a solution of $(\eta^5\text{-C}_5\text{Me}_5)_2\text{HfCl}_2$ (0.14 g, 0.27 mmol) in DME (15 mL) at 0 °C was added LiS^nBu (0.60 mmol) in DME (20 mL), prepared *in situ* at 0 °C from HS^nBu (0.054 g, 0.60 mmol) and $^n\text{BuLi}$ (0.60 mmol in *n*-hexane). The reaction mixture was slowly warmed to 50 °C and stirred at that temperature for 48 h. After removal of the solvent under reduced pressure, the solid residue was extracted with toluene. The toluene extract was subsequently concentrated and addition of hexane followed by cooling gave **1** as pale yellow crystals. Yield: 0.11 g (65%). ^1H NMR (300 MHz, C_6D_6 , 298 K, relative to Me_4Si): δ 0.95 (t, 6H, CH_3), 1.53 (m, 4H, CH_2CH_2), 1.73 (m, 4H, $\text{CH}_2\text{CH}_2\text{CH}_2$), 2.06 (s, 30H, $\text{C}_5\text{-Me}_5$), 2.99 (t, 4H, CH_2S). UV–vis in toluene (298 K), $\lambda_{\text{max}}/\text{nm}$

(1) (a) Kalyanasundaram, K. *Photochemistry of Polypyridine and Porphyrin Complexes*; Academic Press: San Diego, 1992. (b) Roundhill, D. M. *Photochemistry and Photophysics of Metal Complexes*; Plenum Press: New York, 1994.

(2) See for example: (a) Heinselman, K. S.; Hopkins, M. D. *J. Am. Chem. Soc.* **1995**, *117*, 12340. (b) Lee, Y. F.; Kirchhoff, J. R. *J. Am. Chem. Soc.* **1994**, *116*, 3599. (c) Pfennig, B. W.; Thompson, M. E.; Bocarsly, A. B. *J. Am. Chem. Soc.* **1989**, *111*, 8947. (d) Pfennig, B. W.; Thompson, M. E.; Bocarsly, A. B. *Organometallics* **1993**, *12*, 649. (e) Vlcek, A., Jr. *Chemtracts: Inorg. Chem.* **1993**, *5*, 141. (f) Thorn, D. L.; Harlow, R. L. *Inorg. Chem.* **1992**, *31*, 3917. (g) Paulson, S.; Sullivan, B. P.; Caspar, J. V. *J. Am. Chem. Soc.* **1992**, *114*, 6905. (h) Bandy, J. A.; Cloke, F. G. N.; Cooper, G.; Day, J. P.; Girling, R. B.; Graham, R. G.; Green, J. C.; Grinter, R.; Perutz, R. N. *J. Am. Chem. Soc.* **1988**, *110*, 5039.

(3) Yam, V. W. W.; Qi, G. Z.; Cheung K. K. *J. Chem. Soc., Dalton Trans.* **1998**, 1819.

(4) See for example: (a) Yam, V. W. W.; Lee, W. K.; Lai, T. F. *J. Chem. Soc., Chem. Commun.* **1993**, 1571. (b) Yam, V. W. W.; Lo, K. K. W.; Cheung, K. K. *Inorg. Chem.* **1996**, *35*, 3549. (c) Yam, V. W. W.; Lo, K. K. W.; Wang, C. R.; Cheung, K. K. *Inorg. Chem.* **1996**, *35*, 5116. (d) Yam, V. W. W.; Yeung, P. K. Y.; Cheung, K. K. *J. Chem. Soc., Dalton Trans.* **1994**, 2587. (e) Yam, V. W. W.; Yeung, P. K. Y.; Cheung, K. K. *J. Chem. Soc., Chem. Commun.* **1995**, 267. (f) Yam, V. W. W.; Yeung, P. K. Y.; Cheung, K. K. *Angew. Chem., Int. Ed. Engl.* **1996**, *35*, 739. (g) Yam, V. W. W.; Chan, C. L.; Cheung, K. K. *J. Chem. Soc., Dalton Trans.* **1996**, 4019. (h) Yam, V. W. W.; Wong, K. M. C.; Cheung, K. K. *Organometallics* **1997**, *16*, 1729. (i) Yam, V. W. W. *J. Photochem. Photobiol. A* **1997**, *106*, 75. (j) Yam, V. W. W.; Lo, K. K. W.; Wang, C. R.; Cheung, K. K. *J. Phys. Chem. A* **1997**, *101*, 4666. (k) Yam, V. W. W.; Lo, K. K. *Comments Inorg. Chem.* **1997**, *19*, 209.

($\epsilon_{\text{max}}/\text{dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$): 318 (4300), 368 (2300). Positive ion EI-MS: $m/z = 628 \{M\}^+$, 571 $\{M - C_4H_9\}^+$, 538 $\{M - SC_4H_9\}^+$, 481 $\{M - SC_4H_9 - C_4H_9\}^+$. Anal. Found: C, 53.41; H, 7.67. Calcd for $C_{28}H_{48}S_2Hf$: C, 53.56; H, 7.65.

($\eta^5\text{-C}_5\text{Me}_5$)₂Hf(SC₆H₅)₂ (2). The procedure was similar to that of **1** except HSC₆H₅ (0.066 g, 0.60 mmol) was used in place of HSⁿBu to give **2** as yellow crystals. Yield: 0.12 g (67%). ¹H NMR (300 MHz, C₆D₆, 298 K, relative to Me₄Si): δ 1.93 (s, 30H, C₅Me₅), 6.89 (t, $J = 8$ Hz, 2H, aryl protons *para* to S), 6.97 (t, $J = 8$ Hz, 4H, aryl protons *meta* to S), 7.86 (d, $J = 8$ Hz, 4H, aryl protons *ortho* to S). UV-vis in toluene (298 K), $\lambda_{\text{max}}/\text{nm}$ ($\epsilon_{\text{max}}/\text{dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$): 332 (11000), 392 (7600). Positive ion EI-MS: $m/z = 668 \{M\}^+$, 559 $\{M - SC_6H_5\}^+$, 533 $\{M - Cp^*\}^+$. Anal. Found: C, 57.56; H, 5.92. Calcd for $C_{32}H_{40}S_2Hf$: C, 57.58; H, 5.99.

($\eta^5\text{-C}_5\text{Me}_5$)₂Hf(SC₆H₄OMe-*p*)₂ (3). The procedure was similar to that of **1** except HSC₆H₄OMe-*p* (0.084 g, 0.60 mmol) was used in place of HSⁿBu to give **3** as pale yellow green crystals. Yield: 0.13 g (66%). ¹H NMR (300 MHz, C₆D₆, 298 K, relative to Me₄Si): δ 1.97 (s, 30H, C₅Me₅), 3.25 (s, 6H, OMe), 6.63 (d, $J = 8$ Hz, 4H, aryl protons *meta* to S), 7.77 (d, $J = 8$ Hz, 4H, aryl protons *ortho* to S). UV-vis in toluene (298 K), $\lambda_{\text{max}}/\text{nm}$ ($\epsilon_{\text{max}}/\text{dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$): 330 (8800), 396 (5600). Positive ion EI-MS: $m/z = 728 \{M\}^+$, 589 $\{M - SC_6H_4OMe-p\}^+$. Anal. Found: C, 56.28; H, 6.07. Calcd for $C_{34}H_{44}O_2S_2Hf$: C, 56.16; H, 6.06.

($\eta^5\text{-C}_5\text{Me}_5$)₂Hf(SC₆H₄Bu-*p*)₂ (4). The procedure was similar to that of **1** except HSC₆H₄Bu-*p* (0.10 g, 0.60 mmol) was used in place of HSⁿBu to give **4** as pale yellow crystals. Yield: 0.15 g (71%). ¹H NMR (300 MHz, C₆D₆, 298 K, relative to Me₄Si): δ 1.14 (s, 18H, 'Bu), 1.99 (s, 30H, C₅Me₅), 7.03 (d, $J = 8$ Hz, 4H, aryl protons *meta* to S), 7.84 (d, $J = 8$ Hz, 4H, aryl protons *ortho* to S). UV-vis in toluene (298 K), $\lambda_{\text{max}}/\text{nm}$ ($\epsilon_{\text{max}}/\text{dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$): 332 (9700), 394 (6200). Positive ion EI-MS: $m/z = 780 \{M\}^+$, 615 $\{M - SC_6H_4Bu-p\}^+$. Anal. Found: C, 61.77; H, 7.29. Calcd for $C_{40}H_{56}S_2Hf$: C, 61.66; H, 7.19.

Physical Measurements and Instrumentation. UV-visible spectra were obtained on a Hewlett-Packard 8452A diode array spectrophotometer and steady-state excitation and emission spectra on a Spex Fluorolog 111 spectrofluorometer equipped with a Hamamatsu R-928 photomultiplier tube detector. Low-temperature (77 K) spectra were recorded by using an optical Dewar sample holder. ¹H NMR spectra were recorded on a Bruker DPX300 Fourier transform NMR spectrometer. Chemical shifts were reported relative to tetramethylsilane. Positive ion EI mass spectra were recorded on a Finnigan MAT95 mass spectrometer.

Crystal Structure Determination. Crystal Data for 1: $C_{28}H_{48}S_2Hf$, $M_r = 627.30$, monoclinic, space group $C2/c$ (No. 15), $a = 14.843(3)$ Å, $b = 13.233(2)$ Å, $c = 16.332(5)$ Å, $\beta = 116.07(2)^\circ$, $V = 2881(1)$ Å³, $Z = 4$, $D_c = 1.446$ g cm⁻³, $\mu(\text{Mo K}\alpha) = 37.72$ cm⁻¹, $F(000) = 1280$, $T = 301$ K. A pale yellow crystal of dimensions $0.15 \times 0.15 \times 0.30$ mm mounted on a glass fiber was used for data collection at 28 °C on an Enraf-Nonius CAD4 diffractometer with graphite-monochromated Mo K α radiation ($\lambda = 0.71073$ Å) using ω -2 θ scans with ω -scan angle $(0.80 + 0.35 \tan \theta)^\circ$ at a scan speed of 1.50 – 8.24 deg min⁻¹. Intensity data (in the range of $2\theta_{\text{max}} = 48^\circ$; h , 0 to 17; k , 0 to 15; l , -18 to 18 and 3 standard reflections measured after every 300 reflections showed decay of 4.75%) were corrected for decay and for Lorentz and polarization effects and empirical absorption corrections based on the ψ -scan of four strong reflections (minimum and maximum transmission factors 0.836 and 1.000). A total of 2479 reflections were measured, of which 2416 were unique and $R_{\text{int}} = 0.010$; 2167 reflections with $I > 3\sigma(I)$ were considered observed and used in the structural analysis. The space group was determined on the basis of systematic absences and a statistical analysis of intensity distribution, and the successful refinement of the

structure was solved by direct methods (*SIR92*)^{5a} and expanded by the Fourier method; refinement was by full-matrix least squares using the software package *TeXsan*^{5b} on a Silicon Graphics Indy computer. A crystallographic asymmetric unit consists of half of one molecule with the Hf atom at special position. All 16 non-H atoms of the asymmetric unit were refined anisotropically, and the 24 H atoms at calculated positions with thermal parameters equal to 1.3 times that of the attached C atoms were not refined. Convergence for 141 variable parameters by least-squares refinement on F with $w = 4F_o^2/\sigma^2(F_o^2)$, where $\sigma^2(F_o^2) = [\sigma^2(I) + (0.020 F_o^2)^2]$ for 2167 reflections with $I > 3\sigma(I)$ was reached at $R = 0.019$ and $wR = 0.025$ with a goodness-of-fit of 1.58. $(\Delta\rho)_{\text{max}} = 0.01$. The final difference Fourier map was featureless, with maximum positive and negative peaks of 0.51 and 0.55 e Å⁻³, respectively.

Crystal Data for 2: $C_{32}H_{40}S_2Hf$, $M_r = 667.28$, monoclinic, space group $C2/c$ (No. 15), $a = 14.414(5)$ Å, $b = 11.466(3)$ Å, $c = 18.190(3)$ Å, $\beta = 103.18(2)^\circ$, $V = 2927(1)$ Å³, $Z = 4$, $D_c = 1.732$ g cm⁻³, $\mu(\text{Mo K}\alpha) = 37.18$ cm⁻¹, $F(000) = 1344$, $T = 301$ K. A yellow crystal of dimensions $0.25 \times 0.20 \times 0.35$ mm mounted on a glass fiber was used for data collection at 28 °C on an Enraf-Nonius CAD4 diffractometer with graphite-monochromated Mo K α radiation ($\lambda = 0.71073$ Å) using ω -2 θ scans with ω -scan angle $(0.66 + 0.35 \tan \theta)^\circ$ at a scan speed of 1.18 – 5.49 deg min⁻¹. Intensity data (in the range of $2\theta_{\text{max}} = 50^\circ$; h , 0 to 17; k , 0 to 13; l , -21 to 21 and 3 standard reflections measured after every 300 reflections showed no decay) were corrected for Lorentz and polarization effects and empirical absorption corrections based on the ψ -scan of four strong reflections (minimum and maximum transmission factors 0.833 and 1.000). A total of 2831 reflections were measured, of which 2732 were unique and $R_{\text{int}} = 0.011$; 2484 reflections with $I > 3\sigma(I)$ were considered observed and used in the structural analysis. The space group was determined on the basis of systematic absences and a statistical analysis of intensity distribution, and the successful refinement of the structure was solved by direct methods (*SIR92*)^{5a} and expanded by the Fourier method; refinement was by full-matrix least squares using the software package *TeXsan*^{5b} on a Silicon Graphics Indy computer. A crystallographic asymmetric unit consists of half of one molecule with the Hf atom at special position. All 18 non-H atoms of the asymmetric unit were refined anisotropically, and the 20 H atoms at calculated positions with thermal parameters equal to 1.3 times that of the attached C atoms were not refined. Convergence for 159 variable parameters by least-squares refinement on F with $w = 4F_o^2/\sigma^2(F_o^2)$, where $\sigma^2(F_o^2) = [\sigma^2(I) + (0.028 F_o^2)^2]$ for 2484 reflections with $I > 3\sigma(I)$ was reached at $R = 0.024$ and $wR = 0.038$ with a goodness-of-fit of 2.17. $(\Delta\rho)_{\text{max}} = 0.02$. The final difference Fourier map was featureless, with maximum positive and negative peaks of 0.61 and 1.18 e Å⁻³, respectively.

Crystal Data for 3: $C_{32}H_{44}O_2S_2Hf$, $M_r = 727.33$, monoclinic, space group $C2/c$ (No. 15), $a = 18.034(3)$ Å, $b = 11.105(2)$ Å, $c = 16.835(3)$ Å, $\beta = 109.69(2)^\circ$, $V = 3174(1)$ Å³, $Z = 4$, $D_c = 1.522$ g cm⁻³, $\mu(\text{Mo K}\alpha) = 34.40$ cm⁻¹, $F(000) = 1472$, $T = 301$ K. A yellow green crystal of dimensions $0.25 \times 0.15 \times 0.30$ mm on a glass fiber was used for data collection at 28 °C on an MAR diffractometer with a 300 mm image plate detector using graphite-monochromated Mo K α radiation ($\lambda = 0.71073$ Å). Data collection was made with 3° oscillation (60 images) at 120 mm distance and 280 s exposure. The images were interpreted and intensities integrated using the program *DENZO*.^{5c} A total of 3073 unique reflections were obtained

(5) (a) *SIR 92*: Altomare, A.; Cascarano, M.; Giacovazzo, C.; Guagliardi, A.; Burla, M. C.; Polidori, G.; Camalli, M. *J. Appl. Crystallogr.* **1994**, *27*, 435. (b) *TeXsan*: Crystal Structure Analysis Package; Molecular Structure Corporation: The Woodlands, TX, 1985 & 1992. (c) *DENZO*: In *The HKL Manual—A description of programs DENZO, XDISP, and SCALEPACK*; written by Gewirth, D. with the cooperation of the program authors Orwinowski, Z., and Minor, W.; Yale University: New Haven, CT, 1995.

from a total of 13 424 measured reflections ($R_{\text{int}} = 0.033$); 2863 reflections with $I > 3\sigma(I)$ were considered observed and used in the structural analysis. These reflections were in the range h , 0 to 21; k , 0 to 13; l , -20 to 20 with $2\theta_{\text{max}} = 51.0^\circ$. The space group was determined on the basis of systematic absences and a statistical analysis of intensity distribution, and the successful refinement of the structure was solved by direct methods (*SIR92*)^{5a} and expanded by the Fourier method; refinement was by full-matrix least squares using the software package *TeXsan*^{5b} on a Silicon Graphics Indy computer. One crystallographic asymmetric unit consists of half of one molecule with the Hf atom at special position. In the least-squares refinement, all 20 non-H atoms were refined anisotropically, and the 22 H atoms at calculated positions with thermal parameters equal to 1.3 times that of the attached C atoms were not refined. Convergence for 177 variable parameters by least-squares refinement on F with $w = 4F_o^2/\sigma^2(F_o^2)$, where $\sigma^2(F_o^2) = [\sigma^2(I) + (0.032F_o^2)^2]$ for 2863 reflections with $I > 3\sigma(I)$ was reached at $R = 0.025$ and $wR = 0.044$ with a goodness-of-fit of 1.92. $(\Delta/\sigma)_{\text{max}} = 0.01$. The final difference Fourier map was featureless, with maximum positive and negative peaks of 0.78 and 1.51 $\text{e} \cdot \text{\AA}^{-3}$, respectively.

Computational Details. Nonparametrized Fenske–Hall MO calculations⁶ were carried out on the complex $(\eta^5\text{-C}_5\text{Me}_5)_2\text{-Hf}(\text{S}^n\text{Bu})_2$. This method is based on a self-consistent-field method, which is an approximation of the Hartree–Fock–Roothaan procedure. The molecular geometry and the atomic basis sets used completely determine the resulting eigenvalues and eigenvectors. The calculation was based upon crystallographic data of $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Hf}(\text{S}^n\text{Bu})_2$. The relative positions of the two S atoms and the two centroids of Cp* were adjusted to an idealized microsymmetry of C_s (C_1 for overall symmetry, two Cp* rings being staggered). The idealized atomic coordinates were taken such that the x direction of the master coordinate system originating on the Hf atom bisects the S–Hf–S bond angle, the y direction is normal to the HfS_2 plane, and the z direction is normal to the plane that bisects the S–Hf–S angle. The basis sets used were those provided with the Fenske–Hall program package version 5.1. All calculations were carried out on a VAX 780 computer at The University of Hong Kong.

Results and Discussion

Reaction of $(\eta^5\text{-C}_5\text{Me}_5)_2\text{HfCl}_2$ with 2.2 equiv of LiSR in DME gave $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Hf}(\text{SR})_2$ [$R = n\text{Bu}$ (**1**), C_6H_5 (**2**), $\text{C}_6\text{H}_4\text{OMe-}p$ (**3**), $\text{C}_6\text{H}_4^t\text{Bu-}p$ (**4**)] in reasonable yield. However, unlike the zirconium analogues,³ reaction of $(\eta^5\text{-C}_5\text{Me}_5)_2\text{HfCl}_2$ with NaSR instead of LiSR did not give the desired $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Hf}(\text{SR})_2$ complexes. Instead only the monosubstituted $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Hf}(\text{SR})\text{Cl}$ was obtained. All the newly synthesized complexes gave satisfactory elemental analyses and were characterized by ^1H NMR spectroscopy and EI mass spectrometry. The structures of **1–3** have been determined by X-ray crystallography and represent the first report on the X-ray structure of this class of $(\eta^5\text{-Cp}')_2\text{Hf}(\text{SR})_2$ ($\text{Cp}' = \text{C}_5\text{Me}_5$ or C_5H_5) compounds.

The perspective drawings of complexes **1–3** are depicted in Figures 1–3. The crystal and structure determination data for these complexes are collected in Table 1. Selected bond distances and angles are summarized in Tables 2, 3, and 4, respectively. For all three complexes, the coordination geometries about the Hf atom, defined by the pentamethylcyclopentadienyl ring centroids and the S atoms, are distorted tetrahedral. A projection onto the Hf–S(1)–S(1*) plane shows that the pentamethylcyclopentadienyl rings have a staggered




Figure 1. Perspective drawing of **1** with atomic numbering scheme. Thermal ellipsoids are shown at the 50% probability level. Starred atoms have coordinates at $-x$, y , $0.5-z$.




Figure 2. Perspective drawing of **2** with atomic numbering scheme. Thermal ellipsoids are shown at the 50% probability level. Starred atoms have coordinates at $-x$, y , $0.5-z$.




Figure 3. Perspective drawing of **3** with atomic numbering scheme. Thermal ellipsoids are shown at the 50% probability level. Starred atoms have coordinates at $-x$, y , $0.5-z$.

conformation. The Hf–S distances in **1–3** of 2.4761(9), 2.502(1), and 2.5044(8) $\text{\AA}$ are comparable to the values observed in $(\eta^5\text{-C}_5\text{H}_5)_2\text{HfS}_5$ of 2.512 $\text{\AA}$,⁷ while as

(6) Hall, M. B.; Fenske, R. F. *Inorg. Chem.* **1972**, *11*, 768.

(7) Shaver, A.; McCall, J. M.; Day, V. W.; Vollmer, S. *Can. J. Chem.* **1987**, *65*, 1676.

Table 1. Crystal and Structure Determination Data for Complexes 1-3

	Cp* ₂ Hf(S ⁿ Bu) ₂ (1)	Cp* ₂ Hf(SC ₆ H ₅) ₂ (2)	Cp* ₂ Hf(SC ₆ H ₄ OMe- <i>p</i>) ₂ (3)
mol formula	C ₂₈ H ₄₈ S ₂ Hf	C ₃₂ H ₄₀ S ₂ Hf	C ₃₄ H ₄₄ O ₂ S ₂ Hf
<i>M_r</i>	627.3	667.28	727.33
<i>T</i> , K	301	301	301
<i>a</i> , Å	14.843(3)	14.414(5)	18.034(3)
<i>b</i> , Å	13.233(2)	11.466(3)	11.105(2)
<i>c</i> , Å	16.332(5)	18.190(3)	16.835(3)
β, deg	116.07(2)	103.18(2)	109.69(2)
<i>V</i> , Å ³	2881(1)	2927(1)	3174(1)
cryst color	pale yellow	yellow	yellow green
cryst syst	monoclinic	monoclinic	monoclinic
space group	C2/c (No. 15)	C2/c (No. 15)	C2/c (No. 15)
<i>Z</i>	4	4	4
<i>F</i> (000)	1280	1344	1472
<i>D_c</i> , g cm ⁻³	1.446	1.732	1.522
cryst dims, mm	0.15 × 0.15 × 0.30	0.25 × 0.20 × 0.35	0.25 × 0.15 × 0.30
λ, Å (graphite-monochromated, Mo Kα)	0.710 73	0.710 73	0.710 73
μ(Mo Kα), cm ⁻¹	37.72	37.18	34.40
collection range	2θ _{max} = 48° (<i>h</i> , 0 to 17; <i>k</i> , 0 to 15; <i>l</i> , -18 to 18)	2θ _{max} = 50° (<i>h</i> , 0 to 17; <i>k</i> , 0 to 13; <i>l</i> , -21 to 21)	2θ _{max} = 51° (<i>h</i> , 0 to 21; <i>k</i> , 0 to 13; <i>l</i> , -20 to 20)
scan mode and scan speed, deg min ⁻¹	ω-2θ; 1.50-8.24	ω-2θ; 1.18-5.49	oscillation
scan width, deg	0.80 + 0.35 tan θ	0.66 + 0.35 tan θ	
no. of data collected	2479	2831	13 424
no of unique data	2416	2732	3073
no. of data used in refinement, <i>m</i>	2167	2484	2863
no. of parameters refined, <i>p</i>	141	159	177
<i>R</i> ^a	0.019	0.024	0.025
<i>wR</i> ^a	0.025	0.038	0.044
goodness-of-fit, <i>S</i>	1.58	2.17	1.92
maximum shift, (Δ/ <i>σ</i>) _{max}	0.01	0.02	0.01
residual maximum in final difference map, e Å ⁻³	+0.51, -0.55	+0.61, -1.18	+0.78, -1.51

^a *w* = 4*F*_o²/σ²(*F*_o)², where σ²(*F*_o)² = [σ²(*I*) + (0.020*F*_o²)] with *I* > 3σ(*I*) for **1**; *w* = 4*F*_o²/σ²(*F*_o)², where σ²(*F*_o)² = [σ²(*I*) + (0.028*F*_o²)] with *I* > 3σ(*I*) for **2**; *w* = 4*F*_o²/σ²(*F*_o)², where σ²(*F*_o)² = [σ²(*I*) + (0.032*F*_o²)] with *I* > 3σ(*I*) for **3**.

Table 2. Selected Bond Distances (Å) and Bond Angles (deg) for 1

Hf(1)–S(1)	2.4761(9)	Hf(1)–C(1)	2.546(3)
Hf(1)–C(2)	2.529(3)	Hf(1)–C(3)	2.577(3)
Hf(1)–C(4)	2.596(4)	Hf(1)–C(5)	2.555(4)
Hf–Cp* _{centroid}	2.269	S(1)–C(11)	1.828(4)
C(1)–C(2)	1.410(5)	C(2)–C(3)	1.415(5)
C(3)–C(4)	1.410(5)	C(4)–C(5)	1.430(5)
C(1)–C(5)	1.402(5)	C(1)–C(6)	1.507(5)
C(2)–C(7)	1.506(5)	C(3)–C(8)	1.502(5)
C(4)–C(9)	1.500(5)	C(5)–C(10)	1.503(5)
C(11)–C(12)	1.477(6)	C(12)–C(13)	1.536(7)
C(13)–C(14)	1.447(8)		
S(1)–Hf(1)–S(1*)	95.90(6)	Hf(1)–S(1)–C(11)	111.6(2)
S(1)–C(11)–C(12)	112.5(3)	C(11)–C(12)–C(13)	111.8(4)
C(12)–C(13)–C(14)	113.2(5)	Cp* _{centroid} –Hf–Cp* _{centroid}	135.1

a result of lanthanide contraction effect, they are slightly shorter than the Zr–S distances observed in the related zirconocene thiolate complexes (η⁵-C₅Me₅)₂Zr(SR)₂ [R = ⁿBu (2.4987(8) Å);³ R = C₆H₅ (2.522(1) Å)⁸]. The S(1)–Hf(1)–S(1*) angles of 95.90(6)°, 100.52(5)°, and 100.62(4)° in **1–3** are comparable to the corresponding S(1)–Zr(1)–S(1*) angles in (η⁵-C₅Me₅)₂Zr(SR)₂ [R = ⁿBu (96.14)°;³ R = C₆H₅ (101)°⁸] and are consistent with the difference in the steric demand of the –Sⁿ-Bu and –SPh moieties. The Hf–Cp*_{centroid} distances in **1–3** of 2.269, 2.261, and 2.265 Å are slightly shorter than the Zr–Cp*_{centroid} distances of 2.27 Å in (η⁵-C₅Me₅)₂Zr(SⁿBu)₂.³ The Cp*_{centroid}–M–Cp*_{centroid} angles (M = Zr,³ Hf) are also similar and are consistent with the steric demand of the ligands. Similar to the zirconium analogues,^{3,8} complexes **1–3** exhibit an *endo* conforma-

Table 3. Selected Bond Distances (Å) and Bond Angles (deg) for 2

Hf(1)–S(1)	2.502(1)	Hf(1)–C(1)	2.545(4)
Hf(1)–C(2)	2.504(4)	Hf(1)–C(3)	2.566(5)
Hf(1)–C(4)	2.595(4)	Hf(1)–C(5)	2.560(4)
Hf–Cp* _{centroid}	2.261	S(1)–C(11)	1.778(4)
C(1)–C(2)	1.425(7)	C(2)–C(3)	1.436(6)
C(3)–C(4)	1.406(7)	C(4)–C(5)	1.442(6)
C(1)–C(5)	1.416(6)	C(1)–C(6)	1.507(6)
C(2)–C(7)	1.498(7)	C(3)–C(8)	1.510(7)
C(4)–C(9)	1.506(6)	C(5)–C(10)	1.480(5)
C(11)–C(12)	1.390(8)	C(12)–C(13)	1.367(7)
C(13)–C(14)	1.412(10)	C(14)–C(15)	1.340(1)
C(15)–C(16)	1.429(9)		
S(1)–Hf(1)–S(1*)	100.52(5)	Hf(1)–S(1)–C(11)	119.0(1)
S(1)–C(11)–C(12)	123.4(3)	S(1)–C(11)–C(16)	116.6(4)
Cp* _{centroid} –Hf–Cp* _{centroid}	135.54		

tion as reflected by their S(1*)–Hf–S(1)–C(11) torsion angles of 48.17°, 48.79°, and 52.07°, respectively. This is in accord with theoretical calculations on this class of compounds and is favored by d⁰ metal centers in order to maximize the pπ–dπ overlap between sulfur and the metal center.⁹ The larger deviation from 90° of the S–Hf–S–C torsion angle in **1–3** is in line with the steric requirement of the pentamethylcyclopentadienyl units, which direct the alkyl/phenyl substituents further away from the more sterically demanding pentamethylcyclopentadienyl groups.

The luminescent behavior of complexes **1–4** has been studied. Their photophysical data together with those of the precursor complex (η⁵-C₅Me₅)₂HfCl₂ and the

(9) (a) Lauher, J. W.; Hoffmann, R. *J. Am. Chem. Soc.* **1976**, *98*, 1729. (b) Calhorda, M. J.; Carrondo, M. A. A. F. d. C. T.; Dias, A. R.; Frazao, C. F.; Hursthouse, M. B.; Simoes, J. A. M.; Teixeira, C. *Inorg. Chem.* **1988**, *27*, 2513.


(8) Howard, W. A.; Trnka, T. M.; Parkin, G. *Inorg. Chem.* **1995**, *34*, 5900.

Table 4. Selected Bond Distances (Å) and Bond Angles (deg) for 3


Hf(1)–S(1)	2.5044(8)	Hf(1)–C(1)	2.549(4)
Hf(1)–C(2)	2.508(4)	Hf(1)–C(3)	2.576(4)
Hf(1)–C(4)	2.598(4)	Hf(1)–C(5)	2.548(3)
Hf–Cp* _{centroid}	2.265	S(1)–C(11)	1.787(3)
C(1)–C(2)	1.418(6)	C(2)–C(3)	1.421(6)
C(3)–C(4)	1.427(6)	C(4)–C(5)	1.423(6)
C(1)–C(5)	1.421(6)	C(1)–C(6)	1.501(5)
C(2)–C(7)	1.496(6)	C(3)–C(8)	1.495(6)
C(4)–C(9)	1.492(5)	C(5)–C(10)	1.510(4)
C(11)–C(12)	1.388(5)	C(12)–C(13)	1.396(6)
C(13)–C(14)	1.387(6)	C(14)–C(15)	1.376(5)
C(15)–C(16)	1.392(6)	O(1)–C(14)	1.376(5)
O(1)–C(17)	1.419(5)		
S(1)–Hf(1)–S(1*)	100.62(4)	Hf(1)–S(1)–C(11)	117.4(1)
S(1)–C(11)–C(12)	118.3(3)	S(1)–C(11)–C(16)	122.7(4)
Cp* _{centroid} –Hf–Cp* _{centroid}	134.5		

Table 5. Photophysical Data for Hafnium Thiolate Complexes

compound	medium (T/K)	emission (λ _{em} /nm)	lifetime (τ ₀ /μs)
1	PE/Tol (v/v 10:1) glass (77)	542	
	solid (77)	538	
2	Tol (298)	540	<0.1
	PE/Tol (v/v 10:1) glass (77)	526	
3	solid (77)	522	
	solid (298)	513	1.5 ± 0.1
4	PE/Tol (v/v 10:1) glass (77)	528	
	solid (77)	532	
	solid (298)	513	9.0 ± 0.5
(η ⁵ -C ₅ Me ₅) ₂ HfCl ₂	solid (77)	455	<0.1
(η ⁵ -C ₅ H ₅) ₂ HfCl ₂	solid (77)	427	
(η ⁵ -C ₅ Me ₅) ₂ ZrCl ₂ ^a	solid (77)	494	
(η ⁵ -C ₅ H ₅) ₂ ZrCl ₂ ^a	solid (77)	452	

^a From ref 3.**Figure 4.** Excitation spectrum (···) and emission spectrum (—) of **2** in petroleum ether/toluene (v/v 10:1) glass at 77 K.

related complexes (η⁵-C₅H₅)₂HfCl₂, (η⁵-C₅Me₅)₂ZrCl₂, and (η⁵-C₅H₅)₂ZrCl₂ are collected in Table 5. The excitation and emission spectra of **2** in petroleum ether/Et₂O glass at 77 K are shown in Figure 4. The excitation spectrum closely resembles that of the electronic absorption spectrum (Figure 5). The complexes (η⁵-C₅Me₅)₂HfCl₂, (η⁵-C₅H₅)₂HfCl₂, (η⁵-C₅Me₅)₂ZrCl₂, and (η⁵-C₅H₅)₂ZrCl₂ emit at 455, 427, 494, and 452 nm, respectively, in the solid state at 77 K. A comparison

**Figure 5.** Electronic absorption spectrum of **2** in toluene at room temperature.**Table 6. Idealized Bond Lengths (Å) and Angles (deg) of 1 Used for Nonparametrized Fenske–Hall MO Calculations**

Hf–Cp* _{centroid}	2.269
Hf–S	2.48
S–C	1.83
Cp* _{centroid} –Hf–Cp* _{centroid}	135.1
S–Hf–S	95.90
Hf–S–C ₁₁	111.6
C ₁₁ –S–Hf–S*	48.17
C ₁₂ –C ₁₁ –S*–Hf	44.82

of their emission spectra shows that the emission energy follows the order Hf > Zr and C₅H₅ > C₅Me₅. With reference to previous spectroscopic work on the related titanocene derivatives¹⁰ and the observed emission trends, the emissions observed in (η⁵-C₅Me₅)₂MCl₂ are tentatively assigned as derived from the respective C₅Me₅ → M ligand-to-metal charge-transfer (LMCT) states. Complexes **1–4** exhibit intense emission at ca. 520–540 nm upon visible light excitation in the solid state at 77 K. The much lower emission energies in **1–4** relative to (η⁵-C₅Me₅)₂HfCl₂ may suggest an emission of different origin. An assignment of the emission origin in complexes **1–4** as derived from the C₅Me₅ → Hf LMCT transition is not favored since the thiolate ligands, being a better σ-donor than the electronegative chloro group, would be expected to render the Hf center less electron deficient and, hence, a poorer electron acceptor. This would cause the C₅Me₅ → Hf LMCT transition to shift to higher energies instead. The observation of an opposite trend would disfavor such an assignment. A comparison of the emission energies of the complexes in the solid state at 77 K shows that **1** < **4** < **3** < **2**, which is in line with the σ-donating effect of the thiolate ligands SⁿBu > SC₆H₄ⁱBu-*p* > SC₆H₄OMe-*p* > SC₆H₅. A similar trend has also been found in the related zirconium chalcogenolate complexes.³ The observation of such an emission trend is suggestive of either a thiolate to hafnium (SR → Hf) LMCT origin or a thiolate to pentamethylcyclopentadienyl (SR → C₅Me₅) LLCT origin. Although one cannot exclude the possibility of a LLCT state, we prefer its assignment as an origin with predominant LMCT character since the emission energies are found to be sensitive to the nature of the metal center. A comparison of the emission

(10) See for example: (a) Kenney, J. W., III; Boone, D. R.; Striplin, D. R.; Chen, Y. H.; Hamar, K. B. *Organometallics* **1993**, *12*, 3671. (b) Bruce, M. R. M.; Sclafani, A.; Tyler, D. R. *Inorg. Chem.* **1986**, *25*, 2546. (c) Bruce, M. R. M.; Kenter, A.; Tyler, D. R. *J. Am. Chem. Soc.* **1984**, *106*, 639.

Table 7. Energies and Percent Compositions for the Frontier Orbitals of $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Hf}(\text{S}^n\text{Bu})_2$

molecular orbital	energy (eV)	% composition		
		SR	Hf	C ₅ Me ₅
HOMO(88)	-9.48	77.4	0.8	21.8
LUMO(89)	-2.65	16.4	73.2	10.4

spectra of $(\eta^5\text{-C}_5\text{Me}_5)_2\text{M}(\text{SR})_2$ shows that, for the same thiolate ligand, the emission for the hafnium complexes occurs at a higher energy than that of the zirconium counterparts. For example, the 77 K solid-state emission of $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Hf}(\text{S}^n\text{Bu})_2$ occurs at 538 nm, while that of $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Zr}(\text{S}^n\text{Bu})_2$ occurs at 610 nm.³

The electronic structures of this class of compounds have been probed by Fenske–Hall molecular orbital calculation on complex **1**. Table 6 summarizes the idealized bond lengths and angles used for the MO calculations. Table 7 shows the percent composition for the frontier molecular orbitals of complex **1**. The calculation results show that the composition of the

HOMO is approximately 77% SⁿBu and 22% C₅Me₅ in character, while that of the LUMO is approximately 73% Hf and 10% C₅Me₅. The large percent composition of thiolate character in the HOMO has also been found in the analogous $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Zr}(\text{S}^n\text{Bu})_2$ ³ and in the related $(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\text{SH})_2$.¹¹ Based on the results of the calculations, an assignment of the origin of emission as derived from a predominant thiolate-to-hafnium (RS → Hf) LMCT state mixed with some C₅Me₅ → Hf LMCT character is favored. A similar assignment has been made for the related zirconium complexes.³

Acknowledgment. V.W.W.Y. acknowledges financial support from the Research Grants Council and The University of Hong Kong.

Supporting Information Available: Tables giving fractional coordinates and thermal parameters, general displacement parameter expressions (*U*), and all bond distances and bond angles for **1–3** (28 pages). Ordering information is given on any current masthead page.

OM980602F

(11) Petersen, J. L.; Lichtenberger, D. L.; Fenske, R. F.; Dahl, L. F. *J. Am. Chem. Soc.* **1975**, *97*, 6433.