

does the desired "CpCo" transfer occur producing the new dinuclear complex $(\eta^5\text{-C}_5\text{H}_5)\text{Ir}(\mu\text{-CO})_2\text{Co}(\eta^5\text{-C}_5\text{H}_5)$ in addition to 1 but also the homodinuclear iridium analogue of composition $(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}(\mu\text{-CO})_2\text{Ir}(\eta^5\text{-C}_5\text{H}_5)$. Attempts to run this reaction on a preparative scale have, however, been thwarted by the extreme reactivity of compounds of type $(\eta^5\text{-C}_5\text{H}_5)\text{M}(\mu\text{-CO})_2\text{M}'(\eta^5\text{-C}_5\text{H}_5)$ ($\text{M}/\text{M}' = \text{Co}/\text{Co}, \text{Co}/\text{Ir}, \text{Ir}/\text{Ir}$).^{9b,31} Work in this and related areas is ongoing and will be reported in the near future.

Summary and Conclusions

A high-yield synthesis for the title complex $(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}(\mu\text{-CO})_2[\text{Co}(\eta^5\text{-C}_5\text{H}_5)]_2$ (3) is described and a detailed discussion of its properties in solution and in the solid state presented. The solid-state structure of 3 is derived from the unsaturated heterodinuclear precursor complex $(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}(\mu\text{-CO})_2\text{Co}(\eta^5\text{-C}_5\text{H}_5)$ (1), to which a $\text{Co}(\eta^5\text{-C}_5\text{H}_5)$ fragment has been added. This observation fits neatly into the growing number of examples illustrating the isolobal concepts developed by Hoffmann and co-workers.¹⁷ The unique CoCp fragment interacts strongly with the metal atoms of the dinuclear fragment and weakly with the two edge-bridging carbonyl ligands in the solid state. In solution, a low-energy CO scrambling process equilibrates the two CoCp fragments. 3 is paramagnetic in solution and in the solid state.

(31) Reaction of $\text{CpIr}(\text{CO})_2$ with $\text{CpCo}(\text{C}_2\text{H}_4)_2$ has, under all conditions investigated thus far, yielded only a complex mixture of tri- and tetranuclear clusters.

Trinuclear 3 is extremely labile by virtue of its formal 46 valence electron count and the reactivity of one of the CoCp fragments. Loss of this fragment is easily initiated by briefly exposing the complex to oxygen, ethylene, or mononuclear cyclopentadienyl dicarbonyl complexes. In all such cases the heterodinuclear precursor complex 1 is also produced initially. 3 readily adds carbon monoxide and catalyzes the decomposition of diazoalkanes to give saturated 48 valence electron cluster complexes, the latter case having a new μ -methylene group on the trinuclear frame. Thus 3 would appear to be a valuable synthetic precursor for a number of studies. On the one hand, small organic fragments are quite easily incorporated into 3 without disruption of the cluster backbone. Under other circumstances 3 appears to cleanly transfer a CoCp fragment to a variety of interesting substrates. These and other studies of the reactivity of 3 and similar analogues are in progress and will be reported in the near future.

Acknowledgment. This work was generously supported by the Alexander von Humboldt-Stiftung, the NATO, Degussa AG, Hoechst AG, Quandt-Stiftung der Varta AG, and the Deutsche Forschungsgemeinschaft.

Registry No. 1, 89371-07-3; 2, 69393-67-5; 3, 92957-47-6; 4, 92957-48-7; 5, 92957-49-8; $(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}(\text{CO})(\pi\text{-C}_2\text{H}_4)$, 92957-50-1; $\text{CpCo}(\text{CO})(\text{C}_2\text{H}_4)$, 77674-12-5; $(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}(\text{CO})_2$, 32660-96-1; $\text{CpIr}(\text{CO})_2$, 12192-96-0.

Supplementary Material Available: Table of observed and calculated structure factors and complete table (26 pages). Ordering information is given on any current masthead page.

Synthesis and Reactions of Thermally Stable Platinum(II) Carboxylic Acids (Hydroxycarbonyls) Containing Bis(tertiary phosphines)[†]

Martin A. Bennett* and Andrzej Rokicki

Research School of Chemistry, Australian National University, Canberra, A.C.T. 2601, Australia

Received August 23, 1984

Platinum(II) carboxylic acids (or hydroxycarbonyls) of general formula $\text{Pt}(\text{CO}_2\text{H})(\text{C}_6\text{H}_9)(\text{P-P})$ (C_6H_9 = 1-cyclohexenyl; P-P = $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2$ (1), *cis*- $\text{Ph}_2\text{PCH}=\text{CHPPh}_2$ (2), $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{CH}_2\text{PPh}_2$ (3), and $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{PPh}_2$ (4)) have been prepared by the successive reaction of water and carbon monoxide with the cyclohexyne complexes $\text{Pt}(\text{C}_6\text{H}_9)(\text{P-P})$. Complex 3 readily inserts a second molecule of CO into the platinum(II)-cyclohexenyl bond to give an acylplatinum(II) carboxylic acid, $\text{Pt}(\text{CO}_2\text{H})(\text{COC}_6\text{H}_9)(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{CH}_2\text{PPh}_2)$ (5). The complexes, which have been characterized by elemental analysis, IR spectroscopy and ^1H , ^{13}C , and ^{31}P NMR spectroscopy, are monomeric in dichloromethane but appear to be associated through hydrogen bonds in the solid state. Complexes 1-5 do not decarboxylate in the solid state up to ca. 80 °C and do not form carboxylate ions by loss of a proton. These properties probably arise from the presence of nondissociating, strongly electron-donating ligands in the coordination sphere. The platinum(II) carboxylic acids do readily undergo esterification with alcohols, and 3 reacts with dimethylamine or *tert*-butylamine to give carbamoyl complexes $\text{Pt}(\text{CONR}^1\text{R}^2)(\text{C}_6\text{H}_9)(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{CH}_2\text{PPh}_2)$ ($\text{R}^1 = \text{R}^2 = \text{Me}$, 6; $\text{R}^1 = t\text{-Bu}$, $\text{R}^2 = \text{H}$, 7). These reactions, and the ready exchange of ^{13}CO with the $\text{Pt-CO}_2\text{H}$ group, indicate that the carboxylic acids behave as $[\text{Pt}(\text{C}_6\text{H}_9)(\text{CO})(\text{P-P})]\text{OH}$, although the carbonyl cations could not be detected in solutions of the acids. Treatment of 5 with HBF_4 gives $[\text{Pt}(\text{COC}_6\text{H}_9)(\text{CO})(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{CH}_2\text{PPh}_2)]\text{BF}_4$, as expected, whereas 3 undergoes competitive cleavage of the platinum-(II)-cyclohexenyl bond and consequent β -elimination of CO_2 from the $\text{Pt-CO}_2\text{H}$ group.

Introduction

Metalloxy-carboxylic acids (hydroxycarbonyl metal complexes), $\text{L}_n\text{MCO}_2\text{H}$, have been proposed as key intermediates in several important reactions involving carbon

monoxide. They include oxidation of CO (either free or coordinated) by metal ions,¹ the isotopic exchange of metal carbonyls with ^{18}O -labeled water (first observed by Muettterties),² the homogeneously catalyzed water gas shift

[†] Dedicated to Professor Earl Muettterties, whose untimely death has deprived the inorganic and organometallic chemistry community of one of its most original and perceptive contributors.

(1) (a) Harkness, A. C.; Halpern, J. *J. Am. Chem. Soc.* 1961, 83, 1258. (b) Nakamura, S.; Halpern, J. *Ibid.* 1961, 83, 4102. (c) Bercaw, J. E.; Goh, L. Y.; Halpern, J. *Ibid.* 1972, 94, 6534.

Table I. Carbon-13 NMR Data for Platinum(II) Carboxylic Acids $\text{Pt}(\text{CO}_2\text{H})(\text{C}_6\text{H}_9)(\text{P}-\text{P})$ and Derived Carbamoyl Complexes^{a, b}

complex	solv	$\delta_{\text{CO}_2\text{H}/\text{CONR}_2}$	$^1J_{\text{PtC}}$	$^2J_{\text{CP}^1}$	$^2J_{\text{CP}^2}$
1	CD_2Cl_2	196.2	1320	157.3	8.5
2	CD_2Cl_2	195.0	1330	160.7	8.6
	$(\text{CD}_3)_2\text{SO}$	198.6	1291	159.0	8.6
3	CD_2Cl_2	196.4	1297	153.3	11.0
	$(\text{CD}_3)_2\text{SO}$	197.9	1274	152.2	10.3
4	CD_2Cl_2	194.7	1299	155.6	12.0
	$(\text{CD}_3)_2\text{SO}$	196.5	1270	153.9	12.0
5 ^c	CD_2Cl_2	193.1	1265	132.5	12.2
6	CDCl_3	199.3	1185	140.0	11.0
	CD_2Cl_2	198.4	1176	141.0	ca. 11
7	CD_2Cl_2	195.5	1116	138.0	ca. 10

^a Recorded on samples enriched to 90% with ^{13}C , chemical shifts being in parts per million to high frequency of Me_4Si ; coupling constants in hertz. ^b Phosphorus atoms are labeled as in eq 1, 3, and 4 (see text). ^c $\delta_{\text{COC}_6\text{H}_5}$, 244.7; $^1J_{\text{PtC}}$ = 895 Hz, $^2J_{\text{CP}^1}$ = 8.7 Hz, and $^2J_{\text{CP}^2}$ = 105 Hz.

reaction,³ and the reaction of metal carbonyls with base to give metal hydrides.^{3,4} However, relatively few metal-carboxylic acids have been isolated and characterized and little is known about their chemistry. Fairly stable compounds of this class include $\text{IrCl}_2(\text{CO}_2\text{H})(\text{CO})(\text{PMe}_2\text{Ph})_2$,⁵ $(\eta\text{-C}_5\text{H}_5)\text{Fe}(\text{CO}_2\text{H})(\text{CO})(\text{PPh}_3)_3$,⁶ $(\eta\text{-C}_5\text{H}_5)\text{Re}(\text{CO}_2\text{H})(\text{CO})(\text{NO})$,⁷ $(\eta\text{-C}_5\text{H}_5)\text{Re}(\text{CO}_2\text{H})(\text{CO})(\text{N}_2\text{Ar})$,⁸ and $(\eta\text{-C}_5\text{H}_5)\text{Mo}(\text{CO}_2\text{H})(\text{CO})_2(\text{PPh}_3)_3$.⁹ A communication from this laboratory which appeared in 1973¹⁰ reported briefly the preparation of the first platinum(II) carboxylic acids $\text{Pt}(\text{CO}_2\text{H})\text{R}(\text{dppe})$ ($\text{R} = \text{CH}_3$, C_6H_9 ; $\text{C}_6\text{H}_9 = 1\text{-cyclohexenyl}$) by reaction of CO with the corresponding hydroxo complexes and noted their remarkable thermal stability, both in the solid state and in solution. Subsequently, analogous compounds $\text{Pt}(\text{CO}_2\text{H})\text{R}(\text{vdpp})$ ($\text{R} = \text{CH}_2\text{CN}$, CF_3 ; $\text{vdpp} = \text{cis-Ph}_2\text{PCH}=\text{CHPPh}_2$) have been reported,¹¹ and $\text{trans-PtCl}(\text{CO}_2\text{H})(\text{PET}_3)_2$ has been isolated from the reaction of $\text{trans-[PtCl(CO)(PET}_3)_2]^+$ with water.¹² We report here in detail our work on platinum(II) carboxylic acids of the type $\text{Pt}(\text{CO}_2\text{H})\text{R}(\text{P}-\text{P})$ that contain bis(tertiary phosphines).¹³ We have concentrated on complexes containing the 1-cyclohexenyl group ($\text{R} = \text{C}_6\text{H}_9$) because they are generally more stable and more soluble in organic solvents than the corresponding methyl compounds.

Experimental Section

Starting Materials. The following were obtained from the

(2) (a) Muetterties, E. L. *Inorg. Chem.* **1965**, *4*, 1841. (b) Darensbourg, D. J.; Froelich, J. A. *J. Am. Chem. Soc.* **1977**, *99*, 4726. (c) Darensbourg, D. J.; Froelich, J. A. *Inorg. Chem.* **1977**, *17*, 3300. (d) Darensbourg, D. J.; Darensbourg, M. Y.; Walker, N.; Froelich, J. A.; Barros, H. L. C. *Ibid.* **1979**, *18*, 1401.

(3) (a) Ford, P. C. *Acc. Chem. Res.* **1981**, *14*, 31 and references cited therein. (b) Darensbourg, D. J.; Rokicki, A. *Organometallics* **1982**, *1*, 1685.

(4) (a) Kruck, T.; Noack, M. *Chem. Ber.* **1964**, *97*, 1693. (b) Clark, H. C.; Jacobs, W. J. *Inorg. Chem.* **1970**, *9*, 1229. (c) Pearson, R. G.; Mauermann, H. J. *Am. Chem. Soc.* **1982**, *104*, 500.

(5) Deeming, A. J.; Shaw, B. L. *J. Chem. Soc. A* **1969**, 443.

(6) Grice, N.; Kao, S. C.; Pettit, R. *J. Am. Chem. Soc.* **1979**, *101*, 1627.

(7) (a) Casey, C. P.; Andrews, M. A.; Rinz, J. E. *J. Am. Chem. Soc.* **1979**, *101*, 741. (b) Tam, W.; Lin, G.-Y.; Wong, W.-K.; Kiel, W. A.; Wong, V. K.; Gladysz, J. A. *Ibid.* **1982**, *104*, 141. (c) Sweet, J. R.; Graham, W. A. *Organometallics* **1982**, *1*, 982.

(8) Barrientos-Penna, C. F.; Gilchrist, A. B.; Sutton, D. *Organometallics* **1983**, *2*, 1265.

(9) Gibson, D. H.; Owens, K.; Ong, T.-S. *J. Am. Chem. Soc.* **1984**, *106*, 1125.

(10) Appleton, T. G.; Bennett, M. A. *J. Organomet. Chem.* **1973**, *55*, C89.

(11) Michelin, R. A.; Napoli, M.; Ros, R. *J. Organomet. Chem.* **1979**, *175*, 239.

(12) Catellani, M.; Halpern, J. *Inorg. Chem.* **1980**, *19*, 566.

(13) Abbreviations: dppe = 1,2-bis(diphenylphosphino)ethane, $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2$; vdpp = *cis*-vinylene-1,2-bis(diphenylphosphino), *cis-Ph}_2\text{PCH}=\text{CHPPh}_2; dppp = 1,3-bis(diphenylphosphino)propane, $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{CH}_2\text{PPh}_2$; dppb = 1,4-bis(diphenylphosphino)butane, $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{PPh}_2$.*

Table II. Carbon-13 NMR Data for $\text{Pt}(\text{CO}_2\text{H})(\text{C}_6\text{H}_9)(\text{dppp})$ (3) in Various Solvents^a

solv	$\delta_{\text{CO}_2\text{H}}$	$^1J_{\text{PtC}}$	$^2J_{\text{CP}^1}$	$^2J_{\text{CP}^2}$
C_6D_6	196.0	1306	155.3	11.0
C_7D_8	195.5	1311	154.0	11.2
CDCl_3	196.9	1309	153.9	11.7
CD_2Cl_2	196.4	1297	153.3	11.0
$\text{THF}-d_3$	196.4	1298	155.0	12.0
$(\text{CD}_3)_2\text{SO}$	197.9	1274	152.2	10.3

^a Coupling constants in hertz.

commercial suppliers given in parentheses: CO (Matheson), CO enriched to 90% in ^{13}C (Isotope Labeling Corp., Whippany, NJ), 40% aqueous HBF_4 (Ajax Chemicals), $\text{HBF}_4\cdot\text{Et}_2\text{O}$ (Aldrich), dimethylamine (Tokyo Kasei), *tert*-butylamine (EGA Chemie), and the bis(tertiary phosphines) dppe, vdpp, and dppp (Strem). 1,4-Bis(diphenylphosphino)butane (dppb)¹⁴ and the cyclohexyne complex $\text{Pt}(\text{C}_6\text{H}_9)(\text{dppe})$ ¹⁵ were prepared by literature methods. The other cyclohexyne complexes $\text{Pt}(\text{C}_6\text{H}_9)(\text{P}-\text{P})$ were prepared similarly to $\text{Pt}(\text{C}_6\text{H}_9)(\text{dppe})$.¹⁶

Spectroscopic Measurements. The following instruments were used for NMR measurements with an internal deuterium lock: JEOL FX 200 and Bruker CXP 200 (^1H and ^{13}C at 50.10- and 50.29 MHz, respectively) and Bruker CXP 200 (^{31}P at 80.98 MHz). Routine IR spectra were measured on a Perkin-Elmer 683 instrument. High-resolution IR spectra on dichloromethane solutions were recorded on a Perkin-Elmer 225 spectrometer with use of 0.1 mm matched, sealed KBr cells. The spectra were calibrated against the spectra of ammonia in the range 3491.4-3393.9 cm^{-1} and of water vapor in the range 1706.3-1417.5 cm^{-1} . NMR spectroscopic data for complexes 1-7 are in Tables I-III, and IR data for the carboxylic acids 1-5 are in Table IV.

Analyses were performed in the microanalytical laboratories of this school (Miss Brenda Stevenson and associates).

Preparations. (1) **(1,3-Bis(diphenylphosphino)propane)(1-cyclohexenyl)(hydroxycarbonyl)platinum(II), $\text{Pt}(\text{CO}_2\text{H})(\text{C}_6\text{H}_9)(\text{dppp})$ (3).** A solution of the cyclohexyne complex $\text{Pt}(\text{C}_6\text{H}_9)(\text{dppp})$ (200 mg, 0.290 mmol) in THF (20 mL) was treated with water (10 mL). The clear, colorless solution was stirred at room temperature, the course of reaction being monitored by observing the disappearance of the $\nu(\text{C}\equiv\text{C})$ band at ca. 1705 cm^{-1} ; this usually required ca. 20 h. The solution was then evaporated to dryness under reduced pressure to give a white solid, $\text{Pt}(\text{OH})(\text{C}_6\text{H}_9)(\text{dppp})$. This was redissolved in THF (2 mL), and the straw-yellow solution was transferred into a test tube that was fitted with a septum cap secured by insulation tape and that contained a stirring bar. The solution was stirred under a pressure of carbon monoxide slightly above atmospheric for 10 min at room temperature. After the gas had been vented, the solution was treated with *n*-hexane to give a colorless solid. This was filtered off, washed with hexane, and dried in vacuo to give 190 mg (0.243 mmol, 84%) of 3, mp 112-115 °C dec.

(14) Sacconi, L.; Gelsomini, J. *Inorg. Chem.* **1968**, *7*, 291.

(15) Bennett, M. A.; Yoshida, T. *J. Am. Chem. Soc.* **1978**, *100*, 1750.

(16) Bennett, M. A.; Rokicki, A., to be submitted for publication.

Table III. Phosphorus-31 and Proton NMR Data for Platinum(II) Carboxylic Acids Pt(CO₂H)(C₆H₅)(P-P) and Derived Carbamoyl Complexes^{a, b}

	δ_{P^1} (¹ J _{PtP¹)}	δ_{P^2} (¹ J _{PtP²)}	² J _{PP}	$\delta_{=CH of C_6H_5}$ (³ J _{PtH} , ⁴ J _{PH})	δ_{CO_2H}
1	39.7 (1897)	41.0 (1645)		5.25 (76, 9)	9.79
2	nm	nm		5.14 (75, 9)	9.82
3	-1.2 (1874) ^c	-6.0 (1610) ^d	22.6	5.05 (62, 7)	9.33
4	16.8 (1950)	6.6 (1666)	15.6	5.08 (76, 7)	9.26
5	-3.3 (1967)	-8.1 (1495)	26.3	ca. 7.15	9.37
6	-0.2 (1698) ^e	-7.2 (1610) ^f	ca. 20	4.90 (75, 8) ^{g, h}	
7	-1.2 (1519) ⁱ	-5.4 (1700) ^j	ca. 19	4.78 (73, 7) ^{e, k}	

^a $\delta(P)$ in ppm relative to external 85% H₃PO₄ (positive to high frequency) and coupling constants in hertz, measured in CD₂Cl₂; ¹H NMR spectra measured in (CD₃)₂SO, except where indicated; P¹ trans to CO₂H or CONR₂ and P² trans to C₆H₅ or C₆H₄CO. ^b All complexes showed complex multiplets in the range δ 1-3 due to the CH₂ protons of the C₆H₅ group and of the bis(tertiary phosphines) and in the range δ 7-8 due to the aromatic protons. ^c ²J_{CP¹} = 153 Hz. ^d ²J_{CP²} = 11 Hz. ^e ²J_{CP¹} = 140 Hz. ^f ²J_{CP²} = 11 Hz. ^g Measured in CDCl₃. ^h δ 2.28 (s, NMe), 2.59 (1:4:1t, NMe, ⁴J_{PtH} = 5.3 Hz). ⁱ ²J_{CP¹} = 138 Hz. ^j ²J_{CP²} = 9.6 Hz. ^k δ 5.31 (s, NH), 0.93 (s, *t*-Bu).

Table IV. IR Data (cm⁻¹) for Pt(CO₂H)(C₆H₅)(P-P)^{a, b}

complex	labeling of COOH	$\nu(OH)$	$\nu(C=O)$	$\nu(C-O)$
1	¹² COOH	2670 w br	1630 w br	1215 m, 1160 m
	¹³ COOH	2660 w br	1610 m br	1185 m, 1160 m
2	¹² COOH	2670 w br (3441 w)	1651 m, 1597 m (1652 s)	^c
	¹³ COOH	2660 w br	1610 m, 1555 m	^c
3	¹² COOH	2680 w (3441 w)	1645 msh, 1605 s, (1648 s)	1153 m
		2650 w br (3435 w)	1600 msh, 1565 s (1615 s)	1127 m
	¹² CO ¹⁸ OH	2670 w br (3428 w)	1635 msh, 1600 s (1647 m, 1632 m)	^c
4	¹² COOH	2680 w (3431 w)	1645 msh, 1595 s (1652 s)	1145 m
	¹³ COOH	2660 w br	1610 msh, 1560 s	^c
5	¹² COOH	2690 w (3436 w)	1640 msh, 1610 s ^d (1656 s) ^e	1120 m
	¹³ COOH	2665 w br	1625 msh, 1571 s ^f	1153 m, 1137 m (1156 m)
				1135 m, 1115 m

^a Measured in KBr disks; values in parentheses refer to CH₂Cl₂ solution. ^b Abbreviations: br, broad; m, medium; s, strong; sh, shoulder; w, weak. ^c Band due to $\nu(C-O)$ could not be located unequivocally owing to overlap with bands due to the bis(tertiary phosphine), either in KBr disk or in CH₂Cl₂ solution. ^d $\nu(C=O)$ for ¹²COC₆H₅: 1563 s cm⁻¹ (KBr). ^e $\nu(C=O)$ for ¹²COC₆H₅: 1565 cm⁻¹ (CH₂Cl₂). ^f $\nu(C=O)$ for ¹³COC₆H₅: 1532 s cm⁻¹ (KBr).

Anal. Calcd for C₃₄H₃₆O₂P₂Pt·0.7THF: C, 56.33; H, 5.33; P, 7.92; mol wt, 734. Found: C, 56.47; H, 5.26; P, 7.95; mol wt by osmometry, 761 (CH₂Cl₂), 735 (pyridine).

The complexes Pt(CO₂H)(C₆H₅)(P-P) (P-P = dppe (1), vdpp (2), and dppb (4)) were prepared similarly from the appropriate cyclohexyne complexes in yields of 40%, 27%, and 84% respectively. In the case of 1 and 2, the carbonylated solution contained insoluble, unidentified solid. This was removed by centrifugation, and the supernatant liquid was then decanted and treated with hexane.

Anal. Calcd for C₃₃H₃₂O₂P₂Pt·0.5THF (2): C, 55.78; H, 4.81; P, 8.22. Found: C, 55.80; H, 4.77; P, 8.40. Calcd for C₃₅H₃₈O₂P₂Pt·0.1THF (4): C, 56.32; H, 5.18; P, 8.21; mol wt, 748. Found: C, 56.01; H, 5.21; P, 7.76; mol wt (CH₂Cl₂), 864.

1 was characterized on the basis of its IR and ¹³C NMR spectra. In all cases, the presence of THF of crystallization was confirmed by ¹H NMR spectroscopy; the proportions given are those that best fit the analytical data. The ¹³C-labeled analogues of 1-4 were prepared similarly from 90% enriched ¹³CO.

(2) (1,3-Bis(diphenylphosphino)propane)(cyclohexene-carbonyl)(hydroxycarbonyl)platinum(II), Pt(CO₂H)-(COC₆H₅)(dppp) (5). A sample of Pt(OH)(C₆H₅)(dppp) was prepared as described above from Pt(C₆H₅)(dppp) (75 mg, 0.109 mmol) in THF (8 mL) and water (4 mL). The solid was dissolved in benzene (1 mL), and the solution was syringed into a tube under ca. 100 kPa of CO. The pressure was increased to 1200 kPa, and the solution was stirred at room temperature for 5 h. The voluminous white solid was removed by centrifugation, washed with hexane, and dried in vacuo to give 55 mg (0.072 mmol, 66%) of 5.

Anal. Calcd for C₃₅H₃₆O₃P₂Pt: C, 55.19; H, 4.76; P, 8.13; mol wt, 762. Found: C, 54.89; H, 4.85; P, 7.71; mol wt (CH₂Cl₂) 728.

If a pressure of 50-100 kPa is used, the reaction requires ca. 24 h at room temperature for completion.

(3) (1,3-Bis(diphenylphosphino)propane)(1-cyclohexenyl)(dimethylcarbamoyl)platinum(II), Pt(CON-

Me₂)(C₆H₅)(dppp) (6). A stirred solution of 3 (60 mg, 0.078 mmol) in THF (1 mL) was treated with gaseous dimethylamine (10 mL, ca. 0.45 mmol) from a syringe, and the mixture was allowed to stand at room temperature for 24 h. Addition of hexane precipitated a creamy solid that was recrystallized from THF/hexane to give colorless crystals of 6 (50 mg, 0.063 mmol, 81%). The labeled analogue Pt(¹³CONMe₂)(C₆H₅)(dppp) was prepared similarly from 90% ¹³C-enriched 3. Carbon analyses for 6 were inexplicably irreproducible: IR (KBr) 1583 (C=O) cm⁻¹.

Anal. Calcd for C₃₆H₄₁NOP₂Pt: C, 56.84; H, 5.43; N, 1.84. Found: C, 58.22, 55.69; H, 5.54, 5.37; N, 1.67, 1.75.

(4) (1,3-Bis(diphenylphosphino)propane)(1-cyclohexenyl)(*tert*-butylcarbamoyl)platinum(II), Pt(CONH-*t*-Bu)(C₆H₅)(dppp) (7). This was prepared similarly to 6 from 3 (60 mg, 78 mmol) and *tert*-butylamine (52 mg, 64 mmol). After the product had been recrystallized from THF/hexane, the yield of 7 was 52 mg (64 mmol, 82%): IR (KBr) 3420, 1566 (NH), 1538 (C=O) cm⁻¹.

Anal. Calcd for C₃₈H₄₅NOP₂Pt: C, 57.86; H, 5.75; N, 1.78. Found: C, 57.46; H, 5.76; N, 1.50.

Reaction of 3 with Alcohols. (1) **NMR Experiments.** An approximately 0.023 M solution of 3 (0.6 mL) in the appropriate deuterated solvent (see text) was allowed to react in an NMR tube with 0.55 μ L of dry methanol (ca. 0.14 mmol, 10% excess). The progress of the reaction was followed by monitoring the appearance of the Pt-CO₂CH₃ signal at δ 2.98.

(2) **IR Experiments.** In each case, 0.25 mL of ca. 0.16 M 3 in CH₂Cl₂ was mixed with a tenfold excess of the alcohol (ca. 0.4 mmol) and the solution was placed in a 0.1-mm path-length KBr cell. The progress of the reaction over a period of ca. 1 h was followed by monitoring the decay of the band at 1650 cm⁻¹ due to 3 and the concomitant growth of the ester absorption at ca. 1625 cm⁻¹. The reaction with methanol was complete within 1 min, those with ethanol and benzyl alcohol required ca. 25 min, and that with 2-propanol was incomplete after 1 h. With the assumption that the extinction coefficients of the $\nu(C=O)$ bands

Table V. Carbon-13 and Phosphorus-31 NMR Data for $[\text{Pt}(\text{COC}_6\text{H}_9)(\text{CO})(\text{dppp})]\text{BF}_4$ (8)^a

	δ_X	$^1J_{\text{PtX}}$, Hz	other coupling const (J , Hz) ^b
C ¹	221.8	591	P ² C ¹ , 80
C ²	174.9	1446	P ¹ C ² , 115
P ¹	-9.7	3413	P ¹ C ² , 115; P ¹ C ¹ , 4
P ²	-8.3	1420	P ² C ¹ , 80; P ² C ² , 9

^a Measured in CD_2Cl_2 ; see text for atom numbering.^b P¹P², 41 Hz.

in the acid and ester are equal, the extent of reaction with 2-propanol was estimated as 56% after 1 h.

There was no reaction between 3 and *tert*-butyl alcohol or triethylamine under the same conditions.


Reaction of 5 with Acid. A 0.046 M solution of 90% enriched $\text{Pt}(\text{COC}_6\text{H}_9)(\text{COC}_6\text{H}_9)(\text{dppp})$ in dichloromethane (1.7 mL) was treated in an NMR tube with an equimolar quantity of $\text{HBF}_4 \cdot \text{Et}_2\text{O}$ (9.7 μL , 0.078 mmol) dissolved in CD_2Cl_2 (0.1 mL). The sole product of the reaction was identified as $[\text{Pt}(\text{COC}_6\text{H}_9)(\text{CO})(\text{dppp})]\text{BF}_4$ on the basis of its ^{13}C and ^{31}P NMR spectra (Table V). Solvent was removed in vacuo to yield a yellow-brown solid: IR (KBr) 2056 s ($^{13}\text{C}=\text{O}$), 1589 m, 1568 m ($^{13}\text{C}=\text{O}$) cm^{-1} , (CH_2Cl_2) 2060 s ($^{13}\text{C}=\text{O}$), 1590 m, 1568 m ($^{13}\text{C}=\text{O}$) cm^{-1} .

Reaction of 3 with Acid. An ca. 0.16 M solution of 3 in dichloromethane (0.25 mL) was treated with an equimolar quantity of $\text{HBF}_4 \cdot \text{Et}_2\text{O}$ (5 μL , ca. 0.04 mmol) or of 40% aqueous HBF_4 (6 μL). The solution was transferred to a 0.1-mm path-length KBr cell, and its IR spectrum was scanned periodically over a 24-h period. The characteristic bands of 3 were replaced immediately after the addition by bands at ca. 2330 and 2107 cm^{-1} assigned, respectively, to CO_2 and $[\text{Pt}(\text{C}_6\text{H}_9)(\text{CO})(\text{dppp})]^+$. The former grew at the expense of the latter with time. When the experiment was carried out on ^{13}C -labeled 3 in CD_2Cl_2 , a peak due to $^{13}\text{CO}_2$ at δ 125.3 was observed in the ^{13}C NMR spectrum.


Reaction of 3 with CO. In a typical experiment, a 0.033 M solution of 90% enriched $\text{Pt}(\text{COC}_6\text{H}_9)(\text{C}_6\text{H}_9)(\text{dppp})$ in dichloromethane (2 mL) was stirred under CO (50–100 kPa). Samples (ca. 0.2 mL) were withdrawn periodically and their IR spectra recorded in a 0.1-mm path-length KBr cell. Over a period of 3 h the $\nu(^{13}\text{C}=\text{O})$ band of the starting material at 1565 cm^{-1} disappeared and was replaced by $\nu(^{12}\text{C}=\text{O})$ bands at 1605 and 1563 cm^{-1} due to the CO_2H and COC_6H_9 groups, respectively, of $\text{Pt}(\text{CO}_2\text{H})(\text{COC}_6\text{H}_9)(\text{dppp})$ (5).

Results

Preparation and Spectroscopic Characterization. Platinum(II) carboxylic acids of general formula $\text{Pt}(\text{C}_6\text{H}_9)(\text{C}_6\text{H}_9)(\text{P-P})$ have been prepared by insertion of carbon monoxide into the Pt–OH bond of the corresponding hydroxo complexes (eq 1). The latter are made




by protonation of the corresponding (cyclohexene)platinum(0) complexes with water (eq 2)¹⁵ and can be used directly in eq 1 without isolation or purification.



The reaction of CO with the hydroxoplatinum(II) complexes is complete within ca. 10 min at room temperature, but the yields of carboxylic acid depend strongly on the pressure of CO, on the bis(tertiary phosphine) present, on the solvent, and on the concentration of starting complex.

In the case of the complexes of dppp and dppb, treatment of ca. 1.5×10^{-4} M solutions of the hydroxo complexes with CO at slightly above atmospheric pressure gives the corresponding carboxylic acids 3 and 4 quantitatively. Poorer yields are obtained if higher concentrations of the hydroxo complexes are allowed to react with CO at or below atmospheric pressure. Separate experiments show that there is a competing reaction between the starting hydroxo complexes and the product carboxylic acids that gives poorly soluble, as yet uncharacterized solids. In the case of the dppe and vdpp complexes, this competing reaction occurs even under ca. 10 atm of CO, although yields of 1 and 2 can be increased by using dichloromethane in place of THF and by reducing the concentration of the hydroxo complex to ca. 2.5×10^{-5} M.

A further complication is that the initially formed complex $\text{Pt}(\text{CO}_2\text{H})(\text{C}_6\text{H}_9)(\text{dppp})$ (3) reacts further with CO at room temperature to give, after several hours, a second platinum(II) carboxylic acid, $\text{Pt}(\text{CO}_2\text{H})(\text{COC}_6\text{H}_9)(\text{dppp})$ (5), containing an acyl group formed by insertion of CO into the Pt–C₆H₉ bond (eq 3). The corresponding methyl



ester $\text{Pt}(\text{CO}_2\text{Me})(\text{C}_6\text{H}_9)(\text{dppp})$ reacts similarly with CO,¹⁷ and studies in progress show that most 1-cyclohexenyl dppp complexes of the type $\text{PtX}(\text{C}_6\text{H}_9)(\text{dppp})$ (X = various anionic ligands) behave in the same way. For complexes containing different bis(tertiary phosphines), the rate of insertion of CO into the Pt–C₆H₉ bond is in the order dppp > dppb >> dppe > vdpp. For this reason, under the conditions described in the Experimental Section, only the dppp system gives an acyl metallocarboxylic acid.

The platinum(II) carboxylic acids 1 and 2 are colorless, amorphous solids that are poorly soluble in water and most organic solvents. Complexes 3–5 are microcrystalline and insoluble in water but very soluble in CH_2Cl_2 , CHCl_3 , THF, and $(\text{CH}_3)_2\text{SO}$. Osmometric measurements on 3, 4, and 5 in dichloromethane, and on 3 in pyridine, show them to be monomeric in these solvents. The compounds have been characterized largely on the basis of NMR and IR spectroscopy. The ^{13}C NMR spectra of samples prepared by use of 90% enriched ^{13}CO exhibit a doublet of doublets at δ ca. 195, with ^{195}Pt satellites, due to the carboxylic carbon atom that is coupled to cis- and trans-phosphorus atoms [$^2J_{\text{CP}}(\text{trans}) = 130\text{--}160$ Hz; $^2J_{\text{CP}}(\text{cis}) = 8\text{--}12$ Hz] (Table I). The magnitude of $^1J_{\text{PtC}}$, ca. 1300 Hz, is similar to that reported for *trans*- $\text{PtCl}(\text{CO}_2\text{H})(\text{PET}_3)_2$ (1337 Hz)¹² and establishes that the carboxylic acid group is attached to platinum through carbon. The chemical shift of the carboxylic carbon atom varies very little for different bis(tertiary phosphines). It is ca. 20 ppm to higher frequency than that of *trans*- $\text{PtCl}(\text{CO}_2\text{H})(\text{PET}_3)_2$ and at slightly lower frequency than those of $(\eta\text{-C}_5\text{H}_5)\text{Fe}(\text{CO}_2\text{H})(\text{CO})(\text{PPh}_3)$ and $(\eta\text{-C}_5\text{H}_5)\text{Mo}(\text{CO}_2\text{H})(\text{CO})_2(\text{PPh}_3)$.⁹ Both the chemical shift and the magnitude of $^1J_{\text{PtC}}$ show little dependence on solvent (Table II).

The $^{31}\text{P}\{^1\text{H}\}$ NMR spectra of the carboxylic acids, measured at 80.98 MHz, consist of a slightly distorted AX pattern with ^{195}Pt satellites (Table III). The variation of the chemical shifts of the phosphorus nuclei of the dppe (1), dppp (3), and dppb (4) complexes reflects the incor-

poration of these atoms into five-, six-, and seven-membered rings, respectively.¹⁸ In the ³¹P NMR spectrum of Pt(¹³CO₂H)(C₆H₉)(dppp), the resonances at δ -1.2 (*J*_{PtP} = 1874 Hz) and -6.0 (*J*_{PtP} = 1610 Hz) show couplings with ¹³C of 153 and ca. 11 Hz, respectively, establishing that the first resonance is due to P¹ trans to CO₂H and the second signal is due to P² trans to C₆H₉. The resonances due to P¹ and P² for the remaining compounds have been assigned on the assumption that the relative magnitudes of ¹*J*_{PtP¹} and ¹*J*_{PtP²} remain similar. These magnitudes reflect the high trans influence of the C₆H₉ and CO₂H ligands and confirm that the carboxylic acid group is C bonded to platinum. As expected,¹⁹ the magnitude of ¹*J*_{PtP} trans to C₆H₉ (ca. 1600–1650 Hz) is less than that trans to the more electron-withdrawing carboxylic acid group (1870–1970 Hz).

The ¹³C NMR spectrum of the cyclohexenecarbonyl carboxylic acid **5**, made by use of 90% enriched ¹³CO, shows a doublet of doublets with ¹⁹⁵Pt satellites at δ 244.7 due to the acyl carbon atom, in addition to a similar pattern centered at δ 193.1 due to the carboxylic acid carbon. The chemical shift of the acyl carbon and the magnitudes of its associated coupling constants (Table V) are very similar to those reported for the chelate acyl complexes [Pt(OC₆H₄CO)(PBU₃)₂]⁺²⁰ and PtL₂COCOCPh=CH (L = various tertiary phosphines or 1/2 dppp)²¹ and confirm that carbon monoxide has inserted into the Pt–C₆H₉ bond, not into the Pt–CO₂H bond. In the ³¹P{¹H} NMR spectrum of **5**, the magnitude of ¹*J*_{PtP²} trans to C₆H₉CO, 1495 Hz, indicates that the cyclohexenylcarbonyl group has an even higher trans influence than cyclohexenyl. This finding is in agreement with the conclusion drawn from the Pt–Cl distances in *trans*-PtCl(CH₃)(PMePh₂)₂²² and *trans*-PtCl(COCH₃)(PMePh₂)₂²³ that acetyl has a higher trans influence than methyl.

The ¹H NMR spectra of the cyclohexenyl carboxylic acids 1–4 show no OH resonance in common NMR solvents such as CDCl₃, CD₂Cl₂, benzene-*d*₆, or toluene-*d*₈, either owing to self-exchange or because the signal is masked by the phenyl resonances of the bis(tertiary phosphines). However, in (CD₃)₂SO, a solvent that is known to suppress self-exchange of protons in organic compounds and also shifts OH resonances to high frequency, the OH resonances of 1–4 appear as a slightly broad singlet, without ¹⁹⁵Pt satellites, in the region δ 9–10; cf. (η-C₅H₅)Re(CO₂H)(CO)(NO) δ 9.47 (CD₂Cl₂)^{7c} and 8.6 (CD₃CN);^{7a} (η-C₅H₅)Re(N₂Ar)(CO₂H)(CO), δ 8.9 (solvent not specified);⁸ PtCl(CO₂H)(PEt₃)₂, δ 11.0 (toluene-*d*₈), 8.5 (CD₃CN), and 8.6 (benzene-*d*₆).¹² The OH resonance of **5** can be observed in most common NMR solvents, the chemical shifts being δ 8.05 (CD₂Cl₂), 9.06 (benzene-*d*₆), and 9.37 ((CD₃)₂SO). It is not known whether the absence of ¹⁹⁵Pt satellites is due to the small magnitude of ⁴*J*_{PtH}, to self-exchange, or to exchange with traces of moisture. Exchange of water with the carboxylic acid proton was demonstrated by double-resonance experiments. Thus, irradiation of the resonance due to water in (CD₃)₂SO solutions of **3** or **5**, or addition of D₂O to a solution of **3** in (CD₃)₂SO or of **5** in

benzene, caused the PtCO₂H signal to disappear.

The IR spectra of the platinum(II) carboxylic acids (Table IV) show a band due to ν(OH) at ca. 2680 cm⁻¹ in KBr disks that shifts to ca. 3440 cm⁻¹ in dichloromethane. The first value is within the range observed for hydrogen-bonded organic carboxylic acids (2700–2500 cm⁻¹), and the latter is only slightly below the range for monomeric carboxylic acids (3560–3500 cm⁻¹).²⁴ The IR spectrum of **3** in a mull of poly(chlorotrifluoroethylene), which is clear in the range of interest, showed the 2680 cm⁻¹ absorption but there was no band in the 3400–3600 cm⁻¹ region. In pyridine, the OH band of **3** appears at 2670 cm⁻¹, a value that is characteristic of a strongly hydrogen-bonded OH...N system.²⁵ These data suggest that the platinum(II) carboxylic acids are associated through hydrogen bonds in the solid state and confirm that they are monomeric in dichloromethane, a solvent of only moderate polarity. In contrast, organic carboxylic acids are associated through hydrogen bonds even in the gas phase.²⁴ Unfortunately, few data for other metallocarboxylic acids are available for comparison, viz., PtCl(CO₂H)(PEt₃)₂, 3140 cm⁻¹ (solid),¹² IrCl₂(CO₂H)(CO)(PMe₂Ph)₂, 3295 (solid) and 3313 cm⁻¹ (CHCl₃),⁵ (η-C₅H₅)Fe(CO₂H)(CO)(PPh₃), 2700 cm⁻¹ (solid),⁶ and (η-C₅H₅)Re(CO₂H)(CO)(NO), 2960 cm⁻¹ (THF).^{7c}

In a KBr disk, the ν(C=O) absorption of complexes **3**–**5** generally appears as a strong band with a shoulder of medium intensity on the high-frequency side in the region 1600–1650 cm⁻¹. In dichloromethane or pyridine, there is a single strong band at ca. 1650 cm⁻¹. The solid-state spectra of **1** and **2** show only one broad and ill-defined band at ca. 1650 cm⁻¹, perhaps because these complexes are amorphous rather than microcrystalline. A band or pair of bands in the region 1100–1200 cm⁻¹ may be due to a coupled ν(C–O)/δ_{OH} mode of the CO₂H group. Similar ν(C=O) and ν(C–O)/δ_{OH} absorptions have been reported for other metallocarboxylic acids; the former are somewhat lower than those observed for organic carboxylic acids (1650–1740 cm⁻¹).

The frequency of the ν(C=O) band of Pt-(C¹⁸OOH)(C₆H₉)(dppp)²⁶ in a KBr disk is equal, within experimental error, to that of unlabeled **3**, although in solution the band in the labeled compound is split (Table IV). If the CO₂H proton had ionized reversibly, the ¹⁸O label should have been distributed equally between the two possible sites and a band due to ν(C=O) at ca. 1610–1620 cm⁻¹ should have been observed.²⁷ Thus the lack of such a band supports the chemical observations (see below) that the platinum(II) carboxylic acids have negligible acidic properties. The conclusion must be regarded as tentative because we could not locate the band due to ν(C=O) in Pt(C¹⁸OOH)(C₆H₉)(dppp); it may be masked by ligand absorptions.

Chemical Properties. In contrast to *trans*-PtCl-(CO₂H)(PEt₃)₂, the platinum(II) carboxylic acids 1–5 show no tendency to decompose by elimination of CO₂, either in the solid state on exposure to air or in vacuo at tem-

(18) Garrou, P. E. *Chem. Rev.* 1981, 81, 229.

(19) Appleton, T. G.; Bennett, M. A. *Inorg. Chem.* 1978, 17, 738.

(20) Motoschi, H.; Pregosin, P. J. *Organomet. Chem.* 1979, 171, C37.

(21) Hamner, E. R.; Kemmitt, R. D. W.; Smith, M. A. R. *J. Chem. Soc., Dalton Trans.* 1977, 261.

(22) Bennett, M. A.; Ho, K. C.; Robertson, G. B. *Inorg. Chem.* 1979, 18, 1061.

(23) Bennett, M. A.; Ho, K. C.; Jeffery, J. C.; McLaughlin, G. M.; Robertson, G. B. *Aust. J. Chem.* 1982, 35, 1311.

(24) Bellamy, L. J. "The Infrared Spectra of Complex Molecules"; Chapman and Hall: London, 1975; p 183.

(25) Bellamy, L. J. "Advances in Infrared Group Frequencies"; Chapman and Hall: London, 1975; p 241.

(26) Prepared by treating Pt(C₆H₉)(dppp) successively with H₂¹⁸O and CO.


(27) The calculated isotope shift for a C=O bond behaving as a simple harmonic oscillator with a frequency of 1650 cm⁻¹ is 1650 – 1650(20/21)^{1/2}, i.e., 40 cm⁻¹. For ¹⁸O-labeled benzophenone the ν(C=O) band is at 1635 cm⁻¹, compared with 1664 cm⁻¹ in the unlabeled material (Halmann, M.; Pinchas, S. J. *Chem. Soc.* 1958, 1703). The corresponding shifts in the [ν(C=O)] bands at ca. 1570 cm⁻¹ in metal acetylacetonates range from 12 to 26 cm⁻¹ (Pinchas, S.; Silver, B. L.; Lauicht, I. J. *Chem. Phys.* 1967, 46, 1506. Musso, H.; Junge, H. *Tetrahedron Lett.* 1966, 33, 4003).

peratures up to 80 °C. They melt with decomposition above 100 °C, but the decomposition products have not yet been identified; preliminary studies show that a hydride is not formed. Solutions of the compounds in benzene, tetrahydrofuran, or dimethyl sulfoxide are stable in the absence of air, though a solution of **3** in dimethyl sulfoxide decomposes with loss of the CO₂H group at 90 °C. Solutions in dichloromethane or chloroform are stable for several hours but decompose on prolonged storage, probably by reaction with traces of HCl. The platinum(II) carboxylic acids fail the standard tests for organic carboxylic acids; e.g. they do not dissolve in 5% aqueous KOH, they neither dissolve in nor react with aqueous KHCO₃, and they are not deprotonated by triethylamine or 1,8-bis(dimethylamino)naphthalene. Complex **3** does react with 5 M aqueous KOH, *tert*-butyllithium, and triethylaluminum, but irreversible decomposition occurs and the products have not been identified. In contrast, treatment of *trans*-[PtH(CO)(P-*i*-Pr)₂]BPh₄ with an excess of KOH in aqueous THF has been reported to give the potassium salt of a platinum(II) carboxylic acid, *trans*-PtH(CO₂K)(P-*i*-Pr)₂.²⁸

Despite their lack of acidity, the platinum(II) carboxylic acids readily form esters. When a solution of **5** in CDCl₃ is treated with a small excess of methanol, the CO₂H resonance at δ 8.06 rapidly disappears and is replaced by a singlet with ¹⁹⁵Pt satellites at δ 2.98 ($J_{\text{PtH}} = 6.3$ Hz) due to the methyl ester Pt(CO₂CH₃)(COC₆H₅)(dppp). This compound has been made independently by treatment of the methoxo cyclohexenyl complex Pt(OCH₃)(C₆H₅)(dppp) with an excess of CO.¹⁷ Similarly, when a twofold excess of methanol is added to a solution of **3** in (CD₃)₂SO, the CO₂H resonance at δ 9.33 disappears over a period of a week and a new singlet due to Pt(CO₂CH₃)(C₆H₅)(dppp) appears at δ 2.82. Esterification occurs much more rapidly in other solvents, e.g., within minutes at room temperature in CDCl₃ or CD₂Cl₂ and within hours in benzene-*d*₆. The progress of the reaction in dichloromethane can also be followed by observing the decrease in intensity of the ν -(C=O) band of the acid at ca. 1650 cm⁻¹ and the corresponding growth of the ester band at ca. 1625 cm⁻¹. Qualitatively, for different alcohols, the rate of esterification is CH₃OH > C₂H₅OH \approx C₆H₅CH₂OH > (CH₃)₂CH-OH; *tert*-butyl alcohol does not form an ester.

When the methyl ester Pt(CO₂CH₃)(C₆H₅)(dppp) is dissolved in wet (CD₃)₂SO, a weak signal at δ 9.33 due to the corresponding acid is observed after several hours at room temperature. This shows that esterification is reversible, although formation of the methyl, ethyl, and benzyl esters is strongly favored.

Complex **3** also reacts readily with dimethylamine and *tert*-butylamine at room temperature to give the corresponding acid amides **6** and **7** (eq 4). Their IR spectra




show a ν (C=O) band at ca. 1580 cm⁻¹ that is typical for carbamoyl complexes²⁹ and is ca. 100 cm⁻¹ to lower fre-

quency than those for organic acid amides.³⁰ In addition to the usual cyclohexenyl and dppp resonances, the ¹H NMR spectrum of the dimethylcarbamoyl complex **6** shows two distinct N-CH₃ singlets at δ 2.28 and 2.59 owing to restricted rotation about the C-N bond. The lower field signal shows ¹⁹⁵Pt satellites ($J_{\text{PtH}} = 5.3$ Hz) and is assigned to the methyl group *trans* to platinum, following earlier work on platinum(II) carbamoyl complexes.³⁰ The ¹³C NMR spectra of the carbamoyl complexes made from Pt(¹³CO₂H)(C₆H₅)(dppp) show a doublet of doublets with ¹⁹⁵Pt satellites at δ ca. 199 due to the carbamoyl carbon atom ($J_{\text{PtC}} \approx 1200$ Hz, $J_{\text{CP}}(\text{trans}) \approx 140$ Hz, $J_{\text{CP}}(\text{cis}) \approx 11$ Hz), the chemical shift being in the range 183–218 ppm recently reported for a range of organometallic carbamoyl complexes.³¹ The ³¹P{¹H} NMR spectra of the carbamoyl complexes, like those of the parent acids, consist of an AX pattern with ¹⁹⁵Pt satellites. As in the case of **3**, assignment of the ³¹P chemical shifts has been based on the magnitude of J_{CP} derived from the ³¹P spectra of ¹³C-enriched complexes. Surprisingly, the magnitude of J_{PtP} *trans* to the carbamoyl ligands (1698 Hz in **6**; 1519 Hz in **7**) is lower than that *trans* either to CO₂H (1874 Hz) or to CO₂CH₃ (1882 Hz), which indicates that the carbamoyl ligands have a markedly higher *trans* influence than CO₂H or CO₂CH₃. In **7** the magnitude of J_{PtP} *trans* to C₆H₅, 1700 Hz, is ca. 100 Hz larger than in **6** and in a series of complexes of the type PtY(C₆H₅)(dppp).¹⁶ The cause of these effects is not known, although steric repulsion between the mutually *cis* *tert*-butylcarbamoyl and cyclohexenyl substituents probably plays a role.

The other platinum(II) carboxylic acids react with amines similarly to **3**, but in the case of **5** reaction occurs at both the carboxylic acid group and the acyl group, and the resulting mixture of products has not been identified.

The reactions of **1–5** with alcohols and amines to form esters and carbamoyl complexes, respectively, probably proceed by initial dissociation of OH⁻ to give a platinum(II) carbonyl cation (eq 5). This is supported by the order of



reactivity of **3** with alcohols, which closely parallels that found in a kinetic study of the reaction of *trans*-[PtCl(CO)(PPh₃)₂]⁺ with alcohols to form the esters *trans*-PtCl(CO₂R)(PPh₃)₂.³² The acid (η -C₅H₅)Fe(CO₂H)(CO)(PPh₃) ionizes completely in formamide (ϵ 109.5) to give [(η -C₅H₅)Fe(CO)₂(PPh₃)]⁺OH⁻, identified by its ν -(C=O) bands;⁶ in contrast, the IR spectrum of a solution of **3** in *N*-methylformamide ($\epsilon \sim 200$) shows no ν (C=O) band in the region 2500–1800 cm⁻¹, so the carbonyl cation must be present in undetectably low concentration. Further evidence for the ionization described by eq 5 is that dichloromethane solutions of ¹³CO-labeled **3** and **4** exchange completely with ¹²CO in the carboxylic acid group over a period of 1 h at room temperature; in the case of **3** exchange is accompanied by insertion into the Pt-C₆H₅ bond to give **5**. Likewise, complex **5** exchanges with ¹³CO only in the carboxylic acid group, the acyl C=O group being unaffected. The planar platinum(II) cations [Pt(C₆H₅)(CO)(P-P)]⁺ could be expected to exchange rapidly with ¹³CO by an S_N2 process; indeed [PtCl₂(CO)]₂ was reported many years ago to undergo “instantaneous” exchange with ¹⁴CO.³³

(28) Yoshida, T.; Ueda, Y.; Otsuka, S. *J. Am. Chem. Soc.* **1978**, *100*, 3941.

(29) Angelici, R. J. *Acc. Chem. Res.* **1972**, *5*, 335. Green, C. R.; Angelici, R. J. *Inorg. Chem.* **1972**, *11*, 2095.

(30) Reference 24, p 231.

(31) Angelici, R. J.; Formanek, T. *Inorg. Chim. Acta* **1983**, *76*, L9.


(32) Byrd, J. E.; Halpern, J. *J. Am. Chem. Soc.* **1971**, *93*, 1634.

(33) Wojcicki, A.; Basolo, F. *J. Am. Chem. Soc.* **1961**, *83*, 525.

Characteristic of many of the metallocarboxylic acids reported in the literature is the reaction with protons to produce metal carbonyl cations (eq 6), which often is the



reverse of the reaction used to prepare the acids. This reaction takes place quantitatively when **5** is treated with $HBF_4 \cdot OEt_2$ in dichloromethane to give the (cyclohexenecarbonyl)platinum(II) carbonyl cation **8**, which has a ter-




minal $\nu(^{13}CO)$ frequency of 2060 cm^{-1} . The complex has been identified unequivocally on the basis of its ^{13}C NMR spectrum, which shows resonances with the appropriate couplings for the acyl and terminal carbonyl carbon atoms (Table V). In the ^{31}P NMR spectrum, the Pt-P coupling constant for the phosphorus atom P^1 trans to terminal CO has the expected magnitude (3413 Hz), consistent with the low NMR trans influence of CO ,¹⁹ whereas J_{PP^2} is very low (1420 Hz), confirming the very high trans influence of the cyclohexenecarbonyl ligand.

The reaction of the cyclohexenyl complex **3** with either $HBF_4 \cdot OEt_2$ or 40% aqueous HBF_4 is more complex. The expected carbonyl cation $[Pt(C_6H_9)(CO)(dppp)]^+$ is formed, as shown by a $\nu(CO)$ band at 2107 cm^{-1} in the IR spectrum of the solution, but there is also a band at 2330 cm^{-1} assigned to free CO_2 that grows in intensity with time at the expense of the $\nu(CO)$ band. The presence of CO_2 was also confirmed by ^{13}C NMR spectroscopy, but the platinum-containing products could not be identified.

Discussion


The preparation of complexes 1–5 by insertion of CO into a M–OH bond represents an alternative approach to the synthesis of metallocarboxylic acids, which have generally been made by nucleophilic attack of OH^- on a carbonyl group attached to a strongly electron-withdrawing metal center. The difference between the two methods is more apparent than real, however, because our reaction probably goes via a platinum(II) carbonyl intermediate. This could be a salt, $[Pt(CO)(C_6H_9)(P-P)]OH$, or a neutral five-coordinate complex, $Pt(OH)(C_6H_9)(CO)(P-P)$. These intermediates are also likely to be involved in the reactions of the carboxylic acids with alcohols and amines (eq 5).

The spectroscopic data for 1–5, and for most other metallocarboxylic acids, suggest that there is strong electron donation from the metal to the hydroxycarbonyl group. This can be expressed in terms of a resonance contribution to the bonding from a dipolar form:



The clearest evidence comes from the $C=O$ stretching frequencies of 1–5, which are ca. 100 cm^{-1} to lower energy than those of typical organic carboxylic acids. It has been noted²⁹ that the $C=O$ stretching frequencies of carbamoyl complexes are likewise ca. 80 cm^{-1} lower than those of typical carboxamides, an effect we also observe in the case of **6** and **7**, and a similar explanation has been advanced. The ^{13}C chemical shifts of the bound carboxylic carbon atoms in 1–5 are 10–20 ppm to higher frequency (lower

Scheme 1. Decomposition of $Pt(CO_2H)(C_6H_9)(dppp)$ (**3**) by Acid



field) than those of organic carboxylic acids (δ 165–185).³⁴ A contribution from the dipolar resonance form **B** will make the carboxylic carbon atom more carbene-like, and in fact the chemical shifts of 1–5 are in the lower part of the range reported for platinum(II)–carbene complexes.³⁵ The fact that these chemical shifts are also at higher frequency than that of *trans*- $PtCl(CO_2H)(PEt_3)_2$ ¹² could be because tertiary phosphines are better σ -donors than Cl^- , although this does not account for the similarity of the Pt–C coupling constants between 1–5 and *trans*- $PtCl(CO_2H)(PEt_3)_2$.

Complexes 1–5 resemble organic carboxylic acids in being associated through hydrogen bonds in the solid state but differ from them in being monomeric in solution, even in a relatively nonpolar solvent such as dichloromethane. This is probably a reflection of the fact that 1–5 show no tendency to form carboxylate ions and are, therefore, weaker acids ($(\eta-C_5H_5)Re(CO_2H)(CO)(NO)$,^{7c} $(\eta-C_5H_5)Fe(CO_2H)(CO)(PPh_3)$ ⁶ and $(\eta-C_5H_4)Re(CO_2H)(CO)(NO)$,^{7c} both of which are soluble in KOH. Presumably this lack of acidity is a consequence of the presence of strongly electron-donating ligands in 1–5.

Complexes 1–5 differ from $(\eta-C_5H_5)Fe(CO_2H)(CO)(PPh_3)$ and $(\eta-C_5H_5)Re(CO_2H)(CO)(NO)$ in their thermal stability toward decarboxylation. For example, $(\eta-C_5H_5)Fe(CO_2H)(CO)(PPh_3)$ rapidly loses CO_2 on warming in benzene solution, whereas its potassium salt is stable to $100^\circ C$ in dry formamide.⁶ On the other hand, $(\eta-C_5H_5)Re(CO_2H)(CO)(NO)$ does not lose CO_2 in anhydrous acetone at $50^\circ C$; the anion $[(\eta-C_5H_5)Re(CO_2)(CO)(NO)]^-$ is readily decarboxylated on heating in dry solvents but is stabilized by water, presumably owing to hydrogen bonding.^{7c} We suggest that 1–5 are stable to decarboxylation because the bidentate phosphines and the σ -bonded carbon ligand do not readily dissociate, so that it is not easy to create a vacancy in the coordination sphere that could accommodate CO_2 , the product of β -elimination. Although a five-coordinate intermediate $PtH(C_6H_9)(CO_2)(P-P)$ might be thought to form readily, in practice this pathway seems to be energetically unfavorable. Evidence for this is the fact that the first step in the thermal decomposition of bis(alkyl) bis(tertiary phosphine) complexes $PtR_2(PR'_3)_2$ is dissociation of tertiary phosphine.³⁶ The first step in the facile thermal decarboxylation of *trans*- $PtCl(CO_2H)(PEt_3)_2$ ¹² may well be dissociation of Cl^- rather than of PEt_3 , since we have found that the phenyl derivative *trans*- $Pt(CO_2H)(C_6H_5)(PEt_3)_2$ is stable in benzene or dichloromethane at room temperature.¹⁶

The necessity for a vacant coordination site to allow β -elimination to occur also accounts for the formation of

(34) Levy, G. C.; Lichter, R. L.; Nelson, G. L. "Carbon-13 Nuclear Magnetic Resonance Spectroscopy", 2nd ed.; Wiley-Interscience: New York, 1980; p 143.

(35) Mann, B. E.; Taylor, B. E. ^{13}C NMR Data for Organometallic Compounds; Academic Press: London, 1981; p 141.

(36) McCarthy, T. J.; Nuzzo, R. G.; Whitesides, G. M. *J. Am. Chem. Soc.* 1981, 103, 3396 and references cited therein.

CO₂ when complex 3 is treated with acid. In this case, the vacant site is generated by protonation of the platinum(II)-cyclohexenyl bond and elimination of cyclohexene (Scheme I). Evidently the platinum(II)-acyl bond in 5 is not so readily cleaved by acid, so that β -elimination of CO₂ from the CO₂H group cannot occur and the alternative reaction, viz., removal of OH⁻ from the CO₂H group, takes over.

As a general conclusion, metallocarboxylic acids seem to be structurally analogous to organic carboxylic acids, but their chemistry is dominated by their abnormal dissociation to [L_nMCO]⁺OH⁻. Their acid-base behavior and the tendency of the CO₂H group to lose CO₂ depend sensitively on the nature of the metal and of the associated ligands in a way that will only be revealed by further studies on related systems.

Substitutional Reactivity of Dodecacarbonyltrimetal Complexes of Iron and Osmium[†]

Abdolrasul Shojale and Jim D. Atwood*

Department of Chemistry, State University of New York at Buffalo, Buffalo, New York 14214

Received September 12, 1984


The kinetics of the substitution reactions of Fe₃(CO)₁₂ and Os₃(CO)₁₂ have been investigated for L = PPh₃, PBu₃, P(OPh)₃, and P(OMe)₃ in hydrocarbon solution. The substitution of the iron cluster leads to both substitution and fragmentation products with a very small dependence on the entering ligand, typical of a CO dissociative interchange mechanism. Substitution of Fe₃(CO)₁₁PPh₃ leads only to fragmentation at a rate which is faster than substitution on Fe₃(CO)₁₂. Substitution on Os₃(CO)₁₂ occurs in an entering ligand independent reaction to Os₃(CO)₉L₃. The reactivities of these clusters are compared to the ruthenium analogue and to the mononuclear complexes M(CO)₅.

Mechanistic studies of transition-metal clusters are necessary to test the hypothesis that adjacent metal centers may lead to modes of reaction not found in mononuclear complexes.¹⁻⁶ Important questions about the mechanism, such as whether metal-metal bond cleavage occurs, can be answered by careful kinetics studies. The most thoroughly studied cluster system is the substitution reactions of Ir₄(CO)₁₂ and its substituted derivatives.⁷⁻¹¹ The reactivity was strongly influenced by the ligands on the cluster. For Ir₄(CO)₁₁L and Ir₄(CO)₁₀L₂ the donor ability of the ligand L was important to the reactivity while for Ir₄(CO)₉L₃ and Ir₄(CO)₈L₄ the size of L became predominant.⁸⁻¹¹ Both of these effects are found in mononuclear complexes. Only in the substitution reactions of Ir₄(CO)₁₂ which proceeded primarily through a mechanism dependent on the entering ligand did a difference from mononuclear reactions appear.⁷

Darensbourg and Incorvia have investigated the reactions of P(OMe)₃ substituted derivatives of Co₄(CO)₁₂.¹²⁻¹³ Reaction of the mono-substituted complex Co₄(CO)₁₁P-

(OMe)₃ with P(OMe)₃ proceeded by a second-order reaction, while reaction of the bis complex Co₄(CO)₁₀(P(OMe)₃)₂ with P(OMe)₃ to give Co₄(CO)₉(P(OMe)₃)₃ was first order in Co₄(CO)₁₀(P(OMe)₃)₂ and independent of P(OMe)₃.^{12,13}

Candlin and Shortland studied CO substitution of Ru₃(CO)₁₂ by PPh₃, PPh₂Et, PPhEt₂, PBu₃, and PPh₃ (eq 1).¹⁴ These reactions were reported to give only the



tris-substituted complex with no evidence of the mono- or bis-substituted species.¹⁴ The rate law observed was of the form

$$\text{rate} = \{k_1 + k_2[\text{L}]\}[\text{Ru}_3(\text{CO})_{12}]$$

very similar to that observed for mononuclear species such as Mo(CO)₆.^{14,15} Significantly, Ru(CO)₄PPh₃ did not yield Ru₃(CO)₉(PPh₃)₃ under conditions where CO was known to dissociate from Ru(CO)₄PPh₃.¹⁴ CO dissociation from the intact triruthenium complex was suggested as the mechanism for reaction.¹⁴ Subsequent work has confirmed these observations and shown that during reaction with PPh₃ the dissociation of CO is accelerated from Ru₃(CO)₁₁PPh₃ and Ru₃(CO)₁₀(PPh₃)₂ with respect to Ru₃(CO)₁₂.¹⁶⁻²⁰

Reaction of the iron and osmium analogues with phosphines have also been reported.^{21,22} For the osmium

[†] Dedicated to the memory of Earl Muetterties, a dedicated scientist, mentor, and friend.

- (1) King, R. B. *Prog. Inorg. Chem.* **1972**, *15*, 287.
- (2) Chini, R.; Longoni, G.; Albano, V. G. *Adv. Organomet. Chem.* **1976**, *14*, 285.
- (3) Humphries, A. P.; Kaesz, H. D. *Prog. Inorg. Chem.* **1979**, *25*, 145.
- (4) Muetterties, E. L. *Bull. Soc. Chim. Belg.* **1975**, *84*, 959.
- (5) Muetterties, E. L.; Rhodin, T. N.; Band, E.; Brucker, C. F.; Pretzer, W. R. *Chem. Rev.* **1979**, *79*, 91.
- (6) Tachikawa, M.; Muetterties, E. L. *Prog. Inorg. Chem.* **1981**, *28*, 203.
- (7) Sonnenberger, D.; Atwood, J. D. *Inorg. Chem.* **1981**, *20*, 3243.
- (8) Sonnenberger, D.; Atwood, J. D. *J. Am. Chem. Soc.* **1982**, *104*, 2113.
- (9) Sonnenberger, D.; Atwood, J. D. *Organometallics* **1982**, *1*, 694.
- (10) Darensbourg, D. J.; Baldwin-Zuschke, B. J. *J. Am. Chem. Soc.* **1981**, *104*, 3906.
- (11) Darensbourg, D. J.; Baldwin-Zuschke, B. J. *Inorg. Chem.* **1981**, *20*, 3846.
- (12) Darensbourg, D. J.; Incorvia, M. J. *J. Organomet. Chem.* **1979**, *171*, 89.
- (13) Darensbourg, D. J.; Incorvia, M. J. *Inorg. Chem.* **1980**, *19*, 2585.

- (14) Candlin, J. P.; Shortland, A. C. *J. Organomet. Chem.* **1969**, *16*, 289.
- (15) Graham, J. R.; Angelici, R. J. *Inorg. Chem.* **1967**, *6*, 2082.
- (16) Poe, A. J.; Twigg, M. V. *J. Organomet. Chem.* **1973**, *50*, C39.
- (17) Poe, A. J.; Twigg, M. V. *J. Chem. Soc., Dalton Trans.* **1974**, 1860.
- (18) Malik, S. K.; Poe, A. *Inorg. Chem.* **1978**, *17*, 1484.
- (19) Malik, S. K.; Poe, A. *Inorg. Chem.* **1979**, *18*, 1241.
- (20) Keeton, D. P.; Malik, S. K.; Poe, A. *J. Chem. Soc. Dalton Trans.* **1977**, 233.
- (21) Deeming, A. J.; Johnson, B. F. G.; Lewis, J. J. *Chem. Soc. A* **1970**, 897.