

Anomalous Dehydration of 2-Phenylindole-3-carboxamide to 2-Phenylindole-3-carbonitrile by Lithium Aluminum Hydride

Short Communication

Tonson Abraham^{a,*} and Albert J. Sicree^b

^a Research Institute, The University of Dayton, Dayton, OH 45469, U.S.A.

^b Air Force Materials Laboratory, Wright-Patterson Air Force Base, OH 45433, U.S.A.

(Received 18 May 1982. Accepted 2 June 1982)

Steric factors have been shown to be responsible for the anomalous dehydration of 2-phenylindole-3-carboxamide (**1**) to 2-phenylindole-3-carbonitrile (**3**) by lithium aluminum hydride. This steric effect was also reflected in the reactions of 2-phenylindole (**4**) and its derivatives.


Anomale Dehydratisierung von 2-Phenylindol-3-carbonsäureamid zu 2-Phenylindol-3-carbonitril mit Lithiumaluminiumhydrid (Kurze Mitteilung)

Es wird gezeigt, daß für die anomale Dehydratisierung von 2-Phenylindol-3-carbonsäureamid (**1**) zum entsprechenden Nitril **3** durch Lithiumaluminiumhydrid sterische Faktoren verantwortlich sind. Dieser sterische Effekt zeigt sich auch bei Reaktionen von 2-Phenylindol (**4**) und seiner Derivate.

(Keywords: Indoles; Reduction; Steric hindrance)

Carboxamides are normally reduced to amines by lithium aluminum hydride (*LAH*)¹. The unexpected dehydration of an indole carboxamide to the indole carbonitrile in the presence of an excess of *LAH* is reported here.

When 2-phenylindole-3-carboxamide (**1**) was allowed to react with an excess of *LAH* in refluxing tetrahydrofuran (*THF*), 2-phenylindole-3-carbonitrile (**3**) was isolated in 88% yield. Normally, one would expect *LAH* to reduce 1,*N*-unsubstituted indole-3-carboxamides to 3-methyl



derivatives, by analogy with the behavior of 1-unsubstituted indole-3-carboxylic acids towards LAH ². Although we are not aware of any reported reactions of 1,*N*-unsubstituted indole-3-carboxamides with LAH , 1-substituted-*N*-unsubstituted indole-3-carboxamides have been reduced to amines by LAH ³, which is analogous to the reduction of related indole derivatives⁴.

Steric hindrance due to the presence of the 2-phenyl group in **1** appeared to be responsible for the anomalous dehydration of **1** by LAH . This effect was reflected in the reactions of 2-phenylindole (**4**) and its derivatives.

For example, 2-phenylindolylmagnesium iodide (**5**), made from **4** and methylmagnesium iodide, failed to react with carbon dioxide when a *THF* solution of **5** was poured into dry-ice. The outcome was the same even when carbon dioxide was bubbled into a boiling solution of **5** in toluene. In both cases, **4** was recovered⁵. The expected⁶ 2-phenylindole-3-carboxylic acid was not formed. The steric resistance to electrophilic substitution at the 3-position of **4** could be overcome by using more powerful electrophiles. Thus **4** underwent electrophilic substitution in the presence of acetyl cyanide⁷ and reactive isocyanates⁸⁻¹⁰.

There appears to be a delicate balance between electronic and steric effects in the case of the reactions of **4** and its derivatives. While 2-methylindole **6** condensed with the acetyl cyanide at room temperature, **4** does not, unless an acid catalyst is used. The mode of condensation with acetyl cyanide at the 3-indolyl position of **4** and its derivatives then depends upon the substituent at the 1-indolyl position⁷. In another example, 2-oxo-2-(2-phenylindol-3-yl)acetonitrile (**7**) was allowed to react with ethanolic KOH to yield ethyl 2-phenylindole-3-carboxylate (**8**)¹¹. Once the positive character of the carbonyl carbon attached to the 3-indolyl position was reduced, as it was in **8** when compared to **7**, steric factors began to dominate, and **8** could not be hydrolyzed to the acid under normal ester hydrolysis conditions¹¹. Also, 2-phenylindole-3-carboxaldehyde (**9**) was reduced to 3-methyl-2-phenylindole (**10**) by LAH ⁴, whereas **1** resisted reduction under similar conditions.

The intermediate **2**, expected to be formed by the action of LAH on **1**, apparently underwent loss of an oxide of aluminum to yield the nitrile **3**. *N,N*-unsubstituted or *N*-monosubstituted carboxamides were less reactive towards LAH than the corresponding *N,N*-disubstituted derivatives¹², presumably due to the reduced reduction potential¹³ of

the intermediates of type **2**. However, since *LAH* reduced 2-methyl-1-substituted-*N*-unsubstituted indole-3-carboxamides to aminomethyl derivatives³, it appeared that steric rather than electronic factors prevented the reduction of **1**.

Reports of the dehydration of carboxamides to nitriles by the use of excess *LAH* are scarce. *Newman* and *Fukunaga* reported the isolation of nitriles by the reaction of *LAH* with carboxamides (four cases) only when the mole ratio of *LAH* to carboxamide was less than one¹⁴. 10,11-Dihydro-5*H*-dibenzo[*a,d*]cycloheptene-5-carboxamide and 2,2-diphenylacetamide have also been dehydrated to nitriles with *LAH*¹⁵. To our knowledge, these are the only reports of dehydration of carboxamides to carbonitriles with *LAH*. In other related chemistry, certain acyl hydrazines were found to be unreactive towards *LAH*¹⁶.

2-phenylindole-3-carboxamide (**1**) was obtained by the hydrolysis of the chlorosulfonyl intermediate that resulted by the action of 2-phenylindole on chlorosulfonyl isocyanate, as reported by *Mehta*, *Dhar* and *Suri*¹⁰. The reactions of indole derivatives with reactive isocyanates, including chlorosulfonyl isocyanate had previously been reported by *Seefelder*^{8,9}. Moreover, compounds **1**¹⁷ and **3**¹⁷⁻²⁰ reported by *Mehta* et al.¹⁰ were also previously known.

Experimental

2-Phenylindole-3-carboxamide (**1**)

This compound was prepared as previously described¹⁰, mp. 184-184.5° (ethyl acetate/heptane). Lit. 189°¹⁰ and 180-181°¹⁷. IR (KBr, cm⁻¹): 3510, 3400, 3340-2380 (br, $\nu_{\max}$ 3100), 1635, 1590, 1570, 1465, 1450, 1330, 1225, 745, 735, 690. PMR (*DMSO*-*d*₆, δ): 6.3-8.9 (m). MS (chemical ionization with methane): *M* + 1 (237).

2-Phenylindole-3-carbonitrile (**3**)

Under argon, 2-phenylindole-3-carboxamide (1.0 g, 4.2 mmol) was added slowly to a suspension of *LAH* (1.6 g, 42 mmol) in 50 ml of anhydrous *THF* (distilled from sodium/benzophenone). The reaction mixture was then refluxed for 3 h, cooled, and the excess *LAH* destroyed by careful addition of 15% NaOH aq. The aluminate salts were filtered off, and washed with ether. The combined organic phase was washed with water, dried (MgSO₄), and then evaporated to give **3** (88%).

Thin layer chromatography (Macherey-Nagel silica gel pre-coated plastic sheets) showed only one component [*R*_f = 0.5, 40% petroleum ether (bp. 30-60°)/60% ether] with *R*_f value identical with that of recrystallized (alcohol/water) material, mp. 240.5-241.5°. Lit. 240°¹⁰, 236-238°¹⁷, and 243°¹⁸. IR (KBr, cm⁻¹): 3410-2280 (br, $\nu_{\max}$ 3230), 2230, 1495, 1455, 1425, 1245, 735, 685. PMR (*DMSO*-*d*₆, δ): 7.2-8.3 (m). MS (chemical ionization with methane): *M* + 1 (219).

References

- ¹ *Hajós A.*, Complex Hydrides, pp. 94—97. New York: Elsevier. 1979.
- ² *Leete E., Marion L.*, Can. J. Chem. **31**, 775 (1953).
- ³ *Domschke G., Fürst H.*, Chem. Ber. **93**, 2907 (1960).
- ⁴ *Leete E.*, J. Amer. Chem. Soc. **81**, 6023 (1959).
- ⁵ *Abraham T.*, Ph. D. Dissertation, p. 125. Washington, D. C.: The Catholic University of America. 1976.
- ⁶ *Majima R., Kotake M.*, Chem. Ber. **63**, 2237 (1930).
- ⁷ *Kiang A. K., Mann F. G.*, J. Chem. Soc. **1953**, 594.
- ⁸ *Seefelder M.*, Ger. Offen. **1,198,367**. Nov. 12, 1965. C. A. **63**, 14817 c (1965).
- ⁹ *Seefelder M.*, Chem. Ber. **96**, 3243 (1963).
- ¹⁰ *Mehta G., Dhar D. N., Suris S. C.*, Synthesis **1978**, 374.
- ¹¹ *Spiro V., Madonia P.*, Gazz. Chim. ital. **87**, 454 (1957).
- ¹² *Mićović V. M., Michailović M.*, J. Org. Chem. **18**, 1190 (1950).
- ¹³ Chemical and Engineering News **59**, 26 (1981).
- ¹⁴ *Newman M. S., Fukunaga I.*, J. Amer. Chem. Soc. **82**, 693 (1960).
- ¹⁵ *Humber L. G., Davis M. A.*, Can. J. Chem. **44**, 2113 (1966).
- ¹⁶ *Hinman R. L.*, J. Amer. Chem. Soc. **78**, 2463 (1956).
- ¹⁷ *Nishiwaka T., Azechi K., Fujiyana F.*, J. Chem. Soc. Perkin I **1974**, 1867.
- ¹⁸ *Loudon J. D., Tennant G.*, J. Chem. Soc. **1960**, 3466.
- ¹⁹ *Tominaga Y., Natsuki R., Matsuda Y., Kobayashi G.*, Yakugaku Zasshi **93**, 1433 (1973). C. A. **80**, 47769 z (1974).
- ²⁰ *Wentrup C.*, Advances in Heterocyclic Chemistry **28**, 336 (1981).