Tetrahedron Letters 55 (2014) 232-234

Contents lists available at ScienceDirect

Tetrahedron Letters

journal homepage: www.elsevier.com/locate/tetlet

Vanadium-catalyzed oxidative Strecker reaction: α-C–H cyanation of *para*-methoxyphenyl (PMP)-protected primary amines

Chen Zhu, Ji-Bao Xia, Chuo Chen*

Division of Chemistry, Department of Biochemistry, University of Texas Southwestern Medical Center, 5323 Harry Hines Boulevard, Dallas, TX 75390-9038, United States

ARTICLE INFO

ABSTRACT

Article history: Received 26 October 2013 Revised 31 October 2013 Accepted 1 November 2013 Available online 15 November 2013

Keywords: Vanadium catalyst C-H cyanation Oxidative Strecker reaction Primary amine Schiff base ligand

Cyanide, or nitrile, is a valuable one-carbon unit. It can be converted into a variety of polar functional groups, for example, amine, aldehyde, ketone, carboxylic acid, and amide. The Strecker reaction is the most commonly used cyanation reaction, which gives α -aminonitriles from imines.¹ Imines are normally prepared by condensing amines and aldehydes (Fig. 1, top), but can also be generated by oxidizing amines (Fig. 1, bottom). Recently, this oxidative approach has been recognized as a powerful new way to functionalize amines.

The oxidative Strecker reaction is a cross-dehydrogenative coupling (CDC) reaction.² It provides a more direct access to α -aminonitriles than the traditional Strecker reaction. It also uses the amine building blocks in a different way. Instead of cyanating aldehydes at the carbonyl position after imine formation, this one-carbon extension reaction functionalizes amines at the α -C-position. The focus of current research on this type of CDC reaction is the oxidative functionalization of tertiary amines, in particular, *N*,*N*-dialkylaniline and tetrahydroisoquinoline.^{3,4} Issues associated with primary and secondary amines arise from the N–H oxidation step,^{5,6} which often yields nitrones or *N*-oxides instead of imines.⁷

Vanadium complexes have been used widely to catalyze the oxidation of olefins, thioethers, alcohols, and phenols, but not amines.⁸ There is only one report of vanadium-catalyzed CDC reaction, specifically, oxidative Strecker reaction of tertiary amines.⁹ As part of a program to expand the utility of vanadium complexes, we searched for vanadium catalyst systems that promote the N–H oxidation. We report here that vanadium-Schiff base complexes can catalyze oxidative Strecker reaction of PMP-protected primary amines.

We describe an oxidative Strecker reaction that allows for direct cyanation of *para*-methoxyphenyl

(PMP)-protected primary amines. A vanadium(V) complex was used as the catalyst and TBHP as the oxi-

dant. The cyanation occurs at the α -C position bearing either an alkyl or an aromatic group. This method

provides a direct access to α -aminonitrile from amines with one-carbon extension.

We recently found that vanadocene dichloride (Cp₂VCl₂) catalyzes benzylic C–H oxidation selectively and efficiently.¹⁰ We hypothesized that high-valent vanadium complexes could also oxidize the weakly activated α -C–H group of amines and promote a Strecker reaction. We envisioned that methods for cyanating primary amines would be most versatile because the nitrogen center of the products could be further functionalized. We chose to use the electron-rich *para*-methoxyphenyl (PMP) protecting group to promote the oxidation and circumvent the self-condensation issue of primary amines. After C–H cyanation, the *N*-PMP group can be removed easily to give α -aminonitriles that bear a primary amino group.¹¹









etrahedro



© 2013 Elsevier Ltd. All rights reserved.

^{*} Corresponding author. Tel.: +1 214 648 5048; fax: +1 214 648 0320. *E-mail address*: Chuo.Chen@UTSouthwestern.edu (C. Chen).

^{0040-4039/\$ -} see front matter @ 2013 Elsevier Ltd. All rights reserved. http://dx.doi.org/10.1016/j.tetlet.2013.11.012

Table 1

Development of the vanadium-catalyzed oxidative α -C-cyanation reaction^a



 $^a\,$ Reaction conditions: 1 (0.1 mmol), TBHP (70 wt % in water, 0.15 mmol), TMSCN (0.12 mmol), 1 mL solvent.



We began our study by searching for an active vanadium catalyst at 5 mol % loading. We used PMP-benzylamine (**1**) as the substrate, TBHP as the oxidant, and TMSCN as the cyanide source (Table 1). All the vanadium(III) and vanadium(IV) complexes we examined were not able to promote this reaction (entries 1 and 2). However, various vanadium(V) complexes were found active (entry 3), with the vanadium(V)-Schiff base complex **3a** catalyzing the oxidative cyanation of **1** to give **2** in 50% yield (entry 4). Introducing two sterically bulky and slightly electron-donating *tert*-butyl groups to the aromatic ring of the Schiff base ligand resulted in a decreased reaction rate (entry 5), whereas adding an electron-withdrawing nitro group led to a complex mixture of products (entry 6). Acetonitrile was proved to be the solvent of choice (entries 7–11). The yield of **2** could be improved to 61% by increasing the reaction concentration to 0.2 M (entry 12).

The generality of this vanadium-catalyzed oxidative Strecker reaction is shown in Table 2. A variety of PMP-protected primary amines can be cyanated easily. For substituted benzylamines, both electron-donating and withdrawing groups can be tolerated at the *para*, *meta*, or *ortho* position (**4**–**9**). In general, electron-rich benzylamines (**4** and **5**) are more reactive than the electron-deficient ones (**6**–**9**). Aromatic rings other than phenyl can also be tolerated. For example, both the PMP-protected naphthylmethylamine and thiophenylmethylamine were cyanated smoothly (**10** and **11**). Importantly, activation of the α -position by an aromatic group is not necessary. This oxidative Strecker reaction can be used to functionalizing PMP-protected aliphatic amines (**12–14**).

This oxidative Strecker reaction allows us to make use of amine building blocks to prepare α -aminonitriles that cannot be easily accessed by traditional methods. For example, α -C-cyanation of the PMP-protected (+)-dehydroabietylamine (**15**) gave **16** in 61% yield as a 2:1 mixture of diastereomers (Fig. 2). **15** was synthesized by copper-catalyzed *N*-arylation of (+)-dehydroabietylamine (see Supplementary Material for details).

Table 2

Scope of the α -C-cyanation reaction^{a,b}



^a Reaction conditions: amine (0.2 mmol), TBHP (70 wt % in water, 0.3 mmol), TMSCN (0.24 mmol), 1 mL CH₃CN.

^o Isolated yields. PMP = para-methoxyphenyl (p-MeOPh).

Mechanistically, we believe that **3a** first reacted with TBHP to give I as the active catalyst (Fig. 3). The PMP-amine then coordinated to I and formed complex II. Similar to other metal-catalyzed amine oxidation reactions,² a single-electron transfer (SET) from the electron-rich nitrogen center to the vanadium complex occurred to give III, a valence tautomer of II. Subsequent O-O homolysis and C-H abstraction gave IV and the iminium ion, which reacted with cyanide to give the α -aminonitrile. Support for the SET hypothesis followed from the observation that electron deficient acetyl, tosyl, and Boc-protected benzylamines did not react under these conditions. In addition, vanadium(III) complexes could not promote this oxidative Strecker reaction, suggesting that a two-electron vanadium(III/V) redox mechanism was not operative. We also believe that amine oxidation was not mediated by tert-butoxyl radical,¹² because vanadium(IV) complexes did not catalyze the reaction, and vanadium(V) complexes other than 3 gave only less than 10% conversion. For the same reason, we believe that the aminium ion was coordinated to vanadium (III) when the C-H abstraction occurred. The Schiff base ligand likely served as an electron sink to facilitate the SET. We further propose that the silvl transfer from TMSCN to the oxide ligand of III or IV



Figure 2. Functionalization of natural product dehydroabietylamine. Reaction conditions: **15** (0.2 mmol), TBHP (70 wt % in water, 0.3 mmol), TMSCN (0.24 mmol), 1 mL CH₃CN. PMP = *para*-methoxyphenyl (*p*-MeOPh).



Figure 3. Proposed mechanism for the vanadium-catalyzed $\alpha\text{-C-cyanation}$ reaction.

promoted the regeneration of **I** and released the cyanide for the Strecker reaction.

In summary, we have developed a vanadium-catalyzed oxidative Strecker reaction of PMP-protected primary amines. This α -C–H cyanation reaction allows for functionalization of amines at the α -C-position, and is complimentary to the traditional Strecker reaction which functionalizes aldehydes. Efforts are underway to further explore the utility of vanadium-catalyzed CDC reactions.

Acknowledgements

Financial Support was provided by NIH (NIGMS R01-GM079554) and the Welch Foundation (I-1596). C.C. is a South-western Medical Foundation Scholar in Biomedical Research.

Supplementary data

Supplementary data (experimental procedures and characterization data of the reaction products) associated with this article can be found, in the online version, at http://dx.doi.org/10.1016/ j.tetlet.2013.11.012.

References and notes

- (a) Gröger, H. Chem. Rev. 2003, 103, 2795–2828; (b) Enders, D.; Shilvock, J. P. Chem. Soc. Rev. 2000, 29, 359–373; (c) Wang, J.; Liu, X.; Feng, X. Chem. Rev. 2011, 11, 6947–6983; (d) Zuend, S. J.; Coughlin, M. P.; Lalonde, M. P.; Jacobsen, E. N. Nature 2009, 461, 968–970.
- (a) Murahashi, S.-I. Pure Appl. Chem. 1992, 64, 403–412; (b) Murahashi, S.-I. Angew. Chem., Int. Ed. 1995, 34, 2443–2465; (c) Murahashi, S.-I.; Zhang, D. Chem. Soc. Rev. 2008, 37, 1490–1501; (d) Campos, K. R. Chem. Soc. Rev. 2007, 36, 1069–1084; (e) Li, C.-J. Acc. Chem. Res. 2009, 42, 335–344; (f) Yoo, W.-J.; Li, C.-J. Top. Curr. Chem. 2010, 292, 281–302; (g) Yeung, C. S.; Dong, V. M. Chem. Rev. 2011, 111, 1215–1292; (h) Sun, C.-L.; Li, B.-J.; Shi, Z.-J. Chem. Rev. 2011, 111, 1293–1314; (i) Liu, C.; Zhang, H.; Shi, W.; Lei, A. Chem. Rev. 2011, 111, 1780– 1824.
- For examples of CDC reactions of secondary amines: (a) Choi, H.; Doyle, M. P. Chem. Commun. 2007, 745–747; (b) Zhao, L; Li, C.-J. Angew. Chem., Int. Ed. 2008, 47, 7075–7078; (c) Zhao, L; Baslé, O.; Li, C.-J. Proc. Natl. Acad. Sci. 2009, 106, 4106–4111; (d) Xie, J.; Huang, Z.-Z. Angew. Chem., Int. Ed. 2010, 49, 10181– 10185; (e) Zhang, G.; Zhang, Y.; Wang, R. Angew. Chem., Int. Ed. 2011, 50, 10429–10432; (f) Wu, J.-C.; Song, R.-J.; Wang, Z.-Q.; Huang, X.-C.; Xie, Y.-X.; Li, P. D. J.-H. Angew. Chem., Int. Ed. 2012, 51, 3453–3457; (g) Li, K.; Tan, G.; Huang, J.; Song, F.; You, J. Angew. Chem., Int. Ed. 2012. http://dx.doi.org/10.1002/ anie.201306181.
- For examples of oxidative Strecker reaction of secondary amines: (a) Barton, D. H. R.; Billion, A.; Boivin, J. *Tetrahedron Lett.* **1985**, *26*, 1229–1232; (b) Köhler, V.; Bailey, K. R.; Znabet, A.; Raftery, J.; Helliwell, M.; Turner, N. J. *Angew. Chem., Int. Ed.* **2010**, *49*, 2182–2184; (c) Sonobe, T.; Oisaki, K.; Kanai, M. Chem. Sci. **2012**, 3, 3249–3255.
- a Ref. 2a-c; (b) Enders, D.; Shilvock, J. P. Chem. Soc. Rev. 2000, 29, 359–373; (c) Largeron, M. Eur. J. Org. Chem. 2013, 5225–5235.
- (a) Martin, G. M.; Benditt, E. P.; Eriksen, N. Nature 1960, 186, 884–885; (b) Chu, G.; Li, C. Org. Biomol. Chem. 2010, 8, 4716–4719; (c) Murahashi, S.-I.; Naota, T.; Taki, H. J. Chem. Soc. Chem. Commun. 1985, 613–614; (d) Samec, J. S. M.; Éll, A. H.; Bäckvall, J.-E. Chem. Eur. J. 2005, 11, 2327–2334; (e) Murahashi, S.-I.; Okano, Y.; Sato, H.; Nakae, T.; Komiya, N. Synlett 2007, 1675–1678; (f) Nishinaga, r.; Yamazaki, S.; Matsuura, T. Tetrahedron Lett. 1988, 33, 4115–4118; (g) Larsen, J.; Jørgensen, K. A. J. Chem. Soc., Perkin Trans. 1992, 2, 1213–1217; (h) Nicolaou, K. C.; Mathison, C. J. N.; Montagnon, T. Angew. Chem., Int. Ed. 2003, 42, 4077–4082; (i) Patila, R. D.; Adimurthy, S. Adv. Synth. Catal. 2011, 353, 1695–1700; (j) Huang, B.; Tian, H.; Lin, S.; Xie, M.; Yu, X.; Xu, Q. Tetrahedron Lett. 2013, 54, 2861–2864; (k) Largeron, M.; Fleury, M.-B. Angew. Chem., Int. Ed. 2012, 51, 5409– 5412; (m) Wendlandt, A. E.; Stahl, S. S. Org. Lett. 2012, 14, 2850–2853; (n) Ref. 3c.
- (a) Murahashi, S.-I.; Shiota, T. Tetrahedron Lett. **1987**, 28, 2383–2386; (b) Murahashi, S.; Oda, T.; Masui, Y. J. Am. Chem. Soc. **1989**, 111, 5002–5003; (c) Murahashi, S.; Mitsui, H.; Shiota, T.; Tsuda, T.; Watanabe, S. J. Org. Chem. **1990**, 55, 1736–1744; (d) Murahashi, S.-I.; Imada, Y.; Ohtake, H. J. Org. Chem. **1994**, 59, 6170–6172; (e) Imada, Y.; Iida, H.; Ono, S.; Murahashi, S.-I. J. Am. Chem. Soc. **2003**, 125, 2868–2869.
- (a) Sharpless, K. B.; Verhoeven, T. R. Aldrichim. Acta 1979, 12, 63–74; (b) Butler, A.; Clague, M. J.; Meister, G. E. Chem. Rev. 1994, 94, 625–638; (c) Hirao, T. Chem. Rev. 1997, 97, 2707–2724; (d) Conte, V.; Di Furia, F.; Licini, G. Appl. Catal., A 1997, 157, 335–361; (e) Ligtenbarg, A. G. J.; Hage, R.; Feringa, B. L. Coord. Chem. Rev. 2003, 237, 89–101.
- 9. Singhal, S.; Jain, S. L.; Sain, B. Chem. Commun. 2009, 2371–2372.
- 10. Xia, J.-B.; Cormier, K. W.; Chen, C. Chem. Sci. 2012, 3, 2240-2245.
- (a) Verkade, J. M. M.; van Hemert, L. J. C.; Quaedflieg, P. J. L. M.; Alsters, P. L.; van Delfta, F. L.; Rutjesa, F. P. J. T. *Tetrahedron Lett.* **2006**, *47*, 8109–8113; (b) Verkade, J. M. M.; van Hemert, L. J. C.; Quaedflieg, P. J. L. M.; Schoemaker, H. E.; Schümann, M.; van Delft, F. L.; Rutjes, F. P. J. T. *Adv. Synth. Catal.* **2007**, *349*, 1332–1336; (c) Koch, K.; van Weerdenburg, B. J. A.; Verkade, J. M. M.; Nieuwland, P. J.; Rutjes, F. P. J. T.; van Hest, J. C. M. *Org. Process Res. Dev.* **2009**, *13*, 1003–1006.
- 12. Ratnikov, M. O.; Doyle, M. P. J. Am. Chem. Soc. 2013, 135, 1549-1557.