

The Reaction of Malononitrile with Enamines of β -Ketocarbothionic Acid Anilides

Elzbieta Augustyn and Krystyna Bogdanowicz-Szwed *

Department of Organic Chemistry, Jagiellonian University, PL-30060 Kraków,
Poland

(Received 11 October 1982. Accepted 22 February 1983)

The reaction of β -morpholino-thiocinnamic acid anilides **1** with malononitrile yielded 2-arylimino-thiopyrans **3**, which in turn were converted under alkaline conditions into 6-thioxo-pyridines **5**. Their structures were elucidated on the basis of chemical properties and spectral data. The mechanism of the conversion **3** into **5** is discussed.

(Keywords: Thiopyrans; Thioxo-pyridines; Reactions with malonitrile)

Die Reaktion von Malononitril mit Enaminen von β -Ketocarbothionsäureaniliden

Die Reaktion von β -Morpholino-thiozimsäureaniliden **1** mit Malononitril lieferte 2-Arylimino-thiopyrane **3**, die unter alkalischen Bedingungen zu 6-Thioxo-pyridinen **5** umgesetzt wurden. Die Struktur der dabei erhaltenen Verbindungen wurde mittels chemischer und spektroskopischer Eigenschaften zugeordnet. Der Mechanismus der Konversion von **3** zu **5** wird diskutiert.

Introduction


Enamines of β -ketoacid anilides have attracted considerable interest as useful intermediates in organic synthesis; they react with a wide range of compounds providing some heterocyclic systems of pharmacological activity.

Recently we have reported¹ that enamines of cyclic β -ketocarbothionic acid anilides reacted easily with malononitrile yielding *o*-aminonitriles, derivatives of cycloalkeno-thiopyrans and cycloalkenopyridines with excellent yield. This results prompted us to develop this method for synthesis of similar heterocyclic systems from easily available starting materials i. e. β -morpholino-thiocinnamic acid anilides (enamines **1 a–1 e**) and mononitrile.

Results and Discussion


The reaction of enamines **1a-1e** with malononitrile in ethanolic or benzene solution proceed smoothly at reflux temperature affording products **3a-3e** in moderate yield. Taking into account our preliminary findings¹ we expect that this reaction may provide thiopyran **3** or pyridine **4** (Scheme 1). The structure elucidation of the obtained products **3a-3e** as thiopyran derivatives was based on analytical data as well as spectral studies.

Scheme 1




Both IR and ^1H -NMR spectra did not allow unambiguously to establish the structure of compounds **3**, because the isomeric pyridine derivatives **4** may have similar spectral features. Final structural assignment for **3a-3e** as thiopyran derivatives was based upon the detailed analysis of their mass spectra, which exhibited the characteristic fragmentation patterns²⁻⁴. The MS spectra of **3a-3e** revealed peaks of

Scheme 2



Scheme 3




On the basis of literature reports⁶ regarding the isomerisation of 2-arylimino-thiopyrans we suspected a conversion of **3** into **4**, hence the conversion of **3** into **5** was surprising. Nucleophilic attack of a hydroxide anion on the C-2 carbon atom involved an open-chain intermediate, which subsequently cyclized to **5**. The alternative isomerisation of **3** into **4** might proceed by a nucleophilic attack on the C-6 carbon atom. The exclusive formation of **5** suggests that the C-2 atom is the more preferred site for nucleophilic attack. The proposed mechanism of this conversion is outlined in Scheme 4.

In further experiments it was interesting to study the replacement of the sulphur atom by oxygen for compounds **5a-5e**. For this purpose we utilized the easy formation of S-alkyl derivatives by **5a-5e**. Compounds **5a-5e** reacted smoothly with dimethyl sulphate in boiling toluene yielding methylthiopyridines **6a-6e** with good yield. Crude solids of **6a-6e** reacted in turn with a concentrated solution of sodium

hydroxide to give **7a-7e**. The assigned structure was in two cases confirmed additionally by comparison with the samples independently prepared from β -morpholino-cinnamic acid anilides (**8a, 8c**) and malononitrile. S-Methyl derivatives **6a, 6b** underwent also nucleophilic substitution by *sec.* amines^{8,9} e.g. morpholine yielding **9a, 9b** with moderate yield. The reaction sequence is illustrated in Scheme 3.

Scheme 4



Experimental

Melting points are uncorrected. IR spectra were recorded on an UR-10 (Zeiss, Jena) spectrophotometer in Nujol mulls or in KBr disks. ¹H-NMR spectra were taken on a Jeol 100 spectrometer in deuterio-chloroform and DMSO (TMS as internal standard). Mass spectra were taken on an LKB-9000 S spectrometer. Elemental analyses were performed in the Regional Laboratory of Physico-Chemical Analyses and Structural Studies in Kraków. The analytical data for C, H, N, S, for all compounds are in full agreement with the proposed structures.

Enamines **1a-1e** and **8a-8c** were prepared according to the method described in Ref.¹⁰ and α -methylbenzylidenomalononitrile according to Ref.¹¹.

2-Arylimino-4-phenyl-5-cyano-2H-thiopyran (**3a-3e**)

General Procedure

A solution of malononitrile (0.02 mol) in ethanol or benzene (10 ml) was added to a solution of enamine **1** (0.02 mol) in ethanol (100 ml). The reaction mixture was boiled under reflux for 1 h. After cooling, the precipitated

crystalline yellow product was filtered off, washed with cold ethanol and crystallized from ethanol. Yellow prisms; average yield 37-70%. The compounds **3a**, **3c**, **3d** crystallized with one molecule of ethanol. For analyses the compounds were dried under reduced pressure at 80 °C.

3a: m. p. 156-158 °C; 67%; IR (Nujol): 3430, 3320, 3140 (NH₂), 2215 (CN), 1660 cm⁻¹ (C=N); ¹H-NMR (CDCl₃): 5.75 (m, 2 H, NH₂), 6.32 (s, 1 H, CH), 6.98-7.61 (m, 10 H arom.); MS (m/z): 303 (15.9%), 200 (100%).

3b: m. p. 166-167 °C; 53%; IR (Nujol): 3370, 3280, 3180 (NH₂), 2210 (CN), 1625 cm⁻¹ (C=N); ¹H-NMR (CDCl₃): 2.85 (s, 3 H, CH₃), 5.96 (m, 2 H, NH₂), 6.36 (s, 1 H, CH), 7.02-7.67 (m, 9 H arom.); MS (m/z): 317 (22.4%), 200 (100%).

3c: m. p. 187-188 °C; 46%; IR (Nujol): 3450, 3320, 3110 (NH₂), 2205 (CN), 1650 cm⁻¹ (C=N); ¹H-NMR (CDCl₃): 5.92 (m, 2 H, NH₂), 6.44 (s, 1 H, CH), 6.92-7.61 (m, 9 H arom.); MS (m/z): 337 (15.7%), 200 (100%).

3d: m. p. 198-201 °C; 70%; IR (Nujol): 3340, 3290, 3200 (NH₂), 2220 (CN), 1620 cm⁻¹ (C=N); ¹H-NMR (CDCl₃): 5.44 (m, 2 H, NH₂), 6.40 (s, 1 H, CH), 6.72-7.54 (m, 9 H arom.); MS (m/z): 381 (19.4%), 383 (17.1%), 200 (100%).

3e: m. p. 153-156 °C; 37%; IR (Nujol): 3370, 3260, 3220 (NH₂), 2210 (CN), 1625 cm⁻¹ (C=N); ¹H-NMR (CDCl₃): 3.92 (s, 3 H, CH₃), 5.82 (m, 2 H, NH₂), 6.31 (s, 1 H, CH), 7.02-7.56 (m, 9 H arom.); MS (m/z): 333 (33.1%), 200 (100%).

2-Arylamino-4-phenyl-5-cyano-6-thioxo-1,6-dihydropyridine (**5a-5e**)

General Procedure

To a solution of thiopyran **3** (0.05 mol) in 50 ml of ethanol 2 ml of a 5% solution of NaOH was added. The reaction mixture was refluxed for 30 min. The deep yellow solution was poured into 100 ml of ice water and the aqueous suspension was neutralised with dilute hydrochloric acid. The precipitate was filtered off, washed with water and purified by recrystallization from ethanol. Average yield 77-87%.

5a: 324-326 °C; 87%; IR (Nujol): 3460, 3305, 3210 (NH), 2220 cm⁻¹ (CN); ¹H-NMR (DMSO): 6.96 (s, 1 H, CH), 7.16 (s, 1 H, NH), 7.32 (s, 1 H, NH), 7.44-7.68 (m, 10 H arom.); MS (m/z): 303 (100%), 259 (13.8%, M⁺-CS).

5b: m. p. 257-258 °C; 82%; IR (Nujol): 3460, 3390, 3300 (NH), 2205 cm⁻¹ (CN); ¹H-NMR (DMSO): 2.95 (s, 3 H, CH₃), 6.88 (s, 1 H, CH), 7.44 (m, 2 H, 2 NH), 7.24-7.85 (m, 9 H arom.); MS (m/z): 317 (100%), 273 (14.3%, M⁺-CS).

5c: m. p. 267-268 °C; 77%; IR (Nujol): 3430, 3320, 3210 (NH), 2210 cm⁻¹ (CN); ¹H-NMR (DMSO): 6.94 (s, 1 H, CH), 7.61 (m, 2 H, 2 NH), 7.36-7.79 (m, 9 H arom.); MS (m/z): 337 (100%), 293 (15.3%, M⁺-CS).

5d: m. p. 287-288 °C; 83%; IR (Nujol): 3440, 3310, 3220 (NH), 2220 cm⁻¹ (CN); ¹H-NMR (DMSO): 6.82 (s, 1 H, CH), 7.32 (m, 2 H, 2 NH), 7.21-7.87 (m, 9 H arom.); MS (m/z): 381 (100%), 383 (88%), 337 (17.4%, M⁺-CS), 339 (11.7%, M⁺-CS).

5e: m. p. 264-265 °C; 81%; IR (Nujol): 3430, 3310, 3190 (NH), 2200 cm⁻¹ (CN); ¹H-NMR (DMSO): 3.92 (s, 3 H, CH₃), 6.93 (s, 1 H, CH), 7.21-7.32 (m, 2 H, 2 NH), 7.14-7.68 (m, 9 H arom.); MS (m/z): 333 (100%), 289 (16.8%, M⁺-CS).

1-Aryl-2-thioxo-4-phenyl-5-cyano-6-amino-1,2-dihydropyridine (**4a-4e**)

4a was prepared according to the method described in Ref.⁷. Compounds **4a-4e** were unknown and were synthesized in the same manner from α -methylbenzylidenomalononitrile and the appropriate arylisothiocyanate in DMF solution in the presence of triethylamine. Average yield 38-63%. The crude solids boiled in ethanol gave satisfactory analytical values for N. Because

of poor solubility in organic solvents it was difficult to record their ^1H -NMR spectra.

4a: m. p. 220–223 °C; IR (KBr): 3 060–3 330 (NH_2), 2 205 cm^{-1} (CN).

4b: m. p. 185–188 °C; IR (KBr): 3 180–3 460 (NH_2), 2 195 cm^{-1} (CN).

4c: m. p. 217–218 °C; IR (KBr): 3 200–3 450 (NH_2), 2 200 cm^{-1} (CN).

4d: m. p. 225–227 °C; IR (KBr): 3 160–3 370 (NH_2), 2 195 cm^{-1} (CN).

4e: m. p. 196–198 °C; IR (KBr): 3 220–3 450 (NH_2), 2 215 cm^{-1} (CN).

2-Arylamino-4-phenyl-5-cyano-6-methylthiopyridine (**6a–6e**)

A mixture of (1.5 g, 0.005 mol) of pyridine **5a**, dimethyl sulphate (1.3 g, 0.01 mol) and toluene (10 ml) was refluxed for 1 h. The precipitate was filtered off and washed with ligroin. The crude solid was suspended in 100 ml of water and neutralized with a dilute solution of NaOH. The precipitated deep yellow product was filtered off and crystallized from methanol. Yield 1.4 g (89%). Deep yellow prisms, m. p. 214–215 °C. IR (Nujol): 3 290, 3 280 (NH), 2 200 cm^{-1} (CN).

An analogous procedure was applied for the preparation of **6b–6e**. The crude solids were used without purification for further reactions.

2-Arylamino-4-phenyl-5-cyano-6-oxo-1,6-dihydropyridine (**7a–7e**)

a) By Hydrolysis of Compounds **6a–6e**

To a solution of 0.0025 mol of **6** in 20 ml of ethanol 10 ml of a 25% solution of NaOH was added. The reaction mixture was refluxed for 2.5 h. The mixture was dissolved in 100 ml of water and neutralized with dilute HCl. The precipitate was filtered off and crystallized from ethanol. Average yield 40–59%.

IR spectral data and m. p.'s for **7a**, **7b**, **7e** were identical with these reported in Ref.¹².

7c: m. p. 305–306 °C, colourless needles from methanol; IR (Nujol): 1 685 (CO), 2 220 (CN), 3 200, 3 330, 3 470 cm^{-1} (NH).

7d: m. p. 303–305 °C, colourless needles from ethanol; IR (Nujol): 1 670 (CO), 2 210 (CN), 3 190, 3 320, 3 470 cm^{-1} (NH).

b) By Condensation of Enamines **8a** or **8c** with Malononitrile

A mixture of 3.3 g (0.01 mol) of enamine **8a**, 0.07 g malononitrile (0.01 mol) was refluxed in 50 ml of benzene for 1 h. The precipitate was filtered off and crystallized from ethanol. Colourless needles with m. p. 284–286 °C¹²; yield 96%.

An analogous procedure was applied for the synthesis of **7c**. Colourless needles, m. p. 305–306 °C; yield 97%.

The identity of samples obtained by method a and b was confirmed by m. p., mixed m. p., and comparison of IR spectra.

2-Arylamino-4-phenyl-5-cyano-6-(*N*-morpholino)-pyridine (**9a**, **9b**)

To a solution of 1 g (0.003 mol) of **6a** in 10 ml of ethanol 5 g of morpholine was added. The reaction mixture was refluxed for 3 h. After cooling the mixture was poured into ice water and neutralized with dilute HCl. The semisolid product was filtered off and crystallized from ethanol. Pale yellow needlesh, m. p. 186–188 °C; yield 42%. IR (Nujol): 2 200 (NC), 3 350 cm^{-1} (NH).

An analogous procedure was applied for compound **9b**. Pale yellow prisms from ethanol, m. p. 241–243 °C; yield 36%. IR (Nujol): 2 205 (CN), 3 360 cm^{-1} (NH).

References

- ¹ Bogdanowicz-Szwed K., *Monatsh. Chem.* **113**, 583 (1982).
- ² Bogdanowicz-Szwed K., Nagraba K., *Org. Mass Spectrom.*, in press.
- ³ Bogdanowicz-Szwed K., Nagraba K., *Topics in Heterocyclic Chemistry, Proceedings of VIIth Symposium on Chemistry of Heterocyclic Compounds* (J. Kovač, ed.), Bratislava (1981).
- ⁴ Hamming M. C., Foster N. G., *Interpretation of Mass Spectra of Organic Compounds*. New York: Academic Press, 1972.
- ⁵ Otto H. H., Schmelz H., *Monatsh. Chem.* **111**, 53 (1980).
- ⁶ Gewald K., Buchwalder M., Peukert M., *J. prakt. Chem.* **315**, 679 (1973).
- ⁷ Gewald K., Liebscher J., Keydel M., *J. prakt. Chem.* **312**, 533 (1970); the authors did not report IR spectral data for **4a**.
- ⁸ Rudolf W. D., Augustin M., *J. prakt. Chem.* **323**, 55 (1981).
- ⁹ Schweiger K., Zigeuner G., *Monatsh. Chem.* **112**, 459 (1981).
- ¹⁰ Hünig S., Hübner K., Benzing E., *Ber.* **95**, 926 (1962).
- ¹¹ Cope A. C., *Org. Synth. Collectiv*, Vol. IV, 234 (1963).
- ¹² Zaleska B., Slusarska B., *Monatsh. Chem.* **112**, 1187 (1981).