## ChemComm

## COMMUNICATION



View Article Online View Journal | View Issue

## Peptide-catalyzed kinetic resolution of planar-chiral metallocenes<sup>†</sup>

Midori Akiyama,<sup>a</sup> Kengo Akagawa,<sup>a</sup> Hidetake Seino<sup>b</sup> and Kazuaki Kudo\*<sup>a</sup>

Cite this: Chem. Commun., 2014, 50, 7893

Received 2nd May 2014, Accepted 23rd May 2014

DOI: 10.1039/c4cc03266g

www.rsc.org/chemcomm

Kinetic resolution of racemic planar-chiral metallocenes was performed through the conjugate addition of a nucleophile to the enal part of substrates. While no enantiomeric discrimination was found with lowmolecular-weight organocatalysts, a properly designed resin-supported peptide catalyzed the reaction in a highly selective manner.

Planar chirality can be found in the groups of molecules with planar structures such as metallocenes,  $\pi$ -ligand complexes, and cyclophanes, and their enantioselective synthesis is necessary for the applications in chiral chemistry.<sup>1</sup> In spite of the recent remarkable progress in the field of asymmetric catalysis, the development of a versatile method for synthesizing chiral planar compounds in a catalytic manner is still a challenging task.<sup>2</sup>

Planar-chiral ferrocenes represent an important class of chiral compounds, and they have been mainly used as the ligands for metal-catalysts<sup>3</sup> and organocatalysts.<sup>1c,4</sup> Enantioenriched planar-chiral ferrocenes are usually obtained through the resolution of a racemic compound by chiral chromatography,<sup>3,4</sup> diastereo-selective transformation with the aid of a chiral directing group,<sup>5</sup> or the reaction with a stoichiometric amount of a chiral reagent.<sup>5*a*-*c*,6</sup> There are also some reports for the catalytic asymmetric synthesis of planar-chiral ferrocenes.<sup>5*c*,7</sup> In most cases, however, the applicability of these catalytic reactions is quite limited, and elaborately designed substrates are required to attain good enantioselectivity. Because planar chirality is the nature of the whole shape of a molecule, a catalyst with a large-scale asymmetric environment is necessary.

On the other hand, peptide catalysts can provide the large reaction sites suitable for substrate recognition by their specific three-dimensional structures.<sup>8</sup> Miller and co-workers<sup>9</sup> and Kawabata and co-workers<sup>10</sup> have reported regioselective reactions

with peptide- and peptide-related catalysts, respectively, in which small catalysts were not effective. Meanwhile, we have developed a resin-supported peptide catalyst consisting of a turn motif, p-Pro-Aib,<sup>11</sup> and a helical part (Fig. 1).<sup>12</sup> The N-terminal prolyl residue forms the iminium-ion intermediate with an  $\alpha$ , $\beta$ -unsaturated aldehyde to promote Michael addition of nucleophiles.<sup>13</sup> The turn structure of this peptide plays a key role in enantioselectivity of the reaction by covering one face of the peptide-bound substrate. The helical part is indispensable for stabilizing the turn structure. Because of a fairly large molecular size of the peptide, it is expected to be applicable for an enantioselective synthesis of planar chiral compounds. Herein, we report the kinetic resolution of racemic planar-chiral metallocene derivatives in the presence of a resin-supported peptide catalyst.

We first examined the kinetic resolution of racemic 1,2-disubstituted ferrocene **1a** in the 1,4-reduction of the enal part with a Hantzsch ester through the iminium-ion intermediate (Table 1).<sup>14</sup> The formyl group of this substrate is distant from the ferrocenyl moiety, hence it might be difficult for a low-molecular-weight chiral amine catalyst to discriminate the enantiomers. In fact, when the reaction was performed using an imidazolidinone catalyst **4** or **5**,<sup>15</sup> both the recovered starting material **1a** and reduced product **2a** were obtained as racemates (Table 1, entries 1 to 3). In contrast, the kinetic resolution took place in the presence of peptide catalyst **6** to give enantioenriched compounds **1a** and **2a** with a relative reaction rate ( $k_{rel}$ ) value<sup>16</sup> of 4.0 (Table 1, entry 4). The use of peptides 7 and **8**, which lack either the terminal turn structure or the helical part resulted in lower  $k_{rel}$  values (Table 1, entries 5 and 6). This indicates the importance of the structural motif of catalyst **6**.

The fifth amino acid residue from the N-terminus of the peptide was scanned to achieve higher selectivity because, according to our previous results,<sup>12</sup> the residue is likely to be

<sup>&</sup>lt;sup>a</sup> Institute of Industrial Science, University of Tokyo, 4-6-1 Komaba, Meguro-ku, Tokyo 153-8505, Japan. E-mail: kkudo@iis.u-tokyo.ac.jp

<sup>&</sup>lt;sup>b</sup> Faculty of Education and Human Studies, Akita University, Tegata-Gakuenmachi, Akita 010-8502, Japan

<sup>†</sup> Electronic supplementary information (ESI) available: Experimental procedures and full compound characterization, including NMR spectra and HPLC traces (PDF). CCDC 996655. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c4cc03266g

Pro-D-Pro-Aib-Trp-Trp-(AA)n-O :-NH-CH<sub>2</sub>-CH<sub>2</sub>-PEG-PS turn motif helical part amphiphilic resin

Fig. 1 Resin-supported peptide catalyst. Aib = 2-aminoisobutyric acid, AA = amino acid.

 Table 1
 Kinetic resolution of a planar-chiral ferrocene through the reduction of an unsaturated bond



<sup>*a*</sup> Conversion calculated from ees of **1a** and **2a** (ref. 16*a*). <sup>*b*</sup> Selectivity factor:  $k_{\rm rel} = k(R_p)/k(S_p)$  (ref. 16). <sup>*c*</sup> The reaction was performed with 0.5 equiv. of **3** for 3 h. <sup>*d*</sup> The reaction was performed in CHCl<sub>3</sub>. <sup>*e*</sup> Determined from <sup>1</sup>H NMR spectra of the crude mixture. <sup>*f*</sup> The hydrochloric acid salt of **5** was used.

Table 2 Screening for the 5th residue of the peptide catalyst

Fe rac-	I CHO	<b>3</b> (3.0 equiv) TFA•catalyst (20 mol <sup>4</sup> THF/H <sub>2</sub> O (1:2) 30 °C, 24 h	<sup>№</sup> ) онс (S <sub>р</sub> )	$ \begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} \end{array}\\ F_{e}\\ \end{array} + \\ \begin{array}{c} \end{array} + \\ \end{array} + \\ \begin{array}{c} \end{array} + \\ \begin{array}{c} \end{array} + \\ \end{array} + \\ \end{array} + \\ \begin{array}{c} \end{array} + \\ $	сно
catalyst Pro-D-P	: ro-Aib-Trp-A	A-(Leu-Leu-Aib)₂─-€		Nva: R = جَرْجَ Ser(Me): R = جَرْجَ Hse(Me): R = جَرْجَ	0
Entry	AA	<b>1a</b> [% ee]	2 <b>a</b> [% ee]	$\mathrm{C_{HPLC}}^{a}$ [%]	$k_{\rm rel}^{\ b}$
1	Ala	74	40	65	4.9
2	Nva	45	61	43	6.4
3	Ser(Me)	48	66	42	7.8
4	Hse(Me	e) 63	66	49	9.2
5	Met	70	61	53	8.4
6 <sup><i>c</i></sup>	Hse(Me	9 64	72	47	11.7

<sup>*a*</sup> Conversion calculated from ees of **1a** and **2a**. <sup>*b*</sup> Selectivity factor. <sup>*c*</sup> The hydrochloric acid salt of a peptide with (Leu-Leu-Aib)<sub>3</sub> as the helical part was used.

located around the ferrocenyl part of the iminium-ion intermediate (Table 2). The replacement of the fifth tryptophan with alanine increased the selectivity of the reaction (Table 2, entry 1), and introducing an amino acid with an unbranched alkyl side chain further enhanced the  $k_{\rm rel}$  value (entry 2). Moreover, the incorporation of an oxygen or a sulfur atom in the linear side chain was effective to improve the selectivity (Table 2, entries 3 to 5). Among them, the peptide possessing homoserine methyl ether was the best for the  $k_{\rm rel}$  value (Table 2, entry 4). After the optimization of the counter anion of the ammonio group of the terminal prolyl residue and the length of the helical part, the  $k_{\rm rel}$  value of 11.7 was achieved (Table 2, entry 6).

On the basis of the preference for the absolute configurations of the major enantiomers obtained by the peptide-catalyzed reaction, we postulated a possible mechanism for the kinetic



Fig. 2 Plausible mechanism for the kinetic resolution by the peptide catalyst.

resolution of planar-chiral ferrocene **1a** (Fig. 2). The conformation of the iminium-ion intermediate is considered to be regulated to avoid the steric repulsion between the iodo group on the cyclopentadienyl (Cp) ring and the vinylic hydrogen next to the iminium group (Fig. 2a). For the  $S_p$  enantiomer of **1a**, both faces of the iminium-ion intermediate are covered either by the Cp ring or by the peptide chain with a turn structure (Fig. 2b).<sup>12</sup> On the other hand, one face of the intermediate of the  $R_p$  enantiomer is accessible by the reagent, thus, the reaction proceeds faster than in the case of the  $S_p$  isomer. For the highly selective kinetic resolution, the fifth residue of the reductant from the upper side of the  $S_p$  enantiomer, whereas the formation of the iminium-ion intermediate with the  $R_p$  enantiomer should not be prevented by the steric repulsion caused by the Cp ring.

Compared to the case with a bulky side chain at the fifth residue of catalyst **6** (Table 1, entry 4), the introduction of the linear side chain in the catalyst was considered to be effective for this purpose (Table 2, entry 3). Although the role of the oxygen atom is not clear at present, an interaction with the cationic iminium-ion part might enhance the formation of the reactive intermediate for the  $R_p$  isomer, resulting in the higher selectivity (Table 2, entries 4 and 5).<sup>17</sup>

This type of kinetic resolution is potentially extendable to the reactions with other nucleophiles. Because the Michael addition of nitromethane to  $\alpha$ , $\beta$ -unsaturated aldehydes is a convenient method for the synthesis of  $\gamma$ -amino acid derivatives,<sup>18</sup> we employed nitromethane as a nucleophile. In the presence of the optimum catalyst **9**, the kinetic resolution proceeded with higher enantioselectivity than in the case with Hantzsch ester **3** (Scheme **1**). In this reaction, a new stereogenic center is created at the  $\beta$ -position of the aldehyde group. The configuration of the major product was consistent with the proposed model shown in Fig. 2, in which the addition occurred from the *Re* face of the *R*<sub>p</sub> isomer of substrate **1a**. The high diastereoselectivity of the reaction indicates the well-controlled conformation of the iminium-ion intermediate with regard to the direction of the ferrocenyl part (Fig. 2a).



Scheme 1 Peptide-catalyzed kinetic resolution through Michael addition of nitromethane.

Next, the scope of substrates for the kinetic resolution through the Michael addition of nitromethane by using peptide 9 was examined under optimized reaction conditions (Table 3). For the planar-chiral ferrocenes with halogen and alkynyl substituents, both unreacted starting material 1 and product 10 were obtained as highly enantioenriched forms (Table 3, entries 1 to 8). A shorter reaction time afforded product 10 with a high ee value (Table 3, entries 1, 3, 5, and 7). Meanwhile, upon increasing the conversion by elongating the reaction time, starting material 1 was obtained in a highly enantioselective manner (Table 3, entries 2, 4, 6, and 8). As to the substrate with an aromatic group, good enantioselectivity was achieved only for the product (Table 3, entry 9). In this case, a reaction conversion could not be increased even in a prolonged reaction time because of the low solubility of the substrate. In addition to the ferrocenes, ruthenocenyl substrate 1f could be used in this catalytic system (Table 3, entries 10 and 11).

Finally, the derivatizations of the obtained materials were conducted to demonstrate the utility of the present resolution method (Scheme 2). The Pinnick oxidation of product **10b** followed by an esterification afforded compound **11**, which could be further transformed to amino acid derivative **12** *via* a reduction and an N-protection. Because there are only limited examples for the asymmetric synthesis of amino acid derivatives with planar chirality,<sup>19</sup> this procedure offers a valuable way for

Table 3	Pe	otide	-catalyzed kinetic	c resolut	ion of planar-	chiral metallocenes
M M rac	, R , c-1	сно	MeNO <sub>2</sub> (5.0 equi 9 (20 mol%) 1,4-dioxane/H <sub>2</sub> O (1 30 °C	v) ► он 1:2)	c (Sp)-1	+ M (S,R <sub>p</sub> )-10
Entry	1	М	R	<i>t</i> [h]	1 [% ee (% Yield)]	<b>10</b> [% ee <sup><i>a</i></sup> (% Yield, d.r.)]
1	1a	Fe	I	10	65 (44)	92 (36, 98:2)
2	1a	Fe	Ι	12	90 (34)	59 (49, 93:7)
3	1b	Fe	Br	5	35 (53)	89 (25, 97:3)
4	1b	Fe	Br	12	83 (31)	86 (43, 93:7)
5	1c	Fe	Cl	8	31 (48)	93(18, >99:<1)
6	1c	Fe	Cl	14	77 (27)	80 (31, 94:6)
7	1d	Fe	1-Heptynyl	21	60 (38)	83 (37, 88:12)
8	1d	Fe	1-Heptynyl	36	84 (30)	70 (44, 83:17)
9	1e	Fe	p-MeOC <sub>6</sub> H <sub>4</sub>	10	42 (51)	85 (33, 95:5)
10	1f	Ru	Ī	4	48 (55)	87 (27, 92:8)
11	1f	Ru	Ι	5	87 (50)	66 (38, 82:18)

<sup>*a*</sup> ee of the major diastereomer.



Scheme 2 Transformation into planar-chiral derivatives.

their preparation. Halogen-substituted ferrocenyl compounds are synthetically useful, because they can easily be transformed by a coupling reaction.<sup>20</sup> The Suzuki–Miyaura reaction with iodine-containing **1a** and *p*-methoxyphenylboronic acid afforded coupling product **1e**. Alkyl ferrocene **13** could be prepared by the catalytic hydrogenation of product **10d**.

In conclusion, we have successfully shown that the peptide catalyst is applicable for the kinetic resolution of planar-chiral metallocenes which cannot be realized by low-molecular-weight organocatalysts. The use of an amino acid with a sterically less demanding, heteroatom-containing side chain at an appropriate position of the peptide sequence was the key for the high enantioselectivity. Planar chiral ferrocenes could tolerate several reaction conditions including oxidation and reduction. Further application of peptide catalysts for synthesizing other planarchiral compounds can be expected.

This work was financially supported by JSPS KAKENHI (23550116), and by MEXT KAKENHI (24105506).

## Notes and references

- Reviews: (a) C. Bolma and K. Muñiz, Chem. Soc. Rev., 1999, 28, 51;
   (b) S. E. Gibson and J. D. Knighta, Org. Biomol. Chem., 2003, 1, 1256;
   (c) O. R. P. David, in Comprehensive Enantioselective Organocatalysis: Catalysts, Reactions, and Applications, ed. P. I. Dalko, Wiley-VCH, Weinheim, 2013, pp. 195–219.
- (a) Y. Takemoto, Y. Baba, A. Honda, S. Nakao, I. Noguchi, C. Iwata, T. Tanaka and T. Ibuka, *Tetrahedron*, 1998, 54, 15567; (b) K. Kanda, T. Koike, K. Endo and T. Shibata, *Chem. Commun.*, 2009, 1870; (c) A. E. Garcia, S. Ouizem, X. Cheng, P. Romanens and E. P. Kündig, *Adv. Synth. Catal.*, 2010, 352, 2306; (d) M. Murai, J. Uenishi and M. Uemura, *Org. Lett.*, 2010, 12, 4788; (e) K. Tomooka, K. Uehara, R. Nishikawa, M. Suzuki and K. Igawa, *J. Am. Chem. Soc.*, 2010, 132, 9232.
- 3 Reviews: (a) T. Hayashi, in Ferrocenes: homogeneous catalysis, organic synthesis, material science, ed. A. Togni and T. Hayashi, Wiley-VCH,

Weinheim, 1995, pp. 105–142; (b) T. J. Colacot, Chem. Rev., 2003, **103**, 3101; (c) L.-X. Dai, T. Tu, S.-L. You, W.-P. Deng and X.-L. Hou, Acc. Chem. Res., 2003, **36**, 659; (d) P. Barbaro, C. Bianchini, G. Giambastiani and S. L. Parisel, Coord. Chem. Rev., 2004, **248**, 2131; (e) R. G. Arrayás, J. Adrio and J. C. Carretero, Angew. Chem., Int. Ed., 2006, **45**, 7674; (f) N. A. Butt, D. Liu and W. Zhang, Synlett, 2014, 615.

- 4 Reviews: (a) Y. Butsugan, S. Araki and M. Watanabe, in *Ferrocenes:* homogeneous catalysis, organic synthesis, material science, ed.
  A. Togni and T. Hayashi, Wiley-VCH, Weinheim, 1995, pp. 143–169;
  (b) G. C. Fu, Acc. Chem. Res., 2004, 37, 542.
- 5 (a) C. J. Richards and A. J. Locke, *Tetrahedron: Asymmetry*, 1998, 9, 2377; (b) R. C. J. Atkinson, V. C. Gibson and N. J. Long, *Chem. Soc. Rev.*, 2004, 33, 313; (c) D. Schaarschmidt and H. Lang, *Organometallics*, 2013, 32, 5668; (d) T. Aratani, T. Gonda and H. Nozaki, *Tetrahedron*, 1970, 26, 5453; (e) D. Marquarding, H. Klusacek, G. Gokel, P. Hoffmann and I. Ugi, *J. Am. Chem. Soc.*, 1970, 92, 5389; (f) O. Riant, O. Samuel and H. B. Kagan, *J. Am. Chem. Soc.*, 1993, 115, 5835; (g) B. F. Rehipre, O. Riant and L. Ricard, *Angew. Chem., Int. Ed. Engl.*, 1993, 32, 568; (h) Y. Nishibayashi and S. Uemura, *Synlett*, 1995, 79; (i) V. I. Sokolov, L. L. Troitskaya and O. A. Reutov, *J. Organomet. Chem.*, 1979, 182, 537.
- 6 (a) D. Price and N. S. Simpkins, *Tetrahedron Lett.*, 1995, 36, 6135;
  (b) M. Tsukazaki, M. Tinkl, A. Roglans, B. J. Chapell, N. J. Taylor and V. Snieckus, *J. Am. Chem. Soc.*, 1996, 118, 685; (c) R. S. Laufer, U. Veith, N. J. Taylor and V. Snieckus, *Org. Lett.*, 2000, 2, 629;
  (d) T. Fukuda, K. Imazato and M. Iwao, *Tetrahedron Lett.*, 2003, 44, 7503.
- 7 (a) S. Siegel and H.-G. Schmalz, Angew. Chem., Int. Ed. Engl., 1997, 36, 2456; (b) C. Genet, S. J. Canipa, P. O'Brien and S. Taylor, J. Am. Chem. Soc., 2006, 128, 9336; (c) A. Bueno, M. Rosol, J. García and A. Moyano, Adv. Synth. Catal., 2006, 348, 2590; (d) M. Ogasawara, S. Watanabe, L. Fan, K. Nakajima and T. Takahashi, Organo-metallics, 2006, 25, 5201; (e) A. Patti and S. Pedotti, Tetrahedron: Asymmetry, 2010, 21, 2631; (f) P. Steffen, C. Unkelbach, M. Christmann, W. Hiller and C. Strohmann, Angew. Chem., Int. Ed., 2013, 52, 9836; (g) D.-W. Gao, Y.-C. Shi, Q. Gu, Z.-L. Zhao and S.-L.

You, *J. Am. Chem. Soc.*, 2013, **135**, 86; (*h*) C. Pi, Y. Li, X. Cui, H. Zhang, Y. Han and Y. R. Wu, *Chem. Sci.*, 2013, **4**, 2675; (*i*) A.-N. Alba, P. Gómez-Sal, R. Rios and A. Moyano, *Tetrahedron: Asymmetry*, 2009, **20**, 1314; (*j*) R. Deng, Y. Huang, X. Ma, G. Li, R. Zhu, B. Wang, Y.-B. Kang and Z. Gu, *J. Am. Chem. Soc.*, 2014, **136**, 4472; (*k*) D.-W. Gao, Q. Yin, Q. Gu and S.-L. You, *J. Am. Chem. Soc.*, 2014, **136**, 4841.

- 8 Reviews: (a) E. A. C. Davie, S. M. Mennen, Y. Xu and S. J. Miller, *Chem. Rev.*, 2007, **107**, 5759; (b) H. Wennemers, *Chem. Commun.*, 2011, **47**, 12036.
- 9 (a) C. A. Lewis and S. J. Miller, Angew. Chem., Int. Ed., 2006, 45, 5616;
  (b) P. A. Lichtor and S. J. Miller, Nat. Chem., 2012, 4, 990;
  (c) T. P. Pathak and S. J. Miller, J. Am. Chem. Soc., 2013, 135, 8415;
  (d) C. T. Mbofana and S. J. Miller, J. Am. Chem. Soc., 2014, 136, 3285.
- 10 (a) T. Kawabata and T. Furuta, Chem. Lett., 2009, 38, 640;
   (b) K. Yoshida, T. Shigeta, T. Furuta and T. Kawabata, Chem. Commun., 2012, 48, 6981.
- 11 T. S. Haque, J. C. Little and S. H. Gellman, J. Am. Chem. Soc., 1996, 118, 6975.
- 12 K. Akagawa, R. Suzuki and K. Kudo, Adv. Synth. Catal., 2012, 354, 1280.
- 13 Reviews of organocatalysts: (a) A. Erkkilä, I. Majander and P. M. Pihko, *Chem. Rev.*, 2007, **107**, 5416; (b) S. Mukherjee, J. W. Yang, S. Hoffman and B. List, *Chem. Rev.*, 2007, **107**, 5471; (c) S. K. Panday, *Tetrahedron: Asymmetry*, 2011, **22**, 1817.
- 14 Review: S. G. Ouellet, A. M. Walji and D. W. C. MacMillan, Acc. Chem. Res., 2007, 40, 1327.
- 15 T. Passon, Synlett, 2008, 147.
- 16 (a) H. B. Kagan and J. C. Fiaud, in *Topics in stereochemistry*, ed.
  A. L. Allinger and E. L. Eliel, Wiley-VCH, Weinheim, 1988, vol. 18, pp. 249–330; (b) E. Vedejs, M. Jure, R. G. Arrayás, J. Adrio and J. C. Carretero, *Angew. Chem., Int. Ed.*, 2005, 44, 3974.
- 17 K. Akagawa, S. Takigawa, I. S. Nagamine, R. Umezawa and K. Kudo, *Org. Lett.*, 2013, **15**, 4964.
- 18 H. Gotoh, H. Ishikawa and Y. Hayashi, Org. Lett., 2007, 9, 5307.
- 19 L. Tebben, K. Bussmann, M. Hagemann, G. Kehr, R. Fröhlich and G. Erker, *Organometallics*, 2008, 27, 4269.
- 20 V. Mamane and O. Riant, Tetrahedron, 2001, 57, 2555.