

Mono(cyclopentadienyl)titanium complexes containing a sulfide-bridged bis(phenolato) ligand. Molecular structure of $\text{Ti}\{2,2'\text{-S}(\text{OC}_6\text{H}_2\text{-4-Me-6-}^t\text{Bu})_2\}(\eta^5\text{-C}_5\text{H}_5)\text{Cl}$

Francisco Amor ^a, Stefan Fokken ^b, Thomas Kleinhenn ^b, Thomas P. Spaniol ^b,
Jun Okuda ^{b,*}

^a Departamento de Química Inorgánica, Universidad de Alcalá, Campus Universitario, 28871 Alcalá de Henares, Spain

^b Institut für Anorganische Chemie und Analytische Chemie, Johannes Gutenberg-Universität Mainz, Duesbergweg 10-14, 55099 Mainz, Germany

Received 25 August 2000; accepted 22 September 2000

Dedicated to Professor H. Brunner on the occasion of his 65th birthday

Abstract

A series of titanium complexes containing a sulfur-bridged chelating bis(phenolato) ligand, $\text{Ti}\{2,2'\text{-S}(\text{OC}_6\text{H}_2\text{-4-Me-6-}^t\text{Bu})_2\}(\eta^5\text{-C}_5\text{R}_5)\text{Cl}$ ($\text{R} = \text{H, Me}$; $\text{R}_5 = 1,3\text{-(SiMe}_3)_2\text{H}_3$), were prepared by the reaction of the dilithium derivative $\text{Li}_2[2,2'\text{-S}(\text{OC}_6\text{H}_2\text{-4-Me-6-}^t\text{Bu})_2]$ with $\text{Ti}(\eta^5\text{-C}_5\text{R}_5)\text{Cl}_3$. Alkylation of the chloro complexes gave complexes of the type $\text{Ti}\{2,2'\text{-S}(\text{OC}_6\text{H}_2\text{-4-Me-6-}^t\text{Bu})_2\}(\eta^5\text{-C}_5\text{R}_5)\text{R}'$ ($\text{R}' = \text{Me, CH}_2\text{Ph}$). The crystal structure of the chloro complex $\text{Ti}\{2,2'\text{-S}(\text{OC}_6\text{H}_2\text{-4-Me-6-}^t\text{Bu})_2\}(\eta^5\text{-C}_5\text{H}_5)\text{Cl}$ has been determined by X-ray diffraction and shows a monomeric four-legged piano-stool structure with the sulfur and chlorine atoms in *trans* positions. A long titanium–sulfur bond length of 2.907(1) Å was observed. © 2001 Elsevier Science B.V. All rights reserved.

Keywords: Titanium complexes; Sulfur donor; Cyclopentadienyl ligand; Phenolato ligand

1. Introduction

Titanium(IV) half-sandwich complexes of the type $\text{Ti}(\eta^5\text{-C}_5\text{R}_5)\text{X}_3$ ($\text{X} = \text{one-electron ligand}$) can in principle accommodate up to three two-electron ligands L [1]. The full 18-electron configuration is not very frequently encountered, since various degrees of π -donation may occur to attenuate the Lewis-acidity [2]. Alkoxo and phenoxo ligands are particularly variable monoanionic ligands in this respect and have found application as controlling ligands in several instances [3]. We have been interested in the use of linked bis(phenolato)


ligands of the general type $2,2'\text{-Z}(\text{OC}_6\text{H}_2\text{-4-Me-6-}^t\text{Bu})_2$ where the link Z between the two bulky phenolato moieties is varied systematically [4]. In particular, the sulfide bridged system *tbmp* ($\text{Z} = \text{S}$) is important as ancillary ligand in olefin polymerization catalysis [5]. We report here the synthesis of several alkyl derivatives of the $\text{Ti}(\text{tbmp})$ fragment.

2. Results and discussion


As reported earlier [4b], the reaction between cyclopentadienyl lithium and the dichloro titanium complex with a sulfide-linked bis(phenolato) ligand, $[\text{Ti}(\text{tbmp})\text{Cl}_2]_2$ ($\text{tbmp} = 2,2'\text{-thiobis(6-tert-butyl-4-methylphenolato)}$), gives mono(cyclopentadienyl)-titanium complexes $\text{Ti}(\text{tbmp})(\eta^5\text{-C}_5\text{R}_5)\text{Cl}$ ($\text{R} = \text{H, 1; R} = \text{Me, 4}$), as shown in Eq. (1).

* Corresponding author. Tel.: +49-6131-3923737/25333; fax: +49-6131-3925606.

E-mail address: okuda@mail.uni-mainz.de (J. Okuda).



These complexes, together with the new derivative $\text{Ti}(\text{tbmp})\{\eta^5\text{-}1,3\text{-(SiMe}_3)_2\text{C}_5\text{H}_3\}\text{Cl}$ (**6**), are also accessible from the metathesis reaction between the appropriate trichloro titanium derivatives $\text{Ti}(\eta^5\text{-C}_5\text{R}_5)\text{Cl}_3$ ($\text{R} = \text{H, Me}$; $\text{R}_5 = 1,3\text{-(SiMe}_3)_2\text{H}_3$) and the dilithium or disodium salts of the tbmp ligand in approximately the same yields as via the previously reported method (Eq. (2)). These reactions can be carried out in separate steps or as a one-pot reaction with no significant differences in the yields. The chloro derivatives are obtained as red crystals which are soluble in ethers and slightly soluble in aliphatic hydrocarbons.



The ^1H -NMR spectra of the complexes **1**, **4** and **6** show one singlet for all the equivalent methyl groups of the *tert*-butyl substituents on the phenyl rings and one singlet for the two methyl substituents on the phenyl rings, while the protons of the phenyl ring at position 3 and 5 appear as two doublets ($^4J_{\text{HH}}$ ca. 2 Hz). The signals for the protons, methyl and trimethylsilyl substituents on the cyclopentadienyl rings are observed in the expected region. Based on similar mass spectral as well as ^1H - and ^{13}C -NMR spectroscopic data of **4** and **6** to those of the parent complex **1**, a structure analogous to that of crystallographically characterized **1** can be implied for both compounds¹. Crystals of **1** suitable for single-crystal X-ray diffraction were obtained by slow crystallization from a diethyl ether/hexane mixture. Fig. 1 shows an ORTEP plot of the

molecular structure of **1** and Table 1 summarizes selected bond distances and angles.

Complex **1** crystallized with diethyl ether in the unit cell. The coordination sphere around the titanium center in **1** can be described as that of a four-legged piano-stool in which the chlorine atom is located in *trans*-position to the sulfur atom. The angles O–Ti–S and O–Ti–Cl range from 70.42(5) to 95.08(6)° with a sum of 328.51°. The titanium–sulfur bond length of 2.907(1) Å is one of the largest so far reported in the literature [4b,6], although it is still smaller than the sum

of the van der Waals radii of the two atoms [7]. This may suggest a weaker interaction between the sulfur atom and the metal center than those reported for other titanium–thioether complexes adducts as $\text{Ti}(\eta^5\text{-C}_5\text{H}_5)(\text{SCH}_2\text{CH}_2\text{SCH}_2\text{CH}_2\text{S})$ (2.600(1) Å) [8a], $\text{TiCl}_4(\text{HSC}_6\text{H}_{11})_2$ (av. 2.664 Å) [8b], $[\text{Ti}(\eta^5\text{-C}_5\text{H}_5)(\text{SCH}_2\text{CH}_2\text{S})\text{Cl}]_2$ (av. 2.513 Å) [8c], $\text{Ti}(\text{N}^t\text{Bu})\text{Cl}_2(9\text{-aneS}_3)$ (av. 2.638 Å) [8d] and *cis*- $\text{TiCl}_4(\text{S}(\text{CH}_2)_4)_2$ (av. 2.265 Å) [8e]. The tbmp ligand

¹ The X-ray diffraction study on a single crystal of **6** shows a structure similar to that of **1** with a position for the chlorine atom *trans* to the sulfur atom. Red crystal, $0.13 \times 0.53 \times 0.63 \text{ mm}^3$, $a = 9.484(2)$, $b = 20.623(6)$, $c = 18.680(3)$ Å, $\alpha = \gamma = 90^\circ$, $\beta = 99.86(2)^\circ$, $V = 3600(1) \text{ Å}^3$, tentative space group $P2_1/c$. The poor quality of the crystallographic data preclude further discussion of the results.




Fig. 1. ORTEP diagram of the molecular structure of $\text{Ti}(\text{tbmp})(\eta^5\text{-C}_5\text{H}_5)\text{Cl}$ (**1**). Thermal ellipsoids are drawn at 50% probability level. Hydrogen atoms are omitted for the sake of clarity. The Et_2O solvent molecule is also excluded.

Table 1

Selected bond lengths (Å) and bond angles (°) for $\text{Ti}(\text{tbmp})(\eta^5\text{-C}_5\text{H}_5)\text{Cl}$ (**1**)

Bond lengths		Bond angles	
Ti–Cl	2.318(1)	Cl–Ti–O(1)	92.50(6)
Ti–S	2.907(1)	Cl–Ti–O(2)	95.08(6)
Ti–O(1)	1.827(2)	S–Ti–O(1)	70.42(5)
Ti–O(2)	1.827(2)	S–Ti–O(2)	70.51(6)
Ti–C(23)	2.379(3)	S–Ti–Cl	148.23(3)
Ti–C(24)	2.380(3)	O(1)–Ti–O(2)	115.59(8)
Ti–C(25)	2.373(3)	Ti–O(1)–C(1)	141.9(1)
Ti–C(26)	2.349(3)	Ti–O(2)–C(13)	137.8(1)
Ti–C(27)	2.354(4)	C(2)–S–C(12)	107.3 (1)
S–C(2)	1.769(2)	Ti–S–C(2)	91.93(7)
S–C(12)	1.773(2)	Ti–S–C(12)	89.93(8)
O(1)–C(1)	1.344(2)		
O(2)–C(13)	1.349(3)		

adopts the typical boat conformation observed in other complexes [4b,6] with the sulfur atom pointing towards the metal center in the proximity of the cyclopentadienyl ring (see Fig. 2). This is probably due to the electron donating effect of the cyclopentadienyl ligand and the steric hindrance of the *tert*-butyl groups. A similar configuration is adopted by other related linked bis(phenolato) ligands with a sulfinyl [4d], oxo [9], tellurium [10] and methylene bridge [11]. The metal–oxygen bond lengths of 1.827(2) Å are in the expected range for titanium complexes with a linked bis(phenolato) ligand (av. 1.892 Å in $[\text{Ti}\{2,2'\text{-S}(\text{OC}_6\text{H}_2\text{-4-Me-6-}^t\text{Bu})_2\}(\text{O}^i\text{Pr})_2]_2$ and av. 1.855 Å in $\text{Ti}\{2,2'\text{-S}(\text{OC}_6\text{H}_2\text{-4-Me-6-}^t\text{Bu})_2\}(\text{o-CH}_2\text{NMe}_2\text{C}_6\text{H}_4)\text{Cl}$ [4b], av. 1.888 Å in $\text{Ti}\{2,2'\text{-S}(\text{OC}_6\text{H}_2\text{-4-Me-6-}^t\text{Bu})_2\}\text{Br}_2$ [6], av. 1.872 Å in $\text{Ti}\{2,2'\text{-Te}(\text{OC}_6\text{H}_2\text{-4-Me-6-}^t\text{Bu})_2\}(\eta^5\text{-C}_5\text{Me}_5)\text{Cl}$ [10], av.





Fig. 2. View of the eight-membered ring conformation in $\text{Ti}(\text{tbmp})(\eta^5\text{-C}_5\text{H}_5)\text{Cl}$ (**1**).

1.751 Å in $\text{Ti}\{2,2'\text{-CH}_2(\text{OC}_6\text{H}_2\text{-4-Me-6-}^t\text{Bu})_2\}\text{Cl}_2$ [11a], av. 1.776 Å in $\text{Ti}\{2,2'\text{-CH}_2(\text{OC}_6\text{H}_2\text{-4-Me-6-}^t\text{Bu})_2\}(\text{BH}_4)_2$ [11b], av. 1.788 Å in $\text{Ti}\{2,2'\text{-CH}_2(\text{OC}_6\text{H}_2\text{-4-Me-6-}^t\text{Bu})_2\}(\text{o-CH}_2\text{NMe}_2\text{C}_6\text{H}_4)(\text{OSO}_2\text{CF}_3)$ [11d], and av. 1.81 Å in $\text{Ti}\{2,2'\text{-(OC}_6\text{H}_2\text{-4-OMe-6-}^t\text{Bu})_2\}(\text{CH}_2\text{Ph})_2$ [12]. Values larger than those found for titanium complexes with a non-linked alkoxo ligand as $\text{Ti}(\text{OC}_6\text{H}_3\text{-2,6-}^t\text{Bu}_2)\text{Cl}_3(\text{THF})_2$ (1.807(8) Å) [9], $\text{Ti}(\text{OPh})_2\text{Cl}_2$ (1.744(10) Å) [13a], $\text{Ti}(\text{OC}_6\text{H}_3\text{-2,6-Ph}_2)_2\text{Cl}_2$ (1.726(2) Å) [13b], $[\text{Ti}(\text{OC}_6\text{H}_3\text{-2,6-Ph}_2)_2]_2(\eta\text{-Cl})_2$ (av. 1.817 Å) [13c], $\text{Ti}(\text{OC}_6\text{H}_3\text{-2,6-Me}_2)_2\text{Cl}_2(\text{THF})_2$ (av. 1.788 Å) [13d], and $\text{Ti}(\text{OC}_6\text{H}_3\text{-2,6-}^i\text{Pr}_2)(\eta^5\text{-C}_5\text{H}_5)(\text{SPh})\text{Cl}$ (1.80(2) Å) [13e] may suggest a decrease in the π -donation from the oxygen atoms to the metal center [14].

Metathesis reactions of the chloro complexes **1**, **4** and **6** with alkylating reagents MgMeCl or $\text{Mg}(\text{CH}_2\text{Ph})\text{Cl}$ resulted in the formation of alkyl complexes $\text{Ti}(\text{tbmp})(\eta^5\text{-C}_5\text{R}_5)\text{R}'$ ($\text{R} = \text{H}$, $\text{R}' = \text{Me}$ (**2**), CH_2Ph (**3**); $\text{R} = \text{Me}$, $\text{R}' = \text{Me}$ (**5**); $\text{R}_5 = 1,3\text{-(SiMe}_3)_2\text{H}_3$, $\text{R}' = \text{Me}$ (**7**), CH_2Ph (**8**)) in moderate yields (68–75%) (Eq. (3)). The complexes **2** and **5** are orange, the complexes **3** and **7** are dark yellow and the complex **8** is brown. They are all soluble in aliphatic hydrocarbons. The reaction of the chloro complex **4** with $\text{Mg}(\text{CH}_2\text{Ph})\text{Cl}$ in THF at -78°C results in the formation of a yellow reaction mixture, but upon work-up the starting material was recovered.

For the methyl complexes **2** and **7**, the methyl group at the titanium center are observed as singlets at 1.34 and 1.31 ppm, respectively, in the $^1\text{H-NMR}$ spectra, while for complex **5** the corresponding signal is high-field shifted to 0.38 ppm. The $^{13}\text{C-NMR}$ spectroscopic TiCH_3 resonances are observed at 58.1 for **2**, 60.8 for **5**, and 60.5 ppm for **7**. The benzyl derivatives **3** and **8** show only one resonance for the two equivalent methylene protons of the benzyl groups at 3.15 ppm and three

broad signals for the protons of the benzylic phenyl ring. The ^1H -NMR spectra of the alkyl complexes **2**, **3**, **5**, **7** and **8** show patterns for the tbmp ligand similar to those described for the ^1H -NMR spectra of the chloro precursors **1**, **4** and **6**: The methyl groups of the *tert*-butyl substituents on the phenolato groups remain equivalent and appear in the region from 1.26 to 1.41 ppm while the methyl substituents on the phenyl rings appear as a singlet at 2.2 ppm. The proton atoms on the phenyl ring are recorded as doublets with a similar coupling constant as in the chloro precursors.



This work shows that the $\text{Ti}(\text{tbmp})$ fragment is stabilized by a cyclopentadienyl ligand and alkylation of the remaining chloro function becomes feasible. Previously, we had encountered considerable difficulties in synthesizing the ten-electron dialkyls of the type $\text{Ti}(\text{tbmp})\text{R}_2$ by alkylating the dichloro complex $\text{Ti}(\text{tbmp})\text{Cl}_2$. Such problems do not appear in the preparation of the more stable derivatives $\text{Ti}(\text{tbmp})(\eta^5\text{-C}_5\text{R}_5)\text{R}'$ which formally have a 14-valence electron configuration.

3. Experimental

All operations were performed under an inert atmosphere of argon using standard Schlenk-line or glove-box techniques. Diethyl ether was distilled from sodium benzophenone ketyl. Pentane and hexane were purified by distillation over sodium–benzophenone ketyl–triglyme. tbmpH_2 [15], $\text{Ti}(\eta^5\text{-C}_5\text{H}_5)\text{Cl}_3$ [16], $\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\text{Cl}_3$ [17] and $\text{Ti}\{\eta^5\text{-}1,3-(\text{SiMe}_3)_2\text{C}_5\text{H}_3\}\text{Cl}_3$ [18] were synthesized according to published procedures. Li^nBu (Aldrich), NaH (Aldrich), MgMeCl (Aldrich) and $\text{Mg}(\text{CH}_2\text{C}_6\text{H}_5)\text{Cl}$ (Aldrich) were used as received. Lithium [19] and Grignard [20] solutions were titrated prior to use. NMR spectra were recorded on a Bruker DRX 400 or on a Varian Unity 300 spectrometer (^1H , 400.1 MHz, 299.9 MHz; ^{13}C , 75.43, 100.6 MHz) in CDCl_3 at 298 K. Chemical shifts (in ppm) for ^1H and ^{13}C were referenced internally using the residual solvent

resonances and reported relative to tetramethylsilane. Mass spectra were recorded on a Finnigan 8230 spectrometer. Elemental analyses were performed by the microanalytical laboratory of this department.

3.1. $\text{Ti}(\text{tbmp})(\eta^5\text{-C}_5\text{H}_5)\text{Cl}$ (**1**)

A solution of Li^nBu (2.1 ml of a 1.5 M solution in THF) was added to a solution of tbmpH_2 (1.1 g, 3.0 mmol) in hexane (ca. 25 ml) at 0°C and the mixture was

stirred for 16 h. The solvents were filtered off and the white precipitate was washed with hexane (2×10 ml) and dried in vacuo. A solution of $\text{Ti}(\eta^5\text{-C}_5\text{H}_5)\text{Cl}_3$ (0.50 g, 2.28 mmol) in diethyl ether (20 ml) was added to a suspension of $\text{Li}_2(\text{tbmp})$ (0.90 g, 2.4 mmol) in diethyl ether (15 ml) at -78°C . The mixture was slowly warmed up to room temperature (r.t.) and stirred for 16 h. Concentration of the solution followed by addition of hexane (ca. 20 ml) and cooling to -30°C afford 0.92 g of dark red crystals (80%): ^1H -NMR δ 1.25 (s, 18H, 6- $\text{C}(\text{CH}_3)_3$), 2.21 (s, 6H, 4- CH_3), 6.76 (s, 5H, C_5H_5), 6.98 (d, $^4J_{\text{HH}} = 1.6$ Hz, 2H, 5-H), 7.27 (d, $^4J_{\text{HH}} = 1.6$ Hz, 2H, 3-H); $^{13}\text{C}\{^1\text{H}\}$ -NMR δ 20.8 (4- CH_3), 29.8 (6- $\text{C}(\text{CH}_3)_3$), 35.2 (6- $\text{C}(\text{CH}_3)_3$), 119.4 (C_5H_5), 128.3 (C-2), 129.9 (C-5), 130.4 (C-4), 131.6 (C-3), 138.4 (C-6), 167.1 (C-1); EI MS m/z 504 (86, $[\text{M}^+]$), 439 (100, $[\text{M}^+ - \text{C}_5\text{H}_5]$); Anal. Calc. for $\text{C}_{27}\text{H}_{33}\text{ClO}_2\text{STi}$: C, 64.22; H, 6.59. Found: C, 64.35; H, 6.49.

3.2. $\text{Ti}(\text{tbmp})(\eta^5\text{-C}_5\text{H}_5)\text{Me}$ (**2**)

Methylmagnesium chloride (0.23 ml of a 3 M solution in THF) was added to a precooled ethereal solution of **1** (0.33 g, 0.66 mmol) at -78°C (ca. 10 ml) and the reaction mixture was allowed to warm slowly to r.t. overnight. All volatiles were removed in vacuo and the residue was extracted with pentane (ca. 30 ml) and the extracts concentrated. Cooling the solution to -78°C

(3)

overnight precipitated 0.22 g of an orange solid (68%): $^1\text{H-NMR}$ δ 1.32 (s, 18H, 6-C(CH₃)₃), 1.34 (s, TiCH₃), 2.29 (s, 6H, 4-CH₃), 6.56 (s, 5H, C₅H₅), 7.06 (d, $^4J_{\text{HH}} = 2$ Hz, 2H, 5-H), 7.44 (d, $^4J_{\text{HH}} = 2$ Hz, 2H, 3-H); $^{13}\text{C}\{^1\text{H}\}\text{-NMR}$ δ 20.7 (4-CH₃), 29.7 (6-C(CH₃)₃), 35.1 (6-C(CH₃)₃), 58.1 (TiCH₃), 115.4 (C₅H₅), 128.4 (C-2), 128.9 (C-4), 129.4 (C-5), 132.9 (C-3), 137.6 (C-6), 166.1 (C-1); EI MS m/z 469 (100, $\text{M}^+ - \text{CH}_3$); Anal. Calc. for C₂₈H₃₆O₂STi: C, 69.41; H, 7.49. Found: C, 69.34; H, 7.72.

3.3. *Ti(tbmp)($\eta^5\text{-C}_5\text{H}_5$)(CH₂Ph) (3)*

A solution of **1** (0.40 g, 0.79 mmol) in diethyl ether (ca. 15 ml) was treated with benzylmagnesium chloride (0.77 ml of a 1.24 M solution in THF) at -78°C . After stirring for 16 h at r.t., all volatiles were removed in vacuo. The residue was extracted into hexane (ca. 30 ml) and the solvent was removed. Recrystallization from pentane at -30°C afforded dark yellow crystals (0.30 g, 68%): $^1\text{H-NMR}$ δ 1.41 (s, 18H, 6-C(CH₃)₃), 2.26 (s, 6H, 4-CH₃), 3.14 (s, 2H, TiCH₂), 6.31 (s, 5H, C₅H₅), 7.06 (d, $^4J_{\text{HH}} = 2$ Hz, 2H, 5-H), 7.26, 7.28, 7.30 (br, 5H, C₆H₅), 7.40 (d, $^4J_{\text{HH}} = 2$ Hz, 2H, 3-H); $^{13}\text{C}\{^1\text{H}\}\text{-NMR}$ δ 20.7 (4-CH₃), 29.9 (6-C(CH₃)₃), 35.4 (6-C(CH₃)₃), 81.2 (TiCH₂), 117.9 (C₅H₅), 122.6, 126.4 (C₆H₅), 128.0 (C-5), 128.9 (C-2), 129.2 (C-4), 129.6 (C₆H₅), 133.2 (C-3), 137.9 (C-6), 152.4 (*ipso* C₆H₅), 166.7 (C-1); Anal. Calc. for C₃₄H₄₀O₂STi: C, 72.87; H, 7.14. Found: C, 72.96; H, 7.18.

3.4. *Ti(tbmp)($\eta^5\text{-C}_5\text{Me}_5$)Cl (4)*

This compound was prepared following a procedure analogous to that described to prepare **1** from Li₂(tbmp) (0.50 g, 1.35 mmol) and Ti($\eta^5\text{-C}_5\text{Me}_5$)Cl₃ (0.37 g, 1.30 mmol) and isolated as red crystals; yield 0.58 g (79%): $^1\text{H-NMR}$ δ 1.35 (s, 18H, 6-C(CH₃)₃), 2.25 (s, 15H, C₅(CH₃)₅), 2.26 (s, 6H, 4-CH₃), 7.05 (d, $^4J_{\text{HH}} = 2$ Hz, 2H, 5-H), 7.23 (d, $^4J_{\text{HH}} = 2$ Hz, 2H, 3-H); $^{13}\text{C}\{^1\text{H}\}\text{-NMR}$ δ 13.0 (C₅(CH₃)₅), 20.9 (4-CH₃), 31.8 (6-C(CH₃)₃), 34.8 (6-C(CH₃)₃), 126.5 (C-5), 128.0 (C-3), 129.5 (C₅(CH₃)₅), 130.1 (C-4), 131.9 (C-2), 137.7 (C-6), 161.0 (C-1); EI MS m/z 574 (89, $[\text{M}^+]$), 539 (41, $[\text{M}^+ - \text{Cl}]$), 439 (100, $[\text{M}^+ - \text{C}_5\text{Me}_5]$); Anal. Calc. for C₃₂H₄₃ClO₂STi: C, 66.83; H, 7.54. Found: C, 67.05; H, 7.64.

3.5. *Ti(tbmp)($\eta^5\text{-C}_5\text{Me}_5$)Me (5)*

Following a procedure analogous to that described to prepare **2**, the chloro complex **4** (0.21 g, 0.36 mmol) was reacted with methylmagnesium chloride (0.13 ml of a 3 M solution in THF) to give orange crystals; yield 0.15 g (75%): $^1\text{H-NMR}$ δ 0.86 (s, 3H, TiCH₃), 1.27 (s, 18H, 6-C(CH₃)₃), 2.07 (s, 15H, C₅(CH₃)₅), 2.24 (s, 6H,

4-CH₃), 7.01 (d, $^4J_{\text{HH}} = 2$ Hz, 2H, 5-H), 7.34 (d, $^4J_{\text{HH}} = 2$ Hz, 2H, 3-H); $^{13}\text{C}\{^1\text{H}\}\text{-NMR}$ δ 12.5 (C₅(CH₃)₅), 20.7 (4-CH₃), 30.0 (6-C(CH₃)₃), 34.7 (6-C(CH₃)₃), 60.8 (TiCH₃), 124.7 (C₅(CH₃)₅), 125.3 (C-5), 127.9 (C-2), 129.7 (C-4), 131.9 (C-3), 137.7 (C-6), 165.5 (C-1); EI MS m/z 537 (100, $[\text{M}^+ - \text{CH}_3]$); Anal. Calc. for C₃₃H₄₆O₂STi: C, 71.49; H, 8.30. Found: C, 71.74; H, 8.47.

3.6. *Ti(tbmp){ $\eta^5\text{-1,3-(SiMe}_3)_2\text{C}_5\text{H}_3$ }Cl (6)*

A solution of tbmpH₂ (0.87 g, 2.44 mmol) in hexane (ca. 25 ml) was added to a suspension of NaH (0.12 g, 4.88 mmol) in hexane (ca. 20 ml) and the mixture was stirred for 16 h. A solution of Ti($\eta^5\text{-1,3-(SiMe}_3)_2\text{C}_5\text{H}_3$)Cl₃ (0.89 g, 2.44 mmol) in diethyl ether (ca. 60 ml) was added to the mixture and stirred for 4 h. The resulting solution was filtered and concentrated to ca. 10 ml and cooled to -20°C overnight to afford 0.91 g of a dark red solid. From the mother liquor a second crop was obtained (0.49 g); combined yield 1.40 g (81%): $^1\text{H-NMR}$ δ 0.24 (s, 18H, Si(CH₃)₃), 1.27 (s, 18H, 6-C(CH₃)₃), 2.24 (s, 6H, 4-CH₃), 6.99 (br, 2H, 5-H) overlapping 7.01 (br, 1H, 7-H), 7.24 (br, 2H, 9,10-H) overlapping 7.25 (br, 2H, 3-H); $^{13}\text{C}\{^1\text{H}\}\text{-NMR}$ δ -0.2 (Si(CH₃)₃), 20.7 (4-CH₃), 30.2 (6-C(CH₃)₃), 35.2 (6-C(CH₃)₃), 128.1 (C-2), 128.5 (C-5), 129.9 (C-4), 130.0 (C-9,10), 131.0 (C-7), 132.4 (C-3), 137.6 (C-8,11), 138.1 (C-6), 166.5 (C-1); EI MS m/z 648 (100, $[\text{M}^+]$), 439 (96, $[\text{M}^+ - (\text{SiMe}_3)_2\text{C}_5\text{H}_3]$); Anal. Calc. for C₃₃H₄₉ClO₂SSi₂Ti: C, 61.05; H, 7.61. Found: C, 61.21; H, 7.57.

3.7. *Ti(tbmp){ $\eta^5\text{-1,3-(SiMe}_3)_2\text{C}_5\text{H}_3$ }Me (7)*

Following a procedure analogous to that described to prepare **2**, the chloro complex **6** (0.35 g, 0.54 mmol) and methylmagnesium chloride (0.20 ml of a 3 M solution in THF) were reacted to give yellow–red micro-crystals; yield 0.25 g (75%): $^1\text{H-NMR}$ δ 0.18 (s, 18H, Si(CH₃)₃), 1.26 (s, 18H, 6-C(CH₃)₃), 1.31 (s, 3H, TiCH₃), 2.23 (s, 6H, 4-CH₃), 6.68 (t, $^4J_{\text{HH}} = 2$ Hz, 1H, 7-H), 6.76 (d, $^4J_{\text{HH}} = 2$ Hz, 2H, 9,10-H), 6.99 (d, $^4J_{\text{HH}} = 2$ Hz, 2H, 5-H), 7.33 (d, $^4J_{\text{HH}} = 2$ Hz, 2H, 3-H); $^{13}\text{C}\{^1\text{H}\}\text{-NMR}$ δ -0.1 (Si(CH₃)₃), 20.7 (4-CH₃), 30.2 (6-C(CH₃)₃), 35.1 (6-C(CH₃)₃), 60.5 (TiCH₃), 125.4 (C-9,10), 127.6 (C-2), 128.2 (C-5), 128.5 (C-4), 129.6 (C-8,11), 131.5 (C-7), 132.3 (C-3), 137.3 (C-6), 166.8 (C-1); Anal. Calc. for C₃₄H₅₂O₂SSi₂Ti: C, 64.98; H, 8.28. Found: C, 64.94; H, 8.59.

3.8. *Ti(tbmp){ $\eta^5\text{-1,3-(SiMe}_3)_2\text{C}_5\text{H}_3$ }CH₂Ph (8)*

Following a procedure analogous to that described to prepare **3**, the chloro complex **6** (0.35 g, 0.54 mmol) was reacted with benzylmagnesium chloride (0.45 ml of

a 1.24 M solution in THF) to give a yellow–brown solid; yield 0.38 g (70%): $^1\text{H-NMR}$ δ –0.06 (s, 18H, $\text{Si}(\text{CH}_3)_3$), 1.32 (s, 18H, 6- $\text{C}(\text{CH}_3)_3$), 2.18 (s, 6H, 4- CH_3), 3.15 (s, 2H, TiCH_2), 6.47 (d, $^4J_{\text{HH}} = 2$ Hz, 2H, 5-H), 6.96(d, $^4J_{\text{HH}} = 2$ Hz, 2H, 3-H), 7.02 (t, $^4J_{\text{HH}} = 2$ Hz, 1H, 7-H), 7.20, 7.24 (br, 2H, C_6H_5), 7.26 (br, 2H, 9,10-H), 7.27 (br, 2H, C_6H_5); $^{13}\text{C}\{^1\text{H}\}$ -NMR δ –0.2 ($\text{Si}(\text{CH}_3)_3$), 20.8 (4- CH_3), 30.2 (6- $\text{C}(\text{CH}_3)_3$), 35.2 (6- $\text{C}(\text{CH}_3)_3$), 82.6 (TiCH_2), 122.6 (C_6H_5), 125.9 (C-8,11), 126.5, 127.9 (C_6H_5), 128.3 (C-2), 128.5 (C-5), 128.9 (C-4), 129.2 (C-9,10), 131.0 (C-7), 133.4 (C-3), 138.1 (C-6), 153.1 (*ipso* C_6H_5), 166.0 (C-1); EI MS m/z 613 (100, $[\text{M}^+ - \text{C}_7\text{H}_7]$); Anal. Calc. for $\text{C}_{40}\text{H}_{56}\text{O}_2\text{SSi}_2\text{Ti}$: C, 68.20; H, 7.95. Found: C, 67.90; H, 7.80.

3.9. X-ray crystal structure analysis of **1**

Crystallographic data are summarized in Table 2. Single crystals of **1** were grown from a diethyl ether/hexane solution at -30°C . Data were collected using an ENRAF–Nonius CAD4 diffractometer (λ -scan mode, Mo– K_α radiation, $\lambda = 0.7107$ Å). Data correction for Lorentz polarization and absorption (using Ψ -scans) was carried out using the program system MOLEN [21a]. From 7400 measured reflections, 6993 were independent. The structure was solved using direct

methods (SHELXS-86) [21b] and difference Fourier syntheses and refined by full-matrix least-squares against all F_o^2 data (SHELXL-93) [21c] (473 parameters). The non-hydrogen atoms of the complex were refined with anisotropic displacement parameters and the hydrogen atoms were refined with isotropic displacement parameters. The crystal structure shows diethyl ether included around a centre of inversion. Split positions of the severely disordered molecule could be resolved and refined with isotropic displacement parameters. Hydrogen atoms of this molecule were not included in the structure factor calculation. The refinement converged with $R = 0.0448$, $wR_2 = 0.1110$ for all observed F_o data, goodness-of-fit 1.078.

4. Supplementary material

Crystallographic data for the structural analysis have been deposited with the Cambridge Crystallographic Data Centre, CCDC No. 154412. Copies of this information may be obtained free of charge from The Director, CCDC, 12 Union Road, Cambridge CB2 1EZ, UK (fax: +44-1223-336033; e-mail: deposit@ccdc.cam.ac.uk or www: <http://www.ccdc.cam.ac.uk>).

Acknowledgements

We gratefully acknowledge the financial support by the Fonds der Chemischen Industrie.

References

- [1] M.L.H. Green, J. Organomet. Chem. 500 (1995) 127.
- [2] R. Poli, Chem. Rev. 91 (1991) 509.
- [3] (a) M.G. Thorn, Z.C. Etheridge, P.E. Fanwick, I.P. Rothwell, J. Organomet. Chem. 591 (1999) 148. (b) A.E. Firth, J.C. Stewart, A.J. Hoskin, D.W. Stephan, J. Organomet. Chem. 591 (1999) 185. (c) K. Nomura, T. Komatsu, Y. Imanishi, J. Mol. Catal. A Chem. 152 (2000) 249. (d) A.J. Nielson, P. Schwerdtfeger, J.M. Waters, J. Chem. Soc. Dalton Trans. (2000) 529.
- [4] (a) J. Okuda, S. Fokken, H.-C. Kang, W. Massa, Chem. Ber. 128 (1995) 221. (b) S. Fokken, T.P. Spaniol, H.-C. Kang, W. Massa, J. Okuda, Organometallics 15 (1996) 5069. (c) S. Fokken, T.P. Spaniol, J. Okuda, F.G. Sernetz, R. Mülhaupt, Organometallics 16 (1997) 4240. (d) J. Okuda, S. Fokken, H.-C. Kang, W. Massa, Polyhedron 17 (1998) 943. (e) J. Okuda, S. Fokken, T. Kleinhenn, T.P. Spaniol, Eur. J. Inorg. Chem. (2000) 1321.
- [5] (a) T. Miyatake, K. Mizunuma, M. Kakugo, Macromol. Chem. Macromol. Symp. 66 (1993) 203. (b) F.G. Sernetz, R. Mülhaupt, S. Fokken, J. Okuda, Macromolecules 30 (1997) 1562. (c) R.D.J. Froese, D.G. Musaev, T. Matsubara, K. Morokuma, J. Am. Chem. Soc. 119 (1997) 7190. (d) R.D.J. Froese, D.G. Musaev, K. Morokuma, Organometallics 18 (1999) 373.
- [6] L. Porri, A. Ripa, P. Colombo, E. Miano, S. Capelli, S.V. Meille, J. Organomet. Chem. 514 (1996) 213.

Table 2

Crystallographic data for $\text{Ti}(\text{tbmp})(\eta^5\text{-C}_5\text{H}_5)\text{Cl}$ (**1**)

Empirical formula	$\text{C}_{27}\text{H}_{33}\text{ClO}_2\text{STi} \cdot 0.5\text{C}_4\text{H}_{10}\text{O}$
f_w	542.00
Crystal colour	Red
Crystal size (mm)	$0.51 \times 0.51 \times 0.83$
Crystal system	Triclinic
Space group	$P\bar{1}$ (no. 2)
a (Å)	9.715(3)
b (Å)	12.834(3)
c (Å)	12.951(4)
α (°)	67.95(2)
β (°)	83.73(2)
γ (°)	76.20(2)
V (Å ³)	1453.1(7)
Z	2
D_{calc} (mg m ^{−3})	1.239
Absorption coefficient (mm ^{−1})	0.483
$F(000)$	574
Temperature (K)	296(2)
θ range (°)	3 to 28
Index ranges	$0 \leq h \leq 12$; $-16 \leq k \leq 16$; $-17 \leq l \leq 17$
Reflections measured	7400
Independent reflections	6993 ($R_{\text{int}} = 0.0197$)
Observed reflections	5280
Final R indices R_1 , wR_2	0.0448, 0.1110
[$I \geq 2\sigma(I)$]	
Final R indices R_1 , wR_2 (all data)	0.0670, 0.1330
Goodness-of-fit	1.078
Largest e_{max} , e_{min} (e Å ^{−3})	0.348 and −0.273

- [7] L. Pauling, *The Nature of the Chemical Bond*, 3rd edn., Cornell University Press, Ithaca, NY, 1960.
- [8] (a) T.T. Nadasdi, Y. Huang, D.W. Stephan, *Inorg. Chem.* 32 (1993) 347. (b) C.H. Winter, T.S. Lewkebandara, J.W. Proscia, A.L. Rheingold, *Inorg. Chem.* 32 (1993) 3807. (c) T.T. Nadasdi, D.W. Stephan, *Inorg. Chem.* 32 (1993) 5933. (d) P.J. Wilson, A.J. Blake, P. Mountford, M. Schröder, *J. Chem. Soc., Chem. Commun.* (1998) 1007. (e) T.S. Lewkebandara, P.J. McKarns, B.S. Haggerty, G.P.A. Yap, A.L. Rheingold, C.H. Winter, *Polyhedron* 17 (1998) 1.
- [9] M. Mazzanti, C. Floriani, A. Chiesi-Villa, C. Guastini, *J. Chem. Soc. Dalton Trans.* (1989) 1793.
- [10] Y. Nakayama, K. Watanabe, N. Ueyama, A. Nakamura, A. Harada, J. Okuda, *Organometallics* 19 (2000) 2498.
- [11] (a) C. Floriani, F. Corazza, W. Lesueur, A. Chiesi-Villa, C. Guastini, *Angew. Chem.* 101 (1989) 93; *Angew. Chem. Int. Ed. Engl.* 28 (1989) 66. (b) F. Corazza, C. Floriani, A. Chiesi-Villa, C. Guastini, *Inorg. Chem.* 30 (1991) 145. (c) P.J. Toscano, E.J. Schermerhorn, C. Dettelbacher, D. Macherone, J. Zubieta, *J. Chem. Soc. Chem. Commun.* (1991) 933. (d) E.E.C.G. Gielens, T.W. Dijkstra, P. Berno, A. Meetsma, B. Hessen, J.H. Teuben, *J. Organomet. Chem.* 591 (1999) 88. (e) D.R. Mulford, P.E. Fanwick, I.P. Rothwell, *Polyhedron* 19 (2000) 35.
- [12] A.J. van der Linden, C.J. Schaverien, N. Meijboom, C. Ganter, A.G. Orpen, *J. Am. Chem. Soc.* 117 (1995) 3008.
- [13] (a) K. Watenpaugh, C.N. Caughlan, *Inorg. Chem.* 5 (1966) 1782. (b) J.R. Dilworth, J. Hanich, M. Krestel, J. Beck, J. Strahle, *J. Organomet. Chem.* 315 (1986) C9. (c) J.E. Hill, J.M. Nash, P.E. Fanwick, I.P. Rothwell, *Polyhedron* 9 (1990) 1617. (d) N. Kanehisa, Y. Kai, N. Kasai, H. Yasuda, Y. Nakayama, K. Takei, A. Nakamura, *Chem. Lett.* (1990) 2167. (e) A.V. Firth, D.W. Stephan, *Organometallics* 16 (1997) 2183.
- [14] P.T. Wolczaski, *Polyhedron* 14 (1995) 3335.
- [15] T.K. Praskasha, R.O. Day, R.R. Holmes, *J. Am. Chem. Soc.* 115 (1993) 2690.
- [16] A.M. Cardoso, R.J.H. Clark, S. Moorhouse, *J. Chem. Soc. Dalton Trans.* (1980) 1156.
- [17] M. Mena, M.A. Pellinghelli, P. Royo, R. Serrano, A. Tiripichio, *J. Chem. Soc. Dalton Trans.* (1996) 1118.
- [18] (a) P. Jutzi, M. Kuhn, *J. Organomet. Chem.* 173 (1979) 221. (b) C.H. Winter, X.-X. Zhou, D.A. Dobbs, M.J. Heeg, *Organometallics* 10 (1991) 210.
- [19] J. Suffert, *J. Org. Chem.* 54 (1989) 509.
- [20] S.C. Watson, J.F. Eastham, *J. Organomet. Chem.* 9 (1967) 165.
- [21] (a) C.K. Fair, MOLEN, *An Interactive Structure Solution Procedure*, Enraf-Nonius, Delft, 1990. (b) G.M. Sheldrick, SHELXS-86, *Program for the Solution of Crystal Structures*, University of Göttingen, Germany, 1986. (c) G.M. Sheldrick, SHELXL-93, *Program for the Refinement of Crystal Structures*, University of Göttingen, Germany, 1993.