## PHOTOCHEMICAL SYNTHESIS OF TRICYCLIC 1H-1,2-DIAZEPINES

Yoshiro YAMASHITA and Mituo MASUMURA\* Department of Applied Chemistry, Faculty of Engineering, Tokushima University, Minami-Josanjima, Tokushima 770

The photochemical reaction of tricyclic N-iminopyridinium ylides afforded lH-l,2-diazepines containing novel structures by the selective electrocyclic ring closure followed by ring opening.

The photochemical reaction of N-iminopyridinium ylides to give 1H-1,2-diazepines has been extensively studied.<sup>1</sup> This reaction is sensitive to the substituent effects and the successful examples are limited to the ylides possessing an electron-withdrawning chromophore at the imino position<sup>1,2</sup> except for N-iminoquinolinium dimers.<sup>3</sup> Recently we reported the first example of the stable tricyclic N-imino-pyridinium ylides possessing a non-conjugated substituent at the imino position.<sup>4</sup> As a part of detailed investigation of these novel ylides, we have examined their photochemical behaviors.

When a methanol solution of la was irradiated with a 400W-high pressure mercury lamp under nitrogen atmosphere, tricyclic diazepine 2a (mp 103-107°C, 62%) was obtained as the only isolable product.<sup>5</sup> The structure of 2a was deduced on the basis of the spectroscopic data along with chemical evidence. 2a;  $\lambda$ max (EtOH) 293 nm (log $\varepsilon$ =4.04),  $\nu$ max (KBr) 1700 cm<sup>-1</sup>,  $\delta$ ppm (CDCl<sub>3</sub>) 0.88 (s, 3H), 2.19 (s, 3H), 5.03 (dd, Hd, J=6.0, 1.5 Hz), 5.40 (ddd, Hb, J=11.0, 4.0, 1.5), 5.81 (dd, Hc, J=11.0, 6.0), 6.15 (d, Ha, J=4.0), 6.9-7.5 (m, 10H). The fact that the coupling constant J<sub>bc</sub> is distinctly larger than J<sub>ab</sub> and J<sub>cd</sub> is inconsistent with the structure 3a,



4

but can be reasonably accomodated by the structure 2a. Reaction of 2a with 2,5dimethyl-3,4-diphenylcyclopentadienone in refluxing toluene produced a 1:1 adduct 4 (mp 193-194°C, 40%).<sup>6</sup> On the other hand, heating of 2a in chloroform at 120°C in a sealed tube for 1 h afforded the starting ylide 1a in 80% yield. Irradiation of 1b and 1c under the same conditions as that of 1a also afforded the diazepines 2b (mp 201-205°C, 18%) and 2c (oil, ca 10%), respectively, although the reactions are slower and the yields are worse than that of 1a.<sup>7</sup> The formation of the diazepines 2 can be explained by ring opening from the assumed primary photoproducts 5 formed by the selective electrocyclic ring closure of the pyridinium ylides. The selectivity observed here is different from the reported examples in which the ring closure on the less hindered  $\alpha$ -carbon is favored.<sup>1</sup> This may be attributed to the fused five-membered ring which ring formed in the electrocyclic ring closure and may also affect the ring opening.

## REFERENCES AND NOTES

- J. Streith, J. P. Luttringer, and M. Nastasi, J. Org. Chem., <u>36</u>, 2962 (1971);
  T. Sasaki, K. Kanematsu, A. Kakehi, I. Ichikawa, and K. Hayakawa, ibid., <u>35</u>, 426 (1970), A. Balasubramanian, J. M. McIntosh, and V. Snieckus, ibid., <u>35</u>, 433 (1970); R. A. Abramovitch and T. Takaya, ibid., <u>38</u>, 3311 (1973).
- 2. V. Snieckus and G. Kan, Chem. Commun. 172 (1970).
- 3. T. Tsuchiya, J. Kurita, H. Igeta, and V. Snieckus, ibid., 640 (1974).
- 4. Y. Yamashita and M. Masumura, Tetrahedron Lett., <u>1979</u>, 1765. The possibility of the ylide <u>7</u>, the corresponding regioisomers of the ylide <u>1</u>, instead of <u>1</u> could be denied by the inspection of the mass fragmentation pattern in addition to the reported reason, because the fragments due to loss of benzonitrile were observed as follows.

la; m/e 352 (50%, M<sup>+</sup>), 208 (100%), 105 (51%, 8a-PhCN), 104 (44%). lb; m/e 366 (48%, M<sup>+</sup>), 222 (100%), 119 (27%, 8b-PhCN), 118 (34%).  $\chi_{0}^{1}$ 

- 5. Satisfactory elemental analyses were obtained for all new compounds reported in this paper.
- 6. 4; vmax (KBr) 1705, 1780 cm<sup>-1</sup>, &ppm (CDCl<sub>3</sub>) 1.10 (s, 3H), 1.28 (s, 3H), 1.49 (s, 3H), 1.75 (s, 3H), 3.04 (brs, 2H), 5.10 (brs, 1H), 6.4-7.3 (m, 19H), 7.5-7.7 (m, 2H). A 1H-1,2-diazepine is known to react with 2,5-dimethyl-3,4-diphenylcyclopentadienone; T. Mukai, Y. Yamashita, H. Sukawa, and T. Tezuka, Chem. Lett., 1975,423.
- 7. The yields of the diazepines 2 were obtained when the ylides disappeared, and no other products except the diazepines could be isolated.

(Received February 20, 1980)