Published on 01 January 1980. Downloaded by University of Illinois at Chicago on 22/10/2014 22:46:34.

Effective Method for the Conjugate Addition of γ -Acetoxy- $\alpha\beta$ -unsaturated Ketones with RCu-AlCl₃

By Toshiro Ibuka,* Hiroyuki Minakata, Yoshinori Mitsui, Kyoko Kinoshita, and Yumiko Kawami (Faculty of Pharmaceutical Sciences, Kyoto University, Kyoto 606, Japan)

Summary An efficient procedure for the conjugate addition of γ -acetoxy- $\alpha\beta$ -unsaturated ketones with RCu-AlCl₃ is described.

DESPITE the successful use of organocopper(I) reagents such as R_2CuLi in synthetic chemistry, considerable current interest is centred around the development of an effective method for conjugate addition as well as substitution reactions because of several drawbacks of the ordinary organocopper(I) reagents. By using lithium dimethylcuprate (Me₂CuLi), it has been reported that γ -acetoxy- $\alpha\beta$ unsaturated ketones were reduced to give $\alpha\beta$ - and/or $\beta\gamma$ -unsaturated ketones.^{1,2,3} Unfortunately, no successful conjugate addition reaction of γ -acetoxy- $\alpha\beta$ -unsaturated

```
TABLE. Yields of products in the reaction of \gamma-acetoxy-\alpha\beta-unsaturated ketones with Me<sub>2</sub>CuLi and RCu-AlCl<sub>3</sub>.<sup>a</sup>
```

Expt.	Substrate	Reagent	l,4-Adduct yield/%	Reduction yield	n product 1/%
1	(4)	Me ₂ CuLi	b	(6) 14	(7) 25
2	(5)	Me ₂ CuLi		(8) 75	(9) 16
3	(4)	MeCu-AlCl ₃	(10) 69, (11) 4		
4	(4)	PhCu-AlCl ₃	(12) 87 —		
5	(5)	MeCu-AlCl ₃	(13) 76, (14) 5		

^a The following procedure for a 1,4-addition reaction is typical. To a stirred suspension of MeCu (11.25 mmol) in dry ether (13.5 ml) under Ar at -70 °C was added dropwise a solution of AlCl₃ (11.25 mmol) in ether (3 ml) followed by a solution of the keto-acetate (1) (252 mg, 1.5 mmol) in dry ether (2 ml), and the temperature then raised to -25 °C. After each step the solution was stirred for 30 min. After the usual work-up, the product was purified by silica gel column chromatography to yield the pure adduct (2) (226 mg, 82% yield). ^b Though not isolated, gas chromatographic analysis of the reaction mixture showed the presence of < 0.75% of the conjugate adduct (10).



J.C S CHEM COMM, 1980

ketones has been reported, so we describe here our results with MeCu-AlCl₃ and PhCu-AlCl₃

It has been reported that the keto-acetate (1) was reduced with lithium dimethylcuprate to afford a mixture of 3-methylcyclohex-2-enone and 3-methylcyclohex-3-enone¹ In contrast, treatment of (1) with 5-7 5 mol equiv of an equimolar mixture of methylcopper(I) and aluminium trichloride (MeCu-AlCl₃) and phenylcopper(I) and aluminium trichloride (PhCu-AlCl₃) in ether at -70 to -25 ± 5 °C gave the conjugate adducts (2) and (3) in 82 and 95% yields, respectively (Scheme) † In these reactions, we did not detect any reduction product by gas chromatographic analysis

As shown in the Table, though the enones (4) and (5)were reduced with Me₂CuLi to afford $\alpha\beta$ - and $\beta\gamma$ -unsaturated ketones (6) and (7), and (8) and (9), respectively (expts 1 and 2), use of MeCu-AlCl₃ and PhCu-AlCl₃ provided the conjugate adducts in satisfactory yields (expts 3, 4, and 5) Application of the $RCu-AlCl_3$ system to a biologically active alkaloid is currently in progress

(Received, 11th August 1980, Com 882)

† All new compounds exhibited satisfactory spectroscopic and analytical data consistent with the structures

Reagents, 1, MeCu-AlCl₃ 11 PhCu-AlCl₃ 111 Me₂CuL1

- ¹ R A Ruden and W E Litterer, Tetrahedron Lett, 1975, 2043 ² P M Wege, R D Clark and C H Heathcock J Org Chem, 1976, 41, 3144 ³ E W Logusch, Tetrahedron Lett 1979, 3365

Scheme