

S0040-4039(96)00515-1

Synthetic Studies on the Common C25 Long Chain Acid Portion of Zooxanthellatoxins from a Symbiotic Dinoflagellate Symbiodinium sp.

Hideshi Nakamura,* Kazuhiro Fujimaki, and Akio Murai

Division of Chemistry, Graduate School of Science, Hokkaido University, Sapporo 060, Japan

Abstract: The absolute stereochemistry of a portion from C11 to C18 of the common C25 long chain acid part of zooxanthellatoxins was determined by synthesizing the degradation product 2a to be 115, 12R, 13S, 14R, 15R, 17R, and 18R. Copyright © 1996 Elsevier Science Ltd

Marine micro-algae produce various types of compounds including nitrogenous neurotoxins, polyether sea food toxins, sulfonium compounds of dimethylsulfide precursors, and antineoplastic macrolides.¹⁻³ Zooxanthellatoxin-A (ZT-A) and ZT-B have been isolated from a symbiotic marine dinoflagellate *Symbiodinium* sp. (strain No. Y-6).⁴ The structures contained several common structural units and consisted with two portions connected through an amide linkage, *i.e.*, the major amino acid portion forming a 62-membered lactone and the common terminal acid with C25 chain length (1).⁵ ZT-A and ZT-B showed similar biological activities such as vasoconstrictive and platelet aggregation activities.^{3,6}



In order to elucidate the mechanisms of biological activities of non-peptidic large molecules, we started to study on the structure-activity relationships and the absolute stereochemistry of zooxanthellatoxins by synthesizing their fragments obtained during the degradation experiments. Here, we report the synthesis and the absolute stereochemistry of the fragment containing C9-C21 portion of the common C25 long chain acid (1) in zooxanthellatoxins.

The acid portion 1 was obtained in a good yield from both ZT-A and ZT-B by alkaline hydrolysis. The relative stereochemistry at C11-C15 was deduced on the basis of the ${}^{3}J_{H-H}$ coupling constants of H11 to H15 protons.^{5d} Stereochemistry in the acyclic portion was unable to be clarified by spectroscopic methods. Ozonolysis of 1 followed by reduction with Me₂S, acetallization with HCl-MeOH, and acetylation afforded diacetate 2a containing the C9-C21 portion and a 1:1 mixture of methyl acetals of a γ -lactone (3) derived from the C1-C8 unit of zooxanthellatoxins. The structure 2a was confirmed by the synthesis started from known methyl altroside tetraacetate (4) by a sequence of the following reactions.⁷



Reagents and conditions: a) allylsilane, BF₃•OEt₂, TMSOTf, CH₃CN, -10 °C-23°C, 24 h, 71%. b) O₂, PdCl₂, CuCl, DMF-H₂O, rt, 45 h, 92%. c) MeONa, MeOH, 0 °C, 2h; *t*-BuOK,*t*-BuOH, 27 °C, 19 h; 1:1 dimethoxypropane-MeOH, CSA, 23 °C, 8 h, 44% from **6**. d) Tf₂O, Et₃N, CH₂Cl₂, -78 °C, 20 mi; BnOCH₂CH₂CH₂C≡CLi, THF, -78 °C, 1 h, 61%. e) Li, NH₃, THF, -78 °C, 7 h, 90%. f) Swern oxidation, 89%; AD-mix-β, MeSO₂NH₂, 1:1 H₂O-*t*-BuOH, 0 °C, 25 h, 96%.

The glycoside 4 prepared from methyl D-glucoside⁸ was treated with allyltrimethylsilane to afford Cglycoside 5 having the opposite stereochemistry at C11. Wacker oxidation of 5 gave methyl ketone 6^9 and the stereochemistry at C11 was adjusted by a base treatment after deacetylation of 6. Acetallization of the reaction mixture with 1:1 MeOH-dimethoxypropane gave methyl acetal 7^{10} in which the hydroxyl groups at C-13 and 14 and the methyl ketone were protected with an acetonide and a methyl acetal groups, respectively.

Chain elongation at C16 was achieved by the successive three step reactions, triflation, coupling reaction with an alkynyl lithium, and reduction under Birch conditions to yield *trans*-olefin alcohol 9. Introduction of two new chiral centers was conducted after oxidation of the terminal hydroxyl group with Sharpless asymmetric dihydroxylation reagents (AD-mix)¹¹ to give a mixture of diol 10. The diols were converted to dimethylacetal triacetate $2a^{12}$ in 37% yield by the following reactions; 1) removal of an acetonide in 80% AcOH at 23 °C for 24 h, 2) acetallization with HCl-MeOH at 23 °C for 30 min , and 3) acetylation with Ac₂O-Py at 23 °C for 16 h. The synthetic 2a was identical with one from natural products whereas a diacetate from 9 with AD-mix- α and the isomers prepared from *cis*-olefin alcohol with AD-mix- α were not identical.¹³ The stereochemistry at C17 position was established to be *R* by applying advanced Mosher method¹⁴ to a major dimethyl acetal 11¹⁵ prepared from 10. Configurations at C9 and C21 of 2a were determined by the NOESY spectrum (from each two MeO signals to H-12 and H-18 signals, respectively). Thus the stereochemistry of the synthetic 2a was proved as shown in the figure.

Absolute stereochemistry of 2a from zooxanthellatoxins was confirmed to be the same to that of synthetic one by comparing their $[\alpha]_D$ and the CD spectra of their benzoate derivative 2b.¹⁶ The stereochemistry in C11 to C18 of the common C25 long chain acid 1 was established chemically to be 11*S*, 12*R*, 13*S*, 14*R*, 15*R*, 17*R*, and 18*R*.

The C25 acid is the common terminal structure of ZT-A and ZT-B although their main chain structures are different each other in terms of their chain length and their oxidation state. It may be suggested that the C25 acid might be derived biogenetically before amidation with the main chains. Studies on the stereochemistry of C1-C8 portion and total synthesis of the C25 acid 1 are in progress at our laboratory.

Acknowledgments: This work was supported in part by a Grant-in-Aid from the Ministry of Education, Science and Culture, Japan.

References and Notes:

- 1. Faulkner, D., J. Nat. Prod. Rep. 1995, 223-269, and the references cited therein.
- 2. Shimizu, Y. Chem. Rev. 1993, 93, 1685-1698.
- 3. Yasumoto, T.; Murata, M., Chem. Rev. 1993, 93, 1897-1909.
- 4. Nakamura, H.; Asari, T.; Matsuoka, S.; Ohizumi, Y.; Kobayashi, J.; Yamasu, T.; Murai, A. *Toxicon*, **1993**, *31*, 371-376.
- (a) Nakamura, H.; Asari, T.; Fujimaki, K.; Maruyama, K.; Murai, A.; Ohizumi, Y.; Kan, Y. *Tetrahedron Lett.*, **1995**, *36*, 7255-7258. (b) Nakamura, H.; Asari, T.; Murai, A.; Kan, Y.; Kondo, T.; Yoshida, K.; Ohizumi, Y. J. Am. Chem. Soc., **1995**, *117*, 550-551. (c) Nakamura, H.; Asari, T.; Murai, A.; Kondo, T.; Yoshida, K.; Ohizumi, Y. J. Org. Chem., **1993**, *58*, 313-314. (d) Asari, T.; Nakamura, H.; Murai, A.; Kan, Y. Tetrahedron Lett., **1993**, *34*, 4059-4062.
- 6. Rho, M. -C.; Nakahata, N.; Nakamura, H.; Murai, A.; Ohizumi, Y. Br. J. Pharmacol. 1995, 115, 433-440.
- 7. All new compounds gave satisfactory spectral data.

- 8. Richtmyer, N. K., Methods in Carbohydrate Chemistry, 1962, 1, 107-113.
- 9. 6: [α]_D²⁰ +5.3° (c 3.15, CHCl₃); IR (film) 1748, 1374, 1228, and 1176 cm⁻¹; ¹H NMR (270 MHz, CDCl₃) δ 2.05 (3H, s), 2.08 (3H, s), 2.11 (3H, s), 2.13 (3H, s), 2.19 (3H, s), 2.57 (1H, dd, J=16, 4 Hz), 2.87 (1H, dd, J=16, 9 Hz), 4.10-4.20 (2H, m), 4.30 (1H, td, J=9, 4 Hz), 4.49 (1H, dd, J=11, 6 Hz), 5.04 (1H, t, J=9 Hz), and 5.26 (2H, m); HR-FABMS m/z 389.1425 (M+H)⁺. Calcd for C₁₇H₂₅O₁₀ 389.1448.
- 10. 7: [α]_D²² -20.9° (c 8.54, MeOH); ¹H NMR (400 MHz, CDCl₃) δ 1.36 (3H, s), 1.46 (6H, s), 1.97 (1H, br s, exchangeable), 2.06 (1H, d, J=15 Hz), 2.32 (1H, dd, J=15, 6 Hz), 3.23 (1H, m), 3.24 (3H, s), 3.58 (1H, br d, J=11, 6 Hz), 3.84 (1H, br d, J=11 Hz), 3.95 (1H, dd, J=10, 6 Hz), 4.16 (1H, dd, J=3, 1 Hz), 4.19 (1H, dd, J=6, 3 Hz), and 4.44 (1H, br d, J=6 Hz); HR-EIMS m/z 274.1396 Calcd for C₁₃H₂₂O₆ 274.1417.
- 11. Kolb, H. C.; VanNieuwenhze, M., S.; Sharpless, K., B. Chem. Rev., 1994, 94, 2483-2547.
- 12. Synthetic **2a**: $[\alpha]_D^{21}$ -32.9° (c 1.18, CHCl₃); IR (film) 2956, 1750, 1376, 1242, and 1100 cm⁻¹: ¹H NMR (400 MHz, CDCl₃) δ 1.48 (3H, s, H8), 1.56 (1H, m, H19), 1.65 (1H, m, H16), 1.83 (1H, m, H20), 1.92 (1H, m, H16), 1.97 (1H, m, H20), 2.01 (1H, m, H19), 2.01 (1H, m, H10), 2.03 (3H, s), 2.10 (3H, s), 2.11 (3H, s), 2.21 (1H, dd, *J*=14, 6 Hz, H10), 3.23 (3H, s), 3.32 (3H, s), 3.59 (1H, ddd, *J*=10, 9, 2, H15), 3.92 (1H, dd, *J*=4, 3 Hz, H12), 4.11 (1H, dt, *J*=7, 3 Hz, H18), 4.18 (1H, dd, *J*=4, 2 Hz, H11), 4.94 (1H, dd, *J*=9, 3 Hz, H14), 5.19 (1H, d, *J*=5 Hz, H21), 5.21 (1H, ddd, *J*=10, 4, 3 Hz, H17), and 5.40 (1H, t, *J*=3 Hz, H13); HR-FABMS *m*/z 473.1998 (M-H)⁺. Calcd for C₂₂H₃₃O₁₁ 473.2023. **2a** from ZT; $[\alpha]_D^{23}$ -38° (c 0.01, CHCl₃).
- 13. Dihydroxylation of the cis-olefin alcohol with AD-mix- α was not diastereoselective and gave a mixture of possible diol isomers.
- Ohtani, I., Kusumi, T., Kashman, Y., and Kakisawa, H., J. Am. Chem. Soc., 1991, 113, 4092-4096.
- Treatment of 10 with HCl-MeOH for 10 min at 23 °C gave 11 after HPLC separation. 11: ¹H NMR (400 MHz, CDCl₃) δ 1.37 (3H, s), 1.46 (3H, s), 1.47 (3H, s), 1.53 (1H, m), 1.70 (1H, m), 1.80-2.05 (4H, m), 2.02 (1H, d, J=15 Hz), 2.32 (1H, dd, J=15, 6 Hz), 2.55 (1H, d, J=6 Hz), 3.24 (3H, s), 3.34 (3H, s), 3.42 (1H, dt, J=10, 3 Hz), 3.73 (1H, m), 3.86 (1H, dd, J=10, 6 Hz), 3.98 (1H, dt, J=7, 5 Hz), 4.17 (2H, m), 4.43 (1H, d, J=6 Hz), 5.05 (1H, dd, J=5, 1 Hz).
- 16. Deacetylation of 2a followed by benzoylation in CH₂Cl₂ with benzoic acid, DCC, and DMAP gave 2b: ¹H NMR (400 MHz, CDCl₃) δ 1.25 (3H, s), 1.50-1.72 (2H, m), 1.76-1.96 (3H, m), 2.12 (1H, d, J=15 Hz), 2.07-2.17 (1H, m), 2.23 (1H, dd, J=15, 6 Hz), 3.26 (3H, s), 3.28 (3H, s), 3.98 (1H, t, J=9 Hz), 4.11 (1H, t, J=3 Hz), 4.24 (1H, m), 4.32 (1H, m), 5.03 (1H, d, J=5 Hz), 5.34 (1H, dd, J=4, 3 Hz), 5.54 (1H, m), 5.79 (1H, t, J=3 Hz), 7.23 (2H, t, J=8 Hz), 7.32 (2H, t, J=8 Hz), 7.45 (2H, t, J=8 Hz), 7.48 (1H, t, J=8 Hz), 7.49 (1H, t, J=8 Hz), 7.60 (1H, t, J=8 Hz), 7.82 (2H, d, J=8 Hz), 7.84 (2H, d, J=8 Hz), 8.08 (2H, d, J=8 Hz); CD (2-propanol) 237.5 nm (Δε +5.9) and 224.0 nm (-3.0). 2b from 1: CD (2-propanol) 237.5 nm (+3.3) and 224.0 nm (-2.2).

(Received in Japan 13 February 1996; revised 7 March 1996; accepted 15 March 1996)