


The Thermal Decomposition of Amine Imides

By D. G. MORRIS,* B. W. SMITH, and R. J. WOOD

(Department of Chemistry, The University, Sheffield, S3 7HF)

It has been shown that amine oxide pyrolyses proceed almost exclusively *via* a *cis*-elimination,¹ and that Hofmann eliminations of quaternary ammonium compounds can, in certain cases, proceed, at least in part, through a *cis*-elimination.² Both reactions are considered to occur *via* five-membered cyclic transition states.

In the present work thermal decompositions have been carried out on *NN*-dimethyl-2-phenylpropylamine acetimide (I) and *NN*-dimethylcyclo-octylamine acetimide (II). When (I) (with a trace of hydroquinone) and (II) were heated at 125° *in vacuo* the respective olefins, 2-phenylpropene (79%) and cyclo-octene (82%) were isolated. The cyclo-octene was found to consist of 96.5% *cis*- and 3.5% *trans*-cyclo-octene by gas chromatographic



SCHEME

analysis on di-isodecyl phthalate under conditions where no isomerisation occurs.[†] This compares with pure *cis*-cyclo-octene from amine oxide pyrolysis³ and *ca.* 85:15 *cis*:-*trans*-cyclo-octene from Hofmann elimination with phenyl-lithium.⁴ The only other isolable product was 2-acetyl-1,1-dimethylhydrazine which was characterised by comparison with an authentic sample.⁵

The olefins can most reasonably be considered to arise from a concerted *cis*-elimination involving a five-membered transition state (Scheme). Such a mechanism fills an isoelectronic gap in a series of dipolar part structures (III) → (V) capable of

effecting a pyrolytic *cis*-elimination. It is not possible at present to rule out a *cis*-elimination involving the oxygen end of the ambident anion. However such a process involving a seven-membered transition state is considered less likely. Two other types of pyrolytic decomposition of amine-imides are possible, *viz.* a Stevens rearrangement,⁶ and initial isocyanate formation.⁷ These processes have only been observed in structures with alkyl residues not containing β hydrogens and at higher temperatures.

(Received June 26th, 1968; Com. 846.)

[†] We are grateful to Dr. Robert Bach University of Minnesota for an authentic sample of *trans*-cyclo-octene.

¹ D. J. Cram and J. E. McCarty, *J. Amer. Chem. Soc.*, 1954, **76**, 5740.

² A. C. Cope and A. S. Mehta, *J. Amer. Chem. Soc.*, 1963, **85**, 1949; J. L. Coke and M. P. Cooke, *J. Amer. Chem. Soc.*, 1967, **89**, 6701; J. L. Coke, M. P. Cooke, and M. C. Mourning, *Tetrahedron Letters*, 1968, 2247.

³ A. C. Cope, R. A. Pike, and C. F. Spencer, *J. Amer. Chem. Soc.*, 1953, **75**, 3212.

⁴ G. Wittig and T. F. Burger, *Annalen* 1960, **632**, 85.

⁵ R. L. Hinman, *J. Amer. Chem. Soc.*, 1956, **78**, 1645.

⁶ S. Wawzonek and E. Yeakey, *J. Amer. Chem. Soc.*, 1960, **82**, 5718.

⁷ R. F. Smith and P. C. Briggs, *Chem. Comm.*, 1965, 120; M. S. Gibson and A. W. Murray, *J. Chem. Soc.*, 1965, 880.