

Preliminary communication

OXIDATIVE ADDITION OF α -KETOIMIDOYL CHLORIDES TO PALLADIUM(0) AND PLATINUM(0) COMPLEXES

ANTONIO MANTOVANI, GIACOMO FACCHIN,

Cattedre di Chimica, Facoltà di Ingegneria, University of Padova (Italy)

TRISTANO BOSCHI and BRUNO CROCIANI*

Centro Chimica Tecnologia Composti Metallorganici Elementi Transizione C.N.R., Istituto di Chimica Industriale, Via Marzolo, 9 Padova (Italy)





(Received October 13th, 1980)

Summary

The reaction of α -ketoimidoyl chlorides with palladium(0) and platinum(0) derivatives yields *trans*-[MCl{C(=NR)COMe}(PMe_nPh_{3-n})₂] (M = Pd; R = *p*-C₆H₄OMe, *c*-C₆H₄, *t*-Bu; *n* = 0, 1; M = Pt; R = *p*-C₆H₄OMe; *n* = 1) and *cis*-[PtCl{C(=N-*p*-C₆H₄OMe)COMe)COMe](PPh₃)₂], which have been characterized by IR, ¹H and ³¹P NMR spectra and by condensation with MeNH₂.

The reactions of imidoyl chlorides with transition metal substrates have been widely used to prepare complexes containing the imidoyl ligands [1–5]. In previous papers we reported that the acid-catalyzed hydrolysis of 1,4-diaza-3-methylbutadiene-2-yl-palladium(II) compounds yields the corresponding α -ketoimidoyl derivatives [6], from which new 1,4-diazadienyl groups with asymmetrically substituted imino nitrogen atoms can be obtained by condensation with primary amines [7]. We have now found a more convenient route to these compounds based on the oxidative addition of α -ketoimidoyl chlorides [8] to palladium(0) complexes (eq. 1).


Attempts to extend reaction 1 to other *d*¹⁰ metal complexes, such as [M(CH₂=CH₂)(PPh₃)₂] (M = Ni, Pt), [PtL₄] (L = PPh₃, PMePh₂), [RhCl(PPh₃)₃], [M'Cl(CO)(PPh₃)₂] (M' = Rh, Ir), have been successful only for platinum(0) derivatives and for R = *p*-C₆H₄OMe (eq. 2).



All the complexes I–VI give satisfactory elemental analyses and are monomeric in 1,2-dichloroethane. The $\nu(\text{C}=\text{O})$ and $\nu(\text{C}=\text{N})$ bands of the α -ketoimidoyl chlorides (1728–1723 and 1666–1638 cm^{-1} in benzene solution, respectively) are shifted to lower frequencies [ca. 40–50 cm^{-1} for $\nu(\text{C}=\text{O})$ and ca. 70–90 cm^{-1} for $\nu(\text{C}=\text{N})$] in the corresponding metal derivatives I–VI.

The ^{31}P NMR signals appear as a singlet for I–III (CD_2Cl_2 solution: I 40.06, II 37.38, III 41.75 ppm down-field from external $\text{P}(\text{Et}_3)_3$) and as a singlet flanked by ^{195}Pt satellites for VI [24.06 ppm, $^1J(\text{Pt}-\text{P}) = 3037$ Hz]. In the ^1H NMR spectra (CD_2Cl_2 solution) the $\delta(\text{PMe})$ signals occur as a triplet at 1.89 ppm for IV [$^2J(\text{P}-\text{H}) + ^4J(\text{P}'-\text{H}) = 6.8$ Hz] and as a triplet with ^{195}Pt satellites at 2.02 ppm for VI [$^2J(\text{P}-\text{H}) + ^4J(\text{P}'-\text{H}) = 7.0$ Hz, $^3J(\text{Pt}-\text{H}) = 31.4$ Hz]. These results indicate a *trans* configuration for I–IV and VI and the

presence of a plane of symmetry perpendicular to the coordination plane across the Cl-M-C_{imidoyl} unit. Although a time-averaged plane of symmetry might be generated by a fast rotation (on the NMR time scale) around the M-C_{imidoyl} bond, steric and electronic factors suggest a configuration of type A, with a planar O=C-C=N conjugated unit, analogous to the structure of the related compound *trans*-[PdCl{C(=NR)C(Me)=NR}(PPh₃)₂] (R = *p*-C₆H₄OMe) [9]:



(A, M = Pd, L = PPh₃, PMePh₂ ;

M = Pt, L = PMePh₂)

This configuration would also account for the down-field shift of the *ortho* protons of the Np-C₆H₄OMe group, resulting from the deshielding effect of the metal atom in close proximity (I δ = 7.60–7.75 ppm; IV 7.65–7.80 ppm; VI 7.70–7.85 ppm) and for the high-field shift of the methyl protons of the COMe group, due to the shielding effect of the phenyl ring current of two mutually *trans* PPh₃ ligands (I δ = 1.34 ppm; II 1.23 ppm; III 1.31 ppm) [9].

When a PPh₃-platinum(0) substrate is used in reaction 2, a *cis* product V is obtained, as shown by its ³¹P NMR spectrum in CD₂Cl₂ (δ = 32.09 ppm, ¹J(Pt-P) = 4504 Hz, for P *trans* to chlorine; δ = 37.60 ppm, ¹J(Pt-P) = 1729 Hz, for P *trans* to carbon, ²J(P-P') = 17.1 Hz). The different geometry of complexes V and VI is also reflected in the different ν (Pt-Cl) values, 295 and 270 cm⁻¹, respectively, resulting from the higher *trans*-influence of the α -ketoimidoyl group compared to PPh₃. Accordingly, in the *trans* complexes I–IV in the ν (Pd-Cl) band was found in the low frequency range 270–250 cm⁻¹.

The compounds I and III undergo condensation reaction at the carbonyl group by MeNH₂ to give the 1,4-diazadienyl derivatives *trans*-[PdCl-{C(=NR)C(Me)=NMe}(PPh₃)₂] (R = *p*-C₆H₄OMe, *c*-C₆H₁₁) [7]. We are now studying this condensation reaction with the platinum complexes V and VI, and the reaction of α -ketoimidoyl chlorides with some metal carbonyl anions.

References

- 1 M.J. Doyle, M.F. Lappert, G.M. McLaughlin and J. McMeeking, J. Chem. Soc., Dalton, (1974) 1494.
- 2 R.D. Adams, D.F. Chodosh and N.M. Golembeski, J. Organometal. Chem., 139 (1977) C39.
- 3 H. Alper, M. Tanaka and K. Hachem, J. Organometal. Chem., 190 (1980) 95, and references cited therein.
- 4 F. Porta, S. Cenini, P. Del Buttero and S. Maiorana, J. Organometal. Chem., 194 (1980) 211.
- 5 K. Hiraki, Y. Fuchita and T. Masumoto, Bull. Chem. Soc. Jpn., 53 (1980) 1171.
- 6 B. Crociani, Inorg. Chim. Acta, 23 (1977) L1.
- 7 B. Crociani and R.L. Richards, J. Organometal. Chem., 154 (1978) 65.
- 8 I. Ugi and U. Fetzner, Chem. Ber., 94 (1961) 1116.
- 9 B. Crociani, G. Bandoli and D.A. Clemente, J. Organometal. Chem., 184 (1980) 269.