Methane-Producing Hydrogenolysis of $(\eta^{5}$ -Cyclopentadienyl)(triphenylphosphine)dimethylcobalt(III). An Autocatalytic Mechanism Involving a Binuclear Metal Dihydride/Metal Dialkyl Reaction as a **Critical Step**

Andrew H. Janowicz and Robert G. Bergman*

Department of Chemistry, University of California Berkeley, California 94720 and Materials and Molecular Research Division Lawrence Berkelev Laboratorv Berkeley, California 94720

Received December 5, 1980

The cleavage of organotransition metal-carbon bonds by H₂ is an important part of many catalytic processes. For example, the terminating steps in hydroformylation¹ and homogeneous hydrogenation² require cleavage of a metal acyl or alkyl by H_2 , and the (heterogeneous) Fischer-Tropsch reaction requires several steps in which C-H bonds are formed, including the final release of hydrocarbon or oxygenate from the metal surface.³

It has commonly been assumed that M-R hydrogenolysis occurs by oxidative addition of H₂ to the metal center at which the carbon fragment is attached, followed by formation of a new C-H bond by reductive elimination (eq 1).⁴ However, evidence has recently

$$M-R + H_2 \rightarrow [R-MH_2] \rightarrow M-H + R-H$$
(1)

been obtained⁵ that in hydroformylation, the final C-H bond forming step is a binuclear process, involving reaction between a transient cobalt acyl complex and the hydroformylation catalyst, $HCo(CO)_4$. Several other M-H/M-R reactions (eq 2) have also

$$M-R + M-H \rightarrow [R-M-H-M] \rightarrow M-M + R-H \quad (2)$$

been discovered recently.⁶ Despite this, however, studies aimed at the direct investigation of the mechanism of metal-carbon bond hydrogenolysis reactions are rare. In this paper, we wish to report out results on the mechanism of the facile methane-forming hydrogenolysis of cyclopentadienyldimethylcobalt derivatives. We chose these complexes for study because they give methane on hydrogenolysis, and yet it seemed unlikely that this process could be occurring by loss of phosphine followed by oxidative addition of H_2 to give a formally Co(V) intermediate. We find that even though no metal hydrides are added as reagents in this reaction, the new C-H bonds are still formed by a binuclear M-H/M-R reaction.

Reaction of $(\eta^5-C_5H_5)Co(PPh_3)(CH_3)_2$ (1) with dihydrogen in benzene- d_6 at 45 °C yields 2 equiv of CH₄ and 0.5 equiv of $(\eta^5 - C_5 H_5) Co(PPh_3)_2$ (2) (Scheme I). In the presence of an added equivalent of PPh₃, the reaction rate is slower but the hydrogenolysis now gives 2 equiv of CH_4 and nearly 1 equiv of 2 (Scheme I). Treatment of complex 1 with D_2 under the same conditions gives CH₃D, establishing that the new hydrogen atom does in fact come from the hydrogen gas.

In agreement with the notion that a conventional mechanism might not be accessible for this process, monitoring the reaction of 1 with H_2 in the presence of added phosphine revealed the presence of an induction period, whose length was dependent upon the concentration of phosphine (Figure 1). Following the induction period, the rate gradually acclerated until the system began

(1) For a review, see: Orchin, M.; Rupilius, W. Catal. Rev. 1972, 6, 85. (2) See, e.g.: Dolcetti, G.; Hoffman, N. W. Inorg. Chim. Acta. 1974, 9, 269



Scheme II

Scheme I

$$\begin{array}{cccc} (\eta^{5}-C_{5}H_{5})\text{Co}(\text{PPh}_{3})\text{Me}_{2} &\rightleftharpoons (\eta^{5}-C_{5}H_{5})\text{Co}\text{Me}_{2} + \text{PPh}_{3} \\ & 1 & 3 \\ (\eta^{5}-C_{5}H_{5})\text{Co}(\text{PPh}_{3})_{2} &\rightleftharpoons (\eta^{5}-C_{5}H_{5})\text{Co}(\text{PPh}_{3}) + \text{PPh}_{3} \\ & 2 & 4 \\ & 4 + H_{2} &\rightleftharpoons (\eta^{5}-C_{5}H_{5})\text{Co}(\text{PPh}_{3})\text{H}_{2} \\ & 5 \\ & 5 + 3 & \frac{\text{PPh}_{3}}{2} 2\text{CH}_{4} + 2 \end{array}$$

to be depleted in complex 1. Such induction periods are often characteristic of free-radical chain processes; however, the operation of a radical chain in this system seemed somewhat unlikely. because it would presumably have to involve methyl radicals. These very reactive radicals would almost certainly produce some toluene by reaction with benzene solvent; in addition, because of the comparable bond energies of H₂ and methane, H₂ should not transfer hydrogen very rapidly to methyl radicals. Nevertheless, we carried out the hydrogenolysis using D₂, $(\eta^5-C_5H_5)C_0$ -(PPh₃)(CD₃)₂, and 10 equiv (0.95 M) of 1,4-cyclohexadiene, a very reactive reagent for transfer of hydrogen atoms to organic radicals.⁷ Under these conditions the methane produced was still exclusively CD₄. This suggested strongly that a radical chain process was not responsible for the observed induction period.

A second explanation for the induction period is that the hydrogenolysis is autocatalytic; we have been able to obtain much stronger experimental support for this hypothesis. As stated above, this rate behavior is observed when the reaction is performed in the presence of added phosphine. Adding both phosphine and the CpCo(PPh₃)₂ reaction product, however, makes the induction period disappear (although the overall rate still exhibits inhibition by PPh₃), and gradually increasing rates are observed immediately after mixing the reagents (Figure 1). Clearly product 2 is a catalyst (or catalyst precursor) for the hydrogenolysis of 1.

An autocatalytic mechanism consistent with the observations reported above is outlined in Scheme II. We know that the thermal decomposition of 1 is slow at 45 °C and has a complicated mechanism, but some Co(I) species are produced in this reaction.⁸ We believe these initiate the hydrogenolysis, and once some 4 is produced, this takes over as the critical Co(I) species. We sought additional support for this mechanism by attempting to prepare dihydride 5, so that we might investigate its reactions with 1 directly. Unfortunately, reaction of 1 with H_2 , reaction of 2 with excess H_2 , or treatment of CpCo(PR₃)I₂ derivatives with hydride reagents gave no evidence for the formation of stable dihydrides. As an alternative, therefore, we prepared⁹ 6, the pentamethylcyclopentadienyl analogue of 1. When 6 and excess H_2 were allowed to react in benzene- d_6 at 45 °C, methane was again produced, along with a new material whose proton NMR spectrum [δ 1.77 (15 H, s), 6.90–7.10 (9 H, m), 7.70–7.83 (6 H, m), -16.54 $(2 \text{ H}, d, J_{P-H} = 87 \text{ Hz})]$ was consistent with its formulation as $(\eta^5 - C_5 Me_5)Co(PPh_3)H_2$ (7). This material showed a strong absorption in the IR at 1908 cm⁻¹. This was assigned as ν_{M-H} on the basis of its shift to 1376 cm⁻¹ when metal-bound hydrogen was replaced with deuterium.10

^{*} Address correspondence to this author at the University of California.

⁽³⁾ See, e.g.: Masters, C. Adv. Organomet. Chem. 1979, 17, 61.
(4) Schrock, R. R.; Parshall, G. W. Chem. Rev. 1976, 76, 243.

⁽⁵⁾ Alemdaroglu, N. H.; Penninger, J. L. M.; Oltay, E. Monatsh. Chem. 1976, 107, 1153.

^{(6) (}a) Norton, J. R. Acc. Chem. Res. 1979, 12, 139. (b) Okrasinski, S. J.; Norton, J. R. J. Am. Chem. Soc. 1977, 91, 295. (c) Wolczanski, P. T.; Threlkel, R. S.; Bercaw, J. E. Ibid. 1979, 101, 278. (d) Renaut, P.; Tainturier, G.; Gautheron, B. J. Organomet. Chem. 1978, 150, C9.

⁽⁷⁾ Kaplan, L. React. Intermed. 1978, 1, 163.

⁽⁸⁾ Evitt, E. R.; Bergman, R. G. J. Am. Chem. Soc. 1980, 102, 7003. (9) Evit, bet gillari, N. O. S. Am. Chem. Bott. 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, 1966, shown in Scheme III.

⁽¹⁰⁾ The corresponding diphosphine, $(\eta^5-C_5(CH_3)_5)Co(PPh_3)_2$ (11), is not formed, presumably due to steric destabilization.



Figure 1. Concentration vs. time plots for the disappearance of complex 1 (0.10 M initial concentration of 1; 3 atm H₂), in benzene- d_6 at 45 °C, with the following additives: (Δ) no additives, (\diamond) 0.025 M PPh₃, (\Box) 0.05 M PPh₃, (\times) 0.10 M PPh₃, (\bigcirc) 0.10 M PPh₃ and 0.017 M 2.

Scheme III



Complex 7 is one of the few mononuclear dihydridocobalt complexes which have been prepared. Although rapid H₂ loss has prevented us from isolating 7 as a pure solid, benzene solutions of the dihydride are stable for relatively long periods of time. This stability encouraged us to carry out a direct study of the interaction of dimethyl complex 6 with 7. Dialkyl 6 and dihydride 7 reacted readily at 45 °C, as predicted by Scheme II, to give 2 mol of methane. We have not yet been able to characterize the organometallic products of this reaction, but in the presence of excess 7 and PPh₃, the rate of disappearance of 6 (as determined by ¹H NMR spectroscopy) showed good pseudo-first-order kinetics. Steady-state analysis of the mechanism shown in Scheme III gives a rate law which predicts that the reciprocal of the pseudofirst-order rate constant k_{obsd} should depend upon the dihydride and PPh₃ concentrations as shown in eq 3. The rate constant

$$1/k_{\text{obsd}} = (1/k_1) + [k_{-1}[\text{PPh}_3]/(k_1k_2[7])]$$
(3)

 k_{obsd} was therefore determined at several different concentrations of PPh₃ and 7, and plotted according to eq 3. As shown in Figure 2, the predicted linear behavior is observed. From the slope and intercept of the plot, we calculate $k_1 = 1.83 \times 10^{-4} \text{ s}^{-1}$ and $k_{-1}/k_2 = 1.44$.

Scheme III and the corresponding rate law posit a specific level of unsaturation of each reacting partner in the rate-determining step associated with microscopic rate constant k_2 ; i.e., the dialkyl reactant is coordinatively unsaturated, whereas the hydride is saturated. The rate law itself rules out the possibility that both reactants are unsaturated; reversible dissociation of phosphine followed from both 7 and 6, followed by reaction of the two unsaturated species so produced, would give an inverse squared dependence upon the concentration of phosphine, which is not observed. That it is the dialkyl complex, rather than the dihydride, which is unsaturated is suggested by the ease with which PMe₃ replaces PPh₃ in 6. This result parallels that of 1 which undergoes PMe₃ substitution by a dissociative mechanism⁸ through intermediate 3.

In summary, even though the inaccessibility of the formal Co(V) oxidation state prevents oxidative addition of H_2 to cyclopentadienylcobalt(III) derivatives, they still undergo facile hy-



Figure 2. Dependence of the pseudo-first-order rate constant for reaction of complex 6 with dihydride 7 (measured in the presence of excess 7 and PPh₃) upon the ratio of PPh₃ to 7.

drogenolysis, and this hydrogenolysis involves a metal alkyl/metal hydride reaction as a critical step. The lack of a stoichiometric amount of metal hydride reagent present at the beginning of the hydrogenolysis reaction leads to an autocatalytic mechanism; direct reaction between dialkyl and dihydride does not require autocatalysis. As in other reactions involving interaction of a metal hydride with a coordinatively unsaturated alkyl,⁶ we do not yet know the precise structure of the intermediate involved in the M-R/M-H step, but we assume interaction initially involves entry of the hydride end of the M-H bond into the metal coordination sphere of the dialkyl, forming a bridging hydride. The precise structure of this complex, and how reductive elimination occurs within it, is not yet clear.¹¹ It is interesting, however, that on the basis of the relative rate constant ratio k_2/k_{-1} measured in our kinetic study, PPh₃ reacts with the unsaturated intermediate 8 only slightly more rapidly than does the dihydride 7.

Acknowledgment. This work was supported by the Director, Office of Energy Research, Office of Basic Energy Sciences, Chemical Sciences Division of the U.S. Department of Energy under Contract Number W-7405-ENG-48.

Synthesis and Reactions of a Binuclear Cobalt Bridging Methylene (μ -CH₂) Complex. Conversion to μ -CH₂ Rh/Co and Rh/Rh Complexes and Methylene Transfer to Ethylene Involving Activation by a Second Metal Complex

Klaus H. Theopold[†] and Robert G. Bergman*

Department of Chemistry, University of California Berkeley, California 94720

Received January 14, 1981

Bridging methylene complexes are among the most interesting binuclear metallacycles, both for structural reasons¹ and because

⁽¹¹⁾ One possibility (also suggested by both referees) is that the unsaturated dialkyl and saturated dihydride first react to give $CpCo(PPh_3)(CH_3)(H)$ and $CpCo(CH_3)(H)$. These alkyl hydrides can then undergo rapid reductive elimination.

[†]Holder of an Ephraim Weiss Scholarship from the University of California, Berkeley, 1979-present.