


Cyclization

Intramolecular Carbostannylation of Alkynes Catalyzed by Silver(I) Species**

Susana Porcel and Antonio M. Echavarren*

Cyclizations of α,ω -enynes catalyzed by transition- or main-group metals provide highly functionalized carbo- and heterocycles under mild conditions in atom-economical processes.^[1] We have previously shown that reactions of allylsilanes or allylstannanes **1** with different metal catalysts proceed to form hetero- or carbocycles **2** (Scheme 1).^[2,3] The



Scheme 1. Cyclizations of allylsilanes or allylstannanes **1** with different metal catalysts. $M = Pt^II, Pd^II, Cu^I, Ru^II, Ag^I, Au^III$; $L =$ ligand; $Y = SiMe_3, SnBu_3$; $Z = C(CO_2Me)_2, C(SO_2Ph)_2, C(CH_2OR)_2$.

best results were obtained by using $PtCl_2$ or $[Pt(MeCN)_2Cl_2]$ as the catalyst and methanol or acetone as the solvent. The reaction proceeds by *exo* attack of the allyl nucleophile to the alkyne to form carbocycles with five- or six-membered rings. Interestingly, when the reactions of the substrates **1** ($Y = SnBu_3$) were catalyzed by palladium(0) complexes, stannyl derivatives **3** were obtained stereoselectively in a process that involves a very different mechanism to that seen in the oxidative addition of the allylstannane to Pd^0 .^[4] Stannanes **4** have been obtained, along with the *Z* isomers (ca. 9:1 *E/Z*

selectivity), in the radical cyclization of substrates **1** performed in the presence of azobisisobutyronitrile (5 mol %) and Bu_3SnH (10 mol %).^[5] Conversely, when the reaction of **1** was promoted by $GaCl_3$ (1 equiv), products **2** were obtained exclusively, whereas a similar reaction with $InCl_3$ (1 equiv) led to **2** as the major products along with mixtures of **3** and **4** in very low yields.^[6–8]

Reactions of simple 1,6-enynes with platinum(II)^[9] or gold(I)^[10] species have been shown to proceed through cyclopropyl metal–carbene intermediates. In many of the gold(I)-catalyzed cyclizations of enynes and other alkynes, a silver salt such as $AgOTf$, $AgBF_4$, or $AgSbF_6$ ($Tf =$ trifluoromethanesulfonyl) is used to generate in situ a cationic gold(I) species from $[AuCl(L)]$ complexes ($L =$ phosphine or a related ligand).^[10–12] The silver salt has been shown to be catalytically inert for many of these processes,^[10,11] although in a few instances silver(I) species can catalyze cyclization reactions.^[13] Herein we report that, in contrast to gold(I) complexes, silver(I) salts and complexes are catalytically active in the cyclization of **1** and afford stannanes **4** as the exclusive stereoisomers. We also report the first examples of skeletal rearrangements and intramolecular cyclopropanation reactions of 1,6-enynes catalyzed by silver(I) complexes, which suggest that metal–carbene intermediates are also involved in these silver(I)-catalyzed transformations.^[14,15]


When the reaction of **1a** with $AgOTf$ (10 mol %) was carried out in toluene at 70 °C, stannane **4a** was obtained in 29% yield, along with **2a** (Table 1, entry 1). The use of $AgBF_4$ led to **4a** and dimer **5** in low yield, whereas complex **6** was ineffective (Table 1, entries 2 and 3). Remarkably, $AgSbF_6$ or $AgSbF_6/PPh_3$ led to a very fast cyclization of (*Z*)-**1a**, and yielded **4a** (83%) and **2a** (5–12%) (Table 1, entries 4 and 5). The reaction also proceeded satisfactorily in the presence of a variety of silver(I) complexes $[Ag(OTf)L]$ bearing phosphines as the ligands ($L = PPh_3, (o\text{-tolyl})_3P, (naphthyl)_3P, 2\text{-biphenyldicyclohexylphosphine}$ or $[(AgOTf)_2(L-L)]$ ($L-L =$ ethane-1,2-diylbis(diphenylphosphane) (*dppe*), 9,9-dimethyl-4,5-bis(diphenylphosphino)xanthene (*xantphos*), 2,2'-bis(diphenylphosphanyl)-1,1'-binaphthyl (*binap*)),^[16,17] which gave **4a** in 85–91% yield. Consistent results were obtained with the preformed $[(Ag(OTf)(PPh_3))_3]$ complex^[18] as catalyst (Table 1, entry 6). No reaction was observed when the ratio of L to Ag was higher than 1:1.

The reaction of substrates similar to **1a** proceeds satisfactorily regardless of their *E/Z* configuration. For example, (*E*)-**1a** (Table 2, entry 1) reacts similarly to (*Z*)-**1a**, and both (*E*)- and (*Z*)-**1d** gave **4d** in 90–91% yields after 30 minutes (Table 2, entries 4 and 5). This silver(I)-catalyzed reaction tolerates protection of the hydroxy groups with acetate and *tert*-butyldiphenylsilyl (TBDS) groups (Table 2, entries 6

[*] S. Porcel, Prof. Dr. A. M. Echavarren
Institute of Chemical Research of Catalonia (ICIQ)
Av. Països Catalans 16, 43007 Tarragona (Spain)
and
Department of Organic Chemistry
Universidad Autónoma de Madrid
Cantoblanco, 28049 Madrid (Spain)
Fax: (+34) 97-792-0225
E-mail: aechavarren@icq.es



[**] We are grateful to the MEC (project CTQ2004-02869 and Consolider Ingenio 2010, Grant CSD2006-0003), the UAM (predoctoral fellowship to S.P.), the AGAUR (2005 SGR 00993), and the ICIQ Foundation for financial support. We also thank E. Escudero-Adán and Dr. J. Benet-Buchholz (X-ray diffraction unit, ICIQ) for the determination of the structure of complex **8**.

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.

Table 1: Silver(I)-catalyzed carbostannylation of (Z)-**1a**.



Entry	Cat. (mol %)	<i>t</i>	Yield of 4a [%]	Yield of 2a [%]
1 ^[a]	AgOTf (10)	12 h	29	24
2 ^[a]	AgBF ₄ (10)	12 h	29	— ^[b]
3	6 (5)	12 h	—	—
4	AgSbF ₆ (10)	1 min	83	12
5	AgSbF ₆ (10) + PPh ₃ (10)	1 min	83	5
6	[Ag OTf(PPh ₃) ₃] (3)	30 min	90	—

[a] Conversion of 85%. [b] Dimer **5** (7%) was obtained. Tf = trifluoromethanesulfonyl, Mes = 2,4,6-trimethylphenyl.


Table 2: Silver(I)-catalyzed carbostannylation of **1b–h**.^[a]


Entry	Substrate	<i>t</i> [h]	Product(s) (yields, [%])
1	(E)- 1a : Z = C(CO ₂ Me) ₂ , R ¹ = H, R ² = Me	0.5	4a (87) + 2a (3)
2	(E)- 1b : Z = C(CO ₂ Me) ₂ , R ¹ = R ² = H	5	4b (71) + 2b (6)
3	(Z)- 1c : Z = C(SO ₂ Ph) ₂ , R ¹ = R ² = H	3.5	4c (72) + 2c (9)
4	(E)- 1d : Z = C(SO ₂ Ph) ₂ , R ¹ = H, R ² = Me	0.5	4d (90)
5	(Z)- 1d : Z = C(SO ₂ Ph) ₂ , R ¹ = H, R ² = Me	0.5	4d (91)
6	(E)- 1e : Z = C(CH ₂ OAc) ₂ , R ¹ = R ² = H	2.5	4e (69) + 2e (12)
7	(E)- 1f : Z = C(CH ₂ OTBDPS) ₂ , R ¹ = R ² = H	2	4f (93)
8	(E)- 1g : Z = C(CH ₂ OH) ₂ , R ¹ = R ² = H	0.2	4g (31) + 2g (11) + 7 (30)
9 ^[b]	(E)- 1h : Z = C(SO ₂ Ph) ₂ , R ¹ = Ph, R ² = Me	3	4h (50) + 2h (50)


[a] Reaction with [AgOTf(PPh₃)₃] (3 mol %) in toluene at 70°C. [b] Reaction carried out with catalyst **8** (10 mol %) in toluene at 90°C. Cy = cyclohexyl.



and **7**). The free hydroxy groups in substrate **1g** compete in the reaction with the alkyne, leading to a 1:1 ratio of bicyclic acetal **7** and stannane **4g** (Table 2, entry 8). A similar

cyclization of diols with alkynes has been reported by Genêt and co-workers with gold(I) catalysts.^[19] Substrate **1h**, which is substituted at the alkyne with a phenyl group, reacted with the catalyst [Ag(2-biphenyldicyclohexylphosphine)(thf)]SbF₆ (**8**)^[20] to give a 1:1 ratio of **4h** and **2h** in quantitative yield. In this case, extensive destannylation of **4h** was observed. The structure of the cationic silver(I) complex **8** is similar to the related gold(I) complexes.^[10c,20,21] In contrast, the radical reaction of (E)-**1h** reported by Hosomi and co-workers occurred exclusively by a 6-endo-dig pathway.^[5] Alkenyl stannanes **4a–h** were obtained as single stereoisomers, whose configuration was assigned as *E* by comparison of their NMR spectra with those of the *Z* isomers **3**^[4] as well as by reaction of **4c** with DCI in CD₃OD.^[22] Reaction of the analogous allylsilanes under these conditions proceeded rather sluggishly to give only products **2**.

The reaction can be extended for the preparation of six- and seven-membered-ring compounds (Scheme 2). For these cyclizations, complex **8** was found to be the best catalyst


Scheme 2. Silver(I)-catalyzed carbostannylation for the synthesis of six- and seven-membered-ring systems. Ts = toluene-4-sulfonyl.

(Figure 1). Substrate **9**, which has an additional methylene group at the stannane chain, afforded the six-membered-ring derivatives **10** and **11**. A 7-endo-dig cyclization was the predominant pathway in the reaction of tosylamide **12**, which afforded heterocycle **13** as the major product after treatment of the mixture of stannanes with I₂. A seven-membered ring **16** was also obtained in the 7-exo-dig cyclization of **15**.

Although sterically hindered, the alkenylstannanes **4a** and **4d** undergo Stille reactions with iodobenzene in the presence of [Pd(PPh₃)₄] (10 mol %), CuI (10 mol %), and CsF (4 equiv) in THF^[23] to give stereospecifically **18** and **19**, respectively (Scheme 3). Compound **19** is the *E* diastereomer of **2h** (Table 2, entry 9), thus providing further confirmation of the configuration assigned for **4a–h**.





Figure 1. X-ray crystal structure of the cation of silver(I) complex **8**.




Scheme 3. Stillé coupling reactions of stannanes **4a** and **4d**.

A similar reaction of substrates **1** with gold(I) catalysts^[10] led only to destannylated products **2**. For example, reaction of (*Z*)-**1c** with $[\text{AuCl}(\text{PPh}_3)]$ (5 mol %) was very sluggish and led only to **2c** in around 20% yield (toluene, 70°C, 14 h). Conversely, reaction of (*Z*)-**1c** with the cationic complex $[\text{Au}[\text{P}(\text{C}_6\text{H}_4(o\text{-Ph})_2)(\text{tBu})_2]\{\text{NCMe}\}]\text{SbF}_6$ ^[20,21] was complete in 30 minutes at 70°C to give **2c** in 97% yield. Reaction of (*Z*)-**1a** with Lewis acids such as Et_2AlCl and ZrCl_4 (toluene, –78 to 0°C, 12–24 h) led only to the partial destannylation of the starting material.^[24]

The enantioselective cyclization of allylstannanes with alkynes was examined with substrate (*E*)-**1d** using the complexes $[(\text{AgOTf})_2(\text{L-L})]$ with chiral bidentate ligands.^[25] Bidentate ligands such as (+)-(2*S*,3*S*)-bis-(diphenylphosphino)bicyclo[2.2.1]hept-5-ene ((+)-(S,S)-norphos), (–)-(R,R)-*P,P'*-[2,2-dimethyl-1,3-dioxolane-4,5-bis(methylene)]bis(diphenylphosphane) ((–)-(R,R)-diop), (R,R)-1-benzyl-3,4-bis(diphenylphosphino)pyrrolidine ((R,R)-deguphos), (2*S*,4*S*)-*N*-tert-butoxycarbonyl-4-diphenylphosphanyl-2-diphenylphosphanylmethylpyrrolidine ((2*S*,4*S*)-bppm) in combination with AgOTf (20 mol % of AgOTf, 9% mol of ligand) gave low enantiomeric ratios (e.r.), ranging from 53:47 to 63:37. The best results were obtained when isolated complexes $[(\text{AgOTf})_2(\text{R})\text{-binap}]$ or $[(\text{AgOTf})_2(\text{R})\text{-Tol-binap}]$ ^[16b] were employed (Table 3). The nature of the counteranion plays a crucial role; for example, no enantioselectivity was achieved with complex $[(\text{AgSbF}_6)_2(\text{R})\text{-Tol-binap}]$ (Table 3, entry 1), but use of the OTf, which is a better coordinating anion, led to a 89:11 e.r. under the same reaction conditions (Table 3, entry 5). The corresponding complex formed between (*R*)-Tol-binap and AgPF_6 led to lower enantioselectivity (68:32 e.r.), whereas with AgBF_4 , only the destannylated

Table 3: Enantioselective cyclization of (*E*)-**1d**.^[a]



Entry	Catalyst	T [°C]	Yield [%]	e.r. ^[b]
1	$[(\text{AgSbF}_6)_2\text{Tol-binap}]$	30	57	50:50
2	$[(\text{AgOTf})_2\text{binap}]$	70	87	86:13
3	$[(\text{AgOTf})_2\text{Tol-binap}]$	70	87	87:12
4	$[(\text{AgOTf})_2\text{Tol-binap}]$	50	91	89:11
5 ^[c]	$[(\text{AgOTf})_2\text{Tol-binap}]$	30	74	89:11

[a] Reactions with 5 mol % catalyst for 30 min. [b] Determined by HPLC (Daicel Chiralpack AD column). [c] Reaction time: 100 min. Tol-binap = 2,2'-bis(*di-p*-tolylphosphanyl)-1,1'-binaphthyl.


product **2a** was obtained. Reaction of (*Z*)-**1d** under the same conditions as in entry 4 led to stannane **4d** with 80:19 e.r.

Yamamoto and co-workers have suggested that a transmetalation between the silver(I) complex and allyltrimethylsilanes may take place.^[16a-c] However, in the reactions of allylstannanes with aldehydes, the silver(I) complex was proposed to act as a chiral Lewis acid rather than forming an allyl–silver(I) species.^[16d] A transmetalation of the allylstannane with silver(I) would lead to the products **3**, after insertion of the allyl–silver(I) species into the alkyne followed by reductive elimination. To exclude this pathway, we prepared the *Z* isomer of **1a** from the cyclization of (*E*)-**1d** with $[\text{Pd}_2(\text{dba})_3\text{dba}]$ (dba = *trans,trans*-dibenzylideneacetone) as the catalyst.^[4] However, no isomerization of this substrate into **4d** was observed after the substrate was heated with complex **8** (10 mol %) in toluene at 70°C for 14 h.


The formation of silver(I)–acetylide complexes is not a major pathway under these reaction conditions.^[26] Instead, the isolation of the products **4** is consistent with a mechanism in which the silver(I) complex selectively activates the alkyne of **1** to form the cyclopropyl carbene–silver(I) complex **20**, followed by cleavage of the cyclopropane to form the alkenyl–silver(I) complex **21** (Scheme 4).^[2,3,10] Reaction of **21** with Bu_3SnOTf (or a similar electrophile in the case of AgSbF_6 or catalyst **8**) gives the stannanes **4**. Alternatively, formation of **21** might take place in a single step as shown in Scheme 4.

To support the formation of silver carbenes in these cyclizations, we examined the reactions of simple enynes with silver(I) catalysts (Scheme 5). Thus, **22** afforded in quantitative yield diene **23**, the product of a skeletal rearrangement with a single cleavage, as shown in the reaction of deuterated substrate $[\text{D}_1]\text{-22}$ to give $[\text{D}_1]\text{-23}$.^[10c] Dienyne **24** led to **25** in 66% yield, along with the skeletal-rearrangement products as minor compounds. Tetracycle **25** is identical to that obtained before using gold(I) species as the catalyst.^[10d] Conversely, **26** gave rise to a 15:1 mixture of the skeletal-rearrangement derivative **27** and the product of an intramolecular cyclopropanation **28** (94% yield, Scheme 5).^[27]

In summary, we have reported the first intramolecular carbostannylation of alkynes catalyzed by silver(I) species, which gave (*E*)-alkenylstannanes stereoselectively as single isomers in a reaction that appears to be mechanistically



Scheme 4. Mechanistic hypotheses for the silver(I)-catalyzed carbostannylation of alkynes.



Scheme 5. Skeletal-rearrangement and cyclopropanation reactions catalyzed by silver(I). DCE = 1,2-dichloroethane.

similar to that of the reaction of enynes with other electrophilic transition-metal complexes. In this case, the alkenyl–silver(I) intermediate is able to react with the tin electrophile generated in situ, thus leading to stannanes of structure **4** with total control of the stereoselectivity. A 78% *ee* has been achieved by using [(AgOTf)₂(*R*)-Tol-binap] as the catalyst. We have also reported the first examples of skeletal rearrangements and intramolecular cyclopropanation reactions of 1,6-enynes catalyzed by silver(I) species, an observation that indicates that silver–carbene species are probably involved in these reactions.

Received: December 13, 2006
 Published online: March 6, 2007

Keywords: cyclization · enantioselectivity · enynes · silver · stannanes

- [1] For reviews, see: a) G. C. Lloyd-Jones, *Org. Biomol. Chem.* **2003**, *1*, 215–236; b) C. Aubert, O. Buisine, M. Malacria, *Chem. Rev.* **2002**, *102*, 813–834; c) S. T. Diver, A. J. Giessert, *Chem. Rev.* **2004**, *104*, 1317–1382; d) A. M. Echavarren, C. Nevado, *Chem. Soc. Rev.* **2004**, *33*, 431–436; e) E. Jiménez-Núñez, A. M. Echavarren, *Chem. Commun.* **2006**, 333–346.
- [2] a) C. Fernández-Rivas, M. Méndez, A. M. Echavarren, *J. Am. Chem. Soc.* **2000**, *122*, 1221–1222; b) C. Fernández-Rivas, M. Méndez, C. Nieto-Oberhuber, A. M. Echavarren, *J. Org. Chem.* **2002**, *67*, 5197–5201.
- [3] M. Méndez, A. M. Echavarren, *Eur. J. Org. Chem.* **2002**, 15–28.
- [4] a) B. Martín-Matute, E. Buñuel, D. J. Cárdenas, M. Méndez, C. Nieto-Oberhuber, A. M. Echavarren, *J. Organomet. Chem.* **2003**, *687*, 410–419; b) S. Shin, T. V. RajanBabu, *J. Am. Chem. Soc.* **2001**, *123*, 8416–8417.
- [5] K. Miura, N. Fujisawa, H. Saito, H. Nishikori, A. Hosomi, *Chem. Lett.* **2002**, 32–33.
- [6] K. Miura, N. Fujisawa, A. Hosomi, *J. Org. Chem.* **2004**, *69*, 2427–2430.
- [7] The intramolecular addition of allylsilanes to alkynes promoted by indium(III) takes place by a mechanism similar to that proposed in Ref. [2]. See: K. Miura, N. Fujisawa, S. Toyohara, A. Hosomi, *Synlett* **2006**, 1883–1886.
- [8] For the cyclization of allyl bromides with alkynes catalyzed by InCl₃, that involves a halogen transfer, see: G. R. Cook, R. Hayashi, *Org. Lett.* **2006**, *8*, 1045–1048.
- [9] a) M. Méndez, M. P. Muñoz, A. M. Echavarren, *J. Am. Chem. Soc.* **2000**, *122*, 11549–11550; b) M. Méndez, M. P. Muñoz, C. Nevado, D. J. Cárdenas, A. M. Echavarren, *J. Am. Chem. Soc.* **2001**, *123*, 10511–10520.
- [10] a) C. Nieto-Oberhuber, M. P. Muñoz, E. Buñuel, C. Nevado, D. J. Cárdenas, A. M. Echavarren, *Angew. Chem.* **2004**, *116*, 2456–2460; *Angew. Chem. Int. Ed.* **2004**, *43*, 2402–2406; b) C. Nieto-Oberhuber, M. P. Muñoz, S. López, E. Jiménez-Núñez, C. Nevado, E. Herrero-Gómez, M. Raducan, A. M. Echavarren, *Chem. Eur. J.* **2006**, *12*, 1677–1693; c) C. Nieto-Oberhuber, S. López, M. P. Muñoz, D. J. Cárdenas, E. Buñuel, C. Nevado, A. M. Echavarren, *Angew. Chem.* **2005**, *117*, 6302–6304; *Angew. Chem. Int. Ed.* **2005**, *44*, 6146–6148; d) C. Nieto-Oberhuber, S. López, E. Jiménez-Núñez, A. M. Echavarren, *Chem. Eur. J.* **2006**, *12*, 5916–5923.
- [11] See, for example: a) D. J. Gorin, N. R. Davis, F. D. Toste, *J. Am. Chem. Soc.* **2005**, *127*, 11260–11261; b) J. J. Kennedy-Smith, S. T. Staben, F. D. Toste, *J. Am. Chem. Soc.* **2004**, *126*, 4526–4527; c) J. J. Kennedy-Smith, S. T. Staben, F. D. Toste, *J. Am. Chem. Soc.* **2004**, *126*, 4526–4527; d) B. D. Sherry, F. D. Toste, *J. Am. Chem. Soc.* **2004**, *126*, 15978–15979.
- [12] S. Ma, S. Yu, Z. Gu, *Angew. Chem.* **2006**, *118*, 206–209; *Angew. Chem. Int. Ed.* **2006**, *45*, 200–203.
- [13] For leading references of silver(I)-catalyzed cyclizations of alkynes, see: a) R. F. Sweis, M. P. Schramm, S. A. Kozmin, *J. Am. Chem. Soc.* **2004**, *126*, 7442–7443; b) T. J. Harrison, G. R. Dake, *Org. Lett.* **2004**, *6*, 5023–5026; c) T. J. Harrison, J. A. Kozak, M. Corbella-Pané, G. R. Dake, *J. Org. Chem.* **2006**, *71*, 4525–4529; d) J. Sun, S. A. Kozmin, *Angew. Chem.* **2006**, *118*, 5113–5115; *Angew. Chem. Int. Ed.* **2006**, *45*, 4991–4993; e) J. Zhao, C. O. Hughes, F. D. Toste, *J. Am. Chem. Soc.* **2006**, *128*, 7436–7437.
- [14] For leading references on silver(I)–carbene complexes, see: a) J. C. Garrison, W. J. Youngs, *Chem. Rev.* **2005**, *105*, 3978–4008; b) P. de Frémont, N. M. Scott, E. D. Stevens, T. Ramnial, O. C. Lightbody, C. L. B. Macdonald, J. A. C. Clyburne, C. D. Abernethy, S. P. Nolan, *Organometallics* **2005**, *24*, 6301–6309; c) R. R. Julian, J. A. May, B. M. Stoltz, J. L. Beauchamp, *J. Am. Chem. Soc.* **2003**, *125*, 4478–4486; d) A. J. Arduengo, H. V. R.

- Dias, J. C. Calabrese, F. Davidson, *Organometallics* **1993**, *12*, 3405–3409.
- [15] For discussions on π -back bonding in silver(I)–carbene complexes, see: a) X. Hu, I. Castro-Rodriguez, K. Olsen, K. Meyer, *Organometallics* **1993**, *12*, 3405–3409; b) D. Nemcsok, K. Wichmann, G. Frenking, *Organometallics* **2004**, *23*, 3640–3646.
- [16] a) M. Wadamoto, H. Yamamoto, *J. Am. Chem. Soc.* **2005**, *127*, 14556–14557; b) N. Momiyama, H. Yamamoto, *J. Am. Chem. Soc.* **2004**, *126*, 5360–5361; c) M. Wadamoto, N. Ozasa, A. Yanagisawa, H. Yamamoto, *J. Org. Chem.* **2003**, *68*, 5593–5601; d) A. Yanagisawa, H. Nakashima, A. Ishiba, H. Yamamoto, *J. Am. Chem. Soc.* **1996**, *118*, 4723–4724.
- [17] S. E. Denmark, J. Fu, *Chem. Rev.* **2003**, *103*, 2763–2793.
- [18] M. Bardaji, O. Crespo, A. Laguna, A. K. Fischer, *Inorg. Chim. Acta* **2000**, *304*, 7–16.
- [19] S. Antoniotti, E. Genin, V. Michelet, J.-P. Genêt, *J. Am. Chem. Soc.* **2005**, *127*, 9976–9977.
- [20] a) E. Herrero-Gómez, C. Nieto-Oberhuber, S. López, J. Benet-Buchholz, A. M. Echavarren, *Angew. Chem.* **2006**, *118*, 5581–5585; *Angew. Chem. Int. Ed.* **2006**, *45*, 5455–5459; b) CCDC-634526 contains the supplementary crystallographic data for complex **8**. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [21] C. Nieto-Oberhuber, S. López, A. M. Echavarren, *J. Am. Chem. Soc.* **2005**, *127*, 6178–6179.
- [22] For details, see the Supporting Information.
- [23] a) S. P. H. Mee, V. Lee, J. E. Baldwin, *Angew. Chem.* **2004**, *116*, 1152–1156; *Angew. Chem. Int. Ed.* **2004**, *43*, 1132–1136; b) S. P. H. Mee, V. Lee, J. E. Baldwin, *Chem. Eur. J.* **2005**, *11*, 3294–3308.
- [24] M. Yasuhisa, A. Naoki, K. Haruo, Yoshinori, Y. *Tetrahedron* **1999**, *55*, 3779–3790.
- [25] For enantioselective cyclizations of enynes initiated by the coordination of the metal to the alkyne function of enynes, see: platinum(III) catalysis: a) L. Charruault, V. Michelet, R. Taras, S. Gladialli, J.-P. Genêt, *Chem. Commun.* **2004**, 850–851; silver(I) catalysis: b) M. P. Muñoz, J. Adrio, J. C. Carretero, A. M. Echavarren, *Organometallics* **2005**, *24*, 1293–1300.
- [26] a) U. Létinois-Halbes, P. Pale, S. Berger, *J. Org. Chem.* **2005**, *70*, 9185–9190; b) A. Vitèrisi, A. Orsini, J.-M. Weibel, P. Pal, *Tetrahedron Lett.* **2006**, *47*, 2779–2781, and references therein.
- [27] For closely related examples of intramolecular cyclopropanation reactions catalyzed by ruthenium(II) or platinum(II), see: B. P. Peppers, S. T. Diver, *J. Am. Chem. Soc.* **2004**, *126*, 9524–9525.