

Synthesis of highly stable intermediates in Michael-type additions to the double bond in $(\text{SPhPh}_2)_2\text{C}=\text{CH}_2$

Eduardo J. Fernández,^a M. Concepción Gimeno,^b Peter G. Jones,^c Antonio Laguna,^{*,b} José M. López-de-Luzuriaga^a and Elena Olmos^a

^a Departamento de Química, Universidad de La Rioja, Obispo Bustamante 3, E-26001 Logroño, Spain

^b Departamento de Química Inorgánica, Instituto de Ciencia de Materiales de Aragón, Universidad de Zaragoza-CSIC, E-50009 Zaragoza, Spain

^c Institut für Anorganische und Analytische Chemie der Technischen Universität, Postfach 3329, D-38023 Braunschweig, Germany


Treatment of vinylidenebis(diphenylphosphine) with elemental sulfur under reflux led to vinylidenebis(diphenylphosphine) disulfide, $(\text{SPhPh}_2)_2\text{C}=\text{CH}_2$, in good yields. Its co-ordination to a gold(III) centre activates the carbon-carbon double bond. Thus, the chelate complex $[\text{Au}(\text{C}_6\text{F}_5)_2\{(\text{SPhPh}_2)_2\text{C}=\text{CH}_2\}]\text{ClO}_4$ reacted rapidly with carbon- or oxygen-donor nucleophiles (Nu^-), such as acac^- (acetylacetonate), CN^- , C_5H_5^- or OEt^- , forming methanide-type derivatives $[\text{Au}(\text{C}_6\text{F}_5)_2\{(\text{SPhPh}_2)_2\text{CCH}_2\text{Nu}\}]$. These products can be considered as intermediates in Michael-type additions of HNu to the $\text{C}=\text{C}$ bond, which in this particular case display high stabilities, allowing their isolation as solids. The crystal structure of the cyclopentadienyl derivative has been established by X-ray crystallography and displays a square-planar gold(III) centre with a *cis* disposition of the pentafluorophenyl groups and sulfur atoms of the chelating ligand, and sp^2 hybridization of the endocyclic carbon atom with partial $\text{P}-\text{C}$ double-bond character.

The co-ordination chemistry of vinylidenebis(diphenylphosphine)¹⁻⁴ and the reactivity of its carbon-carbon double bond in addition reactions have been studied in the last decade. In the course of these investigations it has been shown that, although the double bond in unco-ordinated vinylidenebis(diphenylphosphine) is scarcely susceptible to nucleophilic attack, except under extreme reaction conditions,⁴⁻⁶ complexation to a metal centre^{2a,3,4,7,8} or quaternization of the phosphorus atoms⁹ activates the double bond in such a manner that various nucleophiles can be added through Michael-type reactions. In these processes the species HNu is added to complexes $[\text{M}-(\text{PPh}_2)_2\text{C}=\text{CH}_2]$ to give the products $[\text{M}-(\text{PPh}_2)_2\text{CHCH}_2\text{Nu}]$, where the first step is always the attack of the nucleophile Nu^- on the C_β atom of the $\text{C}=\text{C}$ bond. However, the product formed in the first step had not been isolated.

We have recently reported that the reaction of $[\text{Au}(\text{C}_6\text{F}_5)_2\text{Cl}\{\text{PPh}_2\text{C}(\text{CH}_2)\text{PPh}_2\}]$ with nucleophilic agents Nu^- allows the isolation of these intermediates, affording methanide-like derivatives $[\text{Au}(\text{C}_6\text{F}_5)_2\{(\text{PPh}_2)_2\text{CCH}_2\text{Nu}\}]$.¹⁰ These complexes are structurally and electronically similar to the bis(diphenylphosphino)methanide derivative $[\text{Au}(\text{C}_6\text{F}_5)_2\{(\text{PPh}_2)_2\text{CH}\}]$, which is formed by extraction of one proton from $[\text{Au}(\text{C}_6\text{F}_5)_2\{(\text{PPh}_2)_2\text{CH}_2\}]\text{ClO}_4$ when treated with a nucleophile Nu^- (H^-); the excess electron density is delocalized over both $\text{P}-\text{C}$ bonds¹¹ (Scheme 1).

In contrast, it has been reported that when the same reaction is carried out with the bis(diphenylphosphino)methane disulfide ligand the initially formed six-membered ring is rapidly transformed into a four-membered ring through the co-ordination of the methanide carbon to the gold(III) centre¹² (see Scheme 2), and this can be considered as electrophilic attack of the gold(III) centre on the unsaturated carbon.

In order to investigate the stability of the intermediates in a Michael addition to vinylidenebis(diphenylphosphine) disulfide we have synthesized the chelate bis(pentafluorophenyl) complex $[\text{Au}(\text{C}_6\text{F}_5)_2\{(\text{SPhPh}_2)_2\text{C}=\text{CH}_2\}]\text{ClO}_4$ to study its reactivity with nucleophiles Nu^- . These reactions provide optimum conditions





Scheme 1 $\text{R} = \text{C}_6\text{F}_5$

for completion of the addition to the double bond, because the electrophile gold(III) is present in the same molecule as the carbanion; thus a similar process to that described for the $\text{Ph}_2\text{P}(\text{S})\text{CH}_2(\text{S})\text{PPh}_2$ derivative could be expected. Surprisingly, the products isolated in these reactions are always those formed in the first step, containing a six-membered ring and displaying both carbanion and electrophile together in the same molecule without any interaction between them (see Scheme 2). This fact reveals that these particular intermediates are not only isolable, but also even more stable than expected.

Results and Discussion

When vinylidenebis(diphenylphosphine) disulfide is treated with $[\{\text{Au}(\text{C}_6\text{F}_5)_2(\mu-\text{Cl})\}_2]$ in a 2 : 1 molar ratio one of the sulfur atoms co-ordinates to the gold centre, rupturing the chlorine bridges to give the monodentate $[\text{Au}(\text{C}_6\text{F}_5)_2\text{Cl}\{\text{SPhPh}_2\text{C}(\text{CH}_2)\text{PPh}_2\text{S}\}]$ **1** (Scheme 3) as a white solid soluble in common organic solvents. It presents a *cis* disposition of the pentafluorophenyl groups, as shown by its ^{19}F NMR spectrum, which shows two different types of C_6F_5 with relative intensities 1 : 1. Its infrared spectrum exhibits the $\nu(\text{Au}-\text{Cl})$ stretching

Scheme 2 R = C₆F₅

Scheme 3 R = C₆F₅. (i) $\frac{1}{2} [\{Au(C_6F_5)_2(\mu-Cl)\}_2]$; (ii) AgClO₄; (iii) [Au(C₆F₅)₂(OEt₂)₂]ClO₄; (iv) Tl(acac); (v) [Au(C₆F₅)₂(acac)]; (vi) KCN; (vii) TlC₅H₅; (viii) NaOEt

band at 334 cm⁻¹ and two ν(P=S) vibrations at 594 and 613 cm⁻¹, corresponding to the single and double bond, respectively (614 cm⁻¹ for the free diphosphine disulfide).

Complex **1** reacts with an equimolecular amount of AgClO₄ to give [Au(C₆F₅)₂{(SPPH₂)₂C=CH₂}]ClO₄ **2**. The same product can be prepared by a ligand-displacement reaction of [Au(C₆F₅)₂(OEt₂)₂]ClO₄ with (SPPH₂)₂C=CH₂, as represented in Scheme 3. Compound **2** is obtained in good yields as a white solid, soluble in chlorinated solvents and acetone and insoluble in diethyl ether and hexane.

As mentioned above, the co-ordination of vinylidenebis(diphenylphosphine) extensively activates its double bond^{2a,3,4,7,8} and the same effect is observed here for vinylidenebis(diphenylphosphine) disulfide. This activation occurs in such a manner that the nucleophilic addition in complex **2** takes place only to the C_β atom of the C=C bond, giving rise to derivatives [Au(C₆F₅)₂{(SPPH₂)₂CCH₂Nu}], which can be considered as the products formed in the first step of a normal Michael addition, displaying their striking high stability. Thus, the reaction between equimolecular amounts of **2** and Tl(acac) (acac = acetylacetonate) gives rise to the precipitation of TlClO₄ and addition of acetylacetonate to the C_β atom, affording the neutral derivative [Au(C₆F₅)₂{(SPPH₂)₂CCH₂CH(OMe)₂}] **3** (Scheme 3). This product can alternatively be prepared by treating (SPPH₂)₂C=CH₂ with [Au(C₆F₅)₂(acac)], which contains the nucleophile.

Carbon- or oxygen-donor nucleophiles can be employed in these reactions. Thus, treatment of complex **2** with QNu

(QNu = KCN, TlC₅H₅ or NaOEt) results in the formation of the corresponding QClO₄ and addition of Nu⁻ to the exocyclic carbon centre to give [Au(C₆F₅)₂{(SPPH₂)₂CCH₂Nu}] (Nu⁻ = CN⁻ **4**, C₅H₅⁻ **5**, or OEt⁻ **6**). Complexes **3–6** are air- and moisture-stable in the solid state and soluble in common organic solvents, except hexane. Their IR spectra show a band in the region 569–579 cm⁻¹ [ν(P=S)], and also bands at 1505m, 967m, 804m and 795m cm⁻¹, due to the presence of C₆F₅ groups linked to a gold(III) centre; the last two confirm the *cis* disposition.¹³

The presence of equivalent pentafluorophenyl groups is confirmed in the ¹⁹F NMR spectra (see Experimental section). The addition of the nucleophile to the C_β of the double bond leads to the presence of an excess of electron density at the C_α atom, which is delocalized through both P–C bonds. This is reflected in the ³¹P-{¹H} NMR spectra, in which the chemical shift for these phosphorus decreases from δ 41.7 in complex **2** to δ 36.0–38.0 for **3–6**. The breaking of the double bond also produces a change in both chemical shift and pattern corresponding to the protons of the CH₂ group in their NMR spectra from an AA'XX' spin system at δ 7.15 to a triplet located between δ 2.66 and 3.47.

The structure of complex **5** has been established by X-ray diffraction studies. The molecule is shown in Fig. 1 with selected bond lengths and angles in Table 1. The gold(III) centre has a planar geometry, being bonded to the two pentafluorophenyl groups and chelated by the two sulfur atoms of the diphosphine disulfide ligand. The major deviation of the angles from the ideal 90° is positive and is observed within the chelate ligand, S–Au–S 100.81(3)°, whereas the narrowest angles are




Fig. 1 Structure of complex **5** in the crystal. Displacement parameter ellipsoids represent 50% probability surfaces. Hydrogen atoms and solvent are omitted for clarity

Table 1 Selected bond lengths (Å) and angles (°) for complex **5**

Au–C(61)	2.032(3)	Au–C(51)	2.033(3)
Au–S(2)	2.3552(9)	Au–S(1)	2.3626(9)
P(1)–C(1)	1.709(3)	P(1)–C(11)	1.807(3)
P(1)–C(21)	1.813(4)	P(1)–S(1)	2.0459(11)
P(2)–C(1)	1.719(3)	P(2)–C(31)	1.802(3)
P(2)–C(41)	1.811(3)	P(2)–S(2)	2.0700(12)
C(1)–C(2)	1.538(4)	C(2)–C(3)	1.494(5)
C(3)–C(4)	1.312(6)	C(3)–C(7)	1.466(5)
C(4)–C(5)	1.460(7)	C(5)–C(6)	1.307(7)
C(6)–C(7)	1.455(6)		
C(61)–Au–C(51)	89.54(13)	C(61)–Au–S(2)	84.22(9)
C(51)–Au–S(2)	173.75(10)	C(61)–Au–S(1)	174.97(9)
C(51)–Au–S(1)	85.43(10)	S(2)–Au–S(1)	100.81(3)
C(1)–P(1)–C(11)	107.5(2)	C(1)–P(1)–C(21)	114.6(2)
C(11)–P(1)–C(21)	107.4(2)	C(1)–P(1)–S(1)	114.09(11)
C(11)–P(1)–S(1)	104.62(12)	C(21)–P(1)–S(1)	107.97(12)
C(1)–P(2)–C(31)	109.2(2)	C(1)–P(2)–C(41)	115.3(2)
C(31)–P(2)–C(41)	107.3(2)	C(1)–P(2)–S(2)	114.40(12)
C(31)–P(2)–S(2)	103.70(11)	C(41)–P(2)–S(2)	106.17(11)
P(1)–S(1)–Au	105.71(4)	P(2)–S(2)–Au	104.28(4)
C(2)–C(1)–P(1)	119.3(2)	C(2)–C(1)–P(2)	120.4(2)
P(1)–C(1)–P(2)	115.8(2)	C(3)–C(2)–C(1)	117.0(3)
C(4)–C(3)–C(2)	107.0(4)	C(4)–C(3)–C(7)	131.7(4)
C(7)–C(3)–C(2)	121.2(3)	C(3)–C(4)–C(5)	110.6(4)
C(6)–C(5)–C(4)	107.7(4)	C(5)–C(6)–C(7)	109.4(4)
C(6)–C(7)–C(3)	105.2(4)		

C–Au–S 85.43(10) and 84.22(9)°. The Au–C bond distances, 2.032(3) and 2.033(3) Å, are shorter than in other gold(III) complexes with diphosphine ligands, probably as a consequence of the higher *trans* influence of the phosphorus compared to the sulfur ligands. The Au–S distances are 2.3552(9) and 2.3626(9) Å, similar to others in gold(III) derivatives with similar ligands, e.g. [Au(C₆F₅)₂(SPPPh₂CHPPh₂)] [2.345(2) Å].¹⁴ The P(1)–C(1) and P(2)–C(1) bond lengths are 1.709(3) and 1.719(3) Å, shorter than those found in complexes with the diphosphine, such as [{AuCl(PPh₂C(=CH₂)PPh₂)]₂] [1.828(6) and 1.833(6) Å],⁴ or in other derivatives where a Michael addition has taken place, such as [PdI₂{(PPh₂)₂CHCH₂OCH₂CH₂(3-SC₄H₉)}] [1.84(2) and 1.90(1) Å].^{3d} This indicates that the negative charge of the methanide carbon is delocalized over the P–C bonds, conferring on them a degree of multiple bonding. The C(1)–C(2) distance, 1.538(4) Å, corresponds to a single bond. The C–C distances within the cyclopentadienyl ring allow an unambiguous assignment of single- and double-bond positions, and there is no indication of disorder.

Experimental

Instrumentation and materials

The IR spectra were recorded on a Perkin-Elmer 883 spectro-

photometer, over the range 4000–200 cm^{−1}, using Nujol mulls between polyethylene sheets. ¹H, ¹⁹F and ³¹P NMR spectra on a Bruker ARX 300 in CDCl₃ solutions, chemical shifts being quoted relative to SiMe₄ (¹H, external), CFCl₃ (¹⁹F, external) and H₃PO₄ (85%) (³¹P, external). The elemental analyses (C, H, N, S) were performed with a Perkin-Elmer 2400 microanalyzer. Mass spectra were recorded on a VG Autospec instrument using FAB techniques and 3-nitrobenzyl alcohol as matrix. All experiments were carried out at room temperature. The starting materials (PPh₂)₂C=CH₂,¹⁵ (SPPPh₂)₂C=CH₂,¹⁶ [{Au(C₆F₅)₂-(μ-Cl)]₂¹⁷ and [Au(C₆F₅)₂(acac)]¹⁸ were prepared according to the literature. The complex [Au(C₆F₅)₂(OEt₂)₂]ClO₄ was prepared *in situ* by reaction of [{Au(C₆F₅)₂(μ-Cl)]₂ with 2 equivalents of AgClO₄ in diethyl ether.

Syntheses

[Au(C₆F₅)₂CH{SPPPh₂C(=CH₂)PPh₂-S}] 1. To a dichloromethane solution (20 cm³) of [{Au(C₆F₅)₂(μ-Cl)]₂ (0.112 g, 0.1 mmol) was added (SPPPh₂)₂C=CH₂ (0.092 g, 0.2 mmol). After stirring for 4 h the solution was concentrated to ca. 5 cm³. Addition of hexane led to complex **1** as a white solid (0.168 g, 82%) (Found: C, 44.0; H, 1.7; S, 6.35. Calc. for C₃₈H₂₂AuClF₁₀P₂S₂: C, 44.45; H, 2.15; S, 6.25%). ³¹P-{¹H} NMR: δ 41.7 (s, 1P, Au–P–S) and 45.5 (s, 1P, P=S). ¹H NMR: δ 6.89 (AA'XX', 2 H, CH₂, *N* = 65.0 Hz). ¹⁹F NMR: δ −122.4 (m, *o*-F), −153.6 (t, *p*-F, *J*_{FF} = 20.0), −159.0 (m, *m*-F), −122.6 (m, *o*-F), −157.7 (t, *p*-F, *J*_{FF} = 20.2 Hz) and −161.9 (m, *m*-F). FAB mass spectrum: *m/z* = 992 (88, [M – Cl]⁺) and 658 (100%, [Au{(SPPPh₂)₂C=CH₂}]⁺).

[Au(C₆F₅)₂{(SPPPh₂)₂C=CH₂}]ClO₄ 2. *Method (a).* To a freshly prepared solution of [Au(C₆F₅)₂(OEt₂)₂]ClO₄ (0.2 mmol) in diethyl ether (20 cm³) was added (SPPPh₂)₂C=CH₂ (0.092 g, 0.2 mmol). After stirring for 3 h a white precipitate of complex **2** was filtered off (0.205 g, 94%).

Method (b). A dichloromethane solution (20 cm³) of complex **1** (0.103 g, 0.1 mmol) was treated with AgClO₄ (0.027 g, 0.1 mmol). After stirring for 3 h the AgCl formed was filtered off. Concentration of the solution to ca. 5 cm³ and addition of diethyl ether (20 cm³) gave complex **2** as a white solid (0.131 g, 60%) (Found: C, 41.65; H, 2.5; S, 5.95. Calc. for C₃₈H₂₂AuClF₁₀O₄P₂S₂: C, 41.85; H, 2.05; S, 5.9%). ³¹P-{¹H} NMR: δ 41.7 (s). ¹H NMR: δ 7.15 (AA'XX', 2 H, CH₂, *N* = 69.2 Hz). ¹⁹F NMR: δ −122.3 (m, *o*-F), −153.7 (t, *p*-F, *J*_{FF} = 20.0 Hz) and −159.1 (m, *m*-F). FAB mass spectrum: *m/z* = 992 (65, M⁺) and 658 (100%, [Au{(SPPPh₂)₂C=CH₂}]⁺).

[Au(C₆F₅)₂{(SPPPh₂)₂CCH₂CH(COMe)₂}] 3. *Method (a).* To a dichloromethane solution (20 cm³) of complex **2** (0.218 g, 0.2 mmol) was added Tl(acac) (0.060 g, 0.2 mmol). The reaction mixture was stirred under nitrogen for 3 h and the TlClO₄ formed was then filtered off. The resulting solution was concentrated in a vacuum and the addition of hexane (15 cm³) led to complex **3** as a yellow solid (0.120 g, 55%).

Method (b). To a solution of (SPPPh₂)₂C=CH₂ (0.092 g, 0.2 mmol) in dichloromethane (20 cm³) was added [Au(C₆F₅)₂-(acac)] (0.126 g, 0.2 mmol). After 1 h of stirring under nitrogen the solvent was evaporated and hexane was added to precipitate complex **3** as a white solid (0.153 g, 70%) (Found: C, 47.75; H, 2.7; S, 5.5. Calc. for C₄₃H₂₉AuF₁₀O₂P₂S₂: C, 47.35; H, 2.7; S, 5.9%). ³¹P-{¹H} NMR: δ 36.3 (s). ¹H NMR: δ 1.26 (s, 6 H, CH₃), 2.66 (dt, 2 H, CH₂, ³*J*_{HP} = 21.5, ³*J*_{HH} = 5.3 Hz) and 2.95 (t, 1 H, CH). ¹⁹F NMR: δ −121.9 (m, *o*-F), −157.1 (t, *p*-F, *J*_{FF} = 20.0 Hz) and −161.4 (m, *m*-F). FAB mass spectrum: *m/z* = 1091 (53, M⁺), 758 (82, [M – 2C₆F₅]⁺) and 658 (100% [Au{(SPPPh₂)₂C=CH₂}]⁺).

[Au(C₆F₅)₂{(SPPPh₂)₂CCH₂CN}] 4. To a suspension of complex **2** (0.218 g, 0.2 mmol) in methanol (10 cm³) and under nitrogen was added KCN (0.012 g, 0.2 mmol) and the mixture was stirred for 2 h. Evaporation of the solvent to dryness and

Table 2 Details of data collection and structure refinement for complex **5**

Chemical formula	C ₄₆ H ₃₄ AuF ₁₀ P ₂ S ₂
<i>M</i>	1099.76
Crystal habit	Irregular orange tablet
Crystal size/mm	0.70 × 0.60 × 0.30
Crystal system	Triclinic
Space group	<i>P</i> $\bar{1}$
<i>a</i> /Å	11.617(2)
<i>b</i> /Å	12.163(2)
<i>c</i> /Å	16.005(2)
α /°	86.313(8)
β /°	75.125(10)
γ /°	84.972(8)
<i>U</i> /Å ³	2175.3(6)
<i>Z</i>	2
<i>D</i> /Mg m ⁻³	1.679
<i>F</i> (000)	1082
<i>T</i> /°C	−100
2 θ_{\max} /°	55
μ (Mo-K α)/mm ⁻¹	3.627
Transmission factor range	0.625–1.0
No. reflections measured	12 427
No. unique reflections	9876
<i>R</i> _{int}	0.035
<i>R</i> ^a [<i>F</i> , <i>F</i> > 4 σ (<i>F</i>)]	0.028
<i>wR</i> ^b (<i>F</i> ² , all reflections)	0.065
No. parameters	535
No. restraints	481
<i>S</i> ^c	1.008
Maximum $\Delta\rho$ /e Å ⁻³	0.91

^a $R = \sum ||F_o| - |F_c|| / \sum |F_o|$. ^b $wR = [\sum w(F_o^2 - F_c^2)^2 / \sum w(F_o^2)^2]^{1/2}$; $w^{-1} = \sigma^2(F_o^2) + (aP)^2 + bP$, where $P = (F_o^2 + 2F_c^2)/3$ and *a* and *b* are constants adjusted by the program. ^c $S = [\sum w(F_o^2 - F_c^2)^2 / (n - p)]^{1/2}$, where *n* is the number of data and *p* the number of parameters.

addition of diethyl ether led to precipitation of the KClO₄ formed in the reaction, which was filtered off over Celite. The solution was then concentrated in a vacuum to ca. 5 cm³ and hexane (15 cm³) were added to precipitate complex **4** as a crystalline yellow solid (0.102 g, 50%) (Found: C, 46.55; H, 2.25; N, 1.2; S, 6.0. Calc. for C₃₉H₂₂AuF₁₀NP₂S₂: C, 46.05; H, 2.2; N, 1.4; S, 6.3%). ³¹P-{¹H} NMR: δ 37.7 (s). ¹H NMR: δ 2.76 (t, 2 H, CH₂, ³*J*_{HP} = 19.2 Hz). ¹⁹F NMR: δ −121.9 (m, *o*-F), −156.7 (t, *p*-F, *J*_{FF} = 20.0 Hz) and −161.1 (m, *m*-F). FAB mass spectrum: *m/z* = 1019 (65, [*M* + H]⁺), 684 (100, [*M* − 2C₆F₅]⁺) and 658 (55%, [Au{(SPPH₂)₂C=CH₂}]⁺).

[Au(C₆F₅)₂{(SPPH₂)₂CCH₂(C₅H₉)}]·0.5C₆H₁₄ **5**. To a dichloromethane solution (20 cm³) of complex **2** (0.218 g, 0.2 mmol) under a nitrogen atmosphere was added TiCl₃H₅ (0.054 g, 0.2 mmol). After stirring for 3 h the TiClO₄ was filtered off and the solution was concentrated to ca. 5 cm³. Addition of hexane led to complex **5** as an orange solid (0.09 g, 41%) (Found: C, 50.05; H, 3.15; S, 5.75. Calc. for C₄₆H₃₄AuF₁₀P₂S₂: C, 50.25; H, 3.1; S, 5.85%). ³¹P-{¹H} NMR: δ 36.0 (s). ¹H NMR: δ 1.99 (s, 2 H, CH=CCCH₂CH=CH), 3.06 (t, 2 H, P₂CCH₂, ³*J*_{HP} = 21.4 Hz), 5.12, 5.92, 6.01 (ABC, 3 H, CH=CCCH₂CH=CH). ¹⁹F NMR: δ −121.8 (m, *o*-F), −157.3 (t, *p*-F, *J*_{FF} = 20.1 Hz) and −161.5 (m, *m*-F). FAB mass spectrum: *m/z* = 1057 (100, *M*⁺), 723 (90, [*M* − 2C₆F₅]⁺) and 658 (98%, [Au{(SPPH₂)₂C=CH₂}]⁺).

[Au(C₆F₅)₂{(SPPH₂)₂CCH₂OEt}] **6**. To a dichloromethane solution (20 cm³) of complex **2** (0.218 g, 0.2 mmol) under nitrogen was added a freshly prepared solution of NaOEt (0.2 mmol). After stirring for 3 h the NaClO₄ was filtered off and the solution was concentrated to ca. 5 cm³. Addition of hexane led to the precipitation of complex **6** as a yellow solid (0.068 g, 33%) (Found: C, 46.8; H, 2.85; S, 6.15. Calc. for C₄₀H₂₇AuF₁₀OP₂S₂: C, 46.35; H, 2.6; S, 6.2%). ³¹P-{¹H} NMR: δ 38.0 (s). ¹H NMR: δ 0.82 (t, 3 H, OCH₂CH₃, ³*J*_{HH} = 7.0), 2.67 (q, 2 H, OCH₂CH₃) and 3.47 (t, 2 H, P₂CCH₂, ³*J*_{HP} = 24.7 Hz). ¹⁹F

NMR: δ −121.8 (m, *o*-F), −157.3 (t, *p*-F, *J*_{FF} = 19.9 Hz) and −161.4 (m, *m*-F). FAB mass spectrum: *m/z* = 991 (50, [*M* − OEt]⁺) and 658 (100%, [Au{(SPPH₂)₂C=CH₂}]⁺).

Crystallography

The crystal of complex **5** was mounted in inert oil on a glass fibre. Data were collected using monochromated Mo-K α radiation (λ = 0.710 73 Å). Diffractometer type: Siemens P4 equipped with an LT-2 low-temperature attachment. Scan type ω . Cell constants were refined from setting angles of 65 reflections in the range 2θ = 6–25°. Absorption corrections were applied on the basis of ψ scans. The structure was solved by the heavy-atom method and refined on *F*² (program SHELXL 93).¹⁹ The complex crystallizes with half a molecule of hexane per asymmetric unit. All non-hydrogen atoms were refined anisotropically, with the exception of the solvent carbon atoms. Hydrogen atoms were included using a riding model. Further details are given in Table 2.

CCDC reference number 186/635.

Acknowledgements

We thank the Dirección General de Investigación Científica y Técnica (no. PB94-0079) and the Fonds der Chemischen Industrie for financial support.

References

- P. A. Dolby, M. M. Harding, N. Newar and A. K. Smith, *J. Chem. Soc., Dalton Trans.*, 1992, 2939.
- (a) G. R. Cooper, D. M. McEwan and B. L. Shaw, *Inorg. Chim. Acta*, 1983, **76**, L165; (b) G. R. Cooper, F. S. M. Hassan, B. L. Shaw and M. Thornton-Pett, *J. Chem. Soc., Chem. Commun.*, 1985, 614; (c) G. R. Cooper, D. M. McEwan and B. L. Shaw, *Inorg. Chim. Acta*, 1986, **122**, 207.
- (a) F. S. M. Hassan, S. J. Higgins, G. B. Jacobsen, B. L. Shaw and M. Thornton-Pett, *J. Chem. Soc., Dalton Trans.*, 1988, 3011; (b) S. J. Higgins and B. L. Shaw, *J. Chem. Soc., Dalton Trans.*, 1989, 1527; (c) A. M. Herring, S. J. Higgins, G. B. Jacobsen and B. L. Shaw, *J. Chem. Soc., Chem. Commun.*, 1986, 882; (d) G. King, S. J. Higgins and A. Hopton, *J. Chem. Soc., Dalton Trans.*, 1992, 3403.
- H. Schmidbaur, R. Herr, G. Müller and J. Riede, *Organometallics*, 1985, **4**, 1208.
- J. L. Bookhan, W. McFarlane and J. Colquhoun, *J. Chem. Soc., Chem. Commun.*, 1986, 1041; J. L. Bookhan, W. McFarlane and M. Thornton-Pett, *J. Chem. Soc., Dalton Trans.*, 1992, 2353.
- H. Schmidbaur, R. Herr and J. Riede, *Angew. Chem., Int. Ed. Engl.*, 1984, **23**, 247.
- F. S. M. Hassan, B. L. Shaw and M. Thornton-Pett, *J. Chem. Soc., Dalton Trans.*, 1988, 89.
- X. L. R. Fontaine, F. S. M. Hassan, S. J. Higgins, G. B. Jacobsen, B. L. Shaw and M. Thornton-Pett, *J. Chem. Soc., Chem. Commun.*, 1985, 1635.
- H. Schmidbaur, R. Herr, Th. Pollok, A. Schier, G. Müller and J. Riede, *Chem. Ber.*, 1985, **118**, 3105.
- E. J. Fernández, M. C. Gimeno, P. G. Jones, A. Laguna and M. E. Olmos, *Organometallics*, 1997, **16**, 1130.
- R. Usón, A. Laguna, M. Laguna, B. R. Manzano, P. G. Jones and G. M. Sheldrick, *J. Chem. Soc., Dalton Trans.*, 1984, 839.
- A. Laguna, M. Laguna, A. Rojo and M. N. Fraile, *J. Organomet. Chem.*, 1986, **315**, 269.
- R. Usón, A. Laguna, J. García and M. Laguna, *Inorg. Chim. Acta*, 1979, **37**, 201.
- R. Usón, A. Laguna, M. Laguna, M. N. Fraile, P. G. Jones and C. F. Erbrügger, *J. Chem. Soc., Dalton Trans.*, 1989, 73.
- I. J. Colquhoun and W. McFarlane, *J. Chem. Soc., Dalton Trans.*, 1982, 1915; H. Schmidbaur and Th. Pollok, *Helv. Chim. Acta*, 1984, **67**, 2175.
- H. Schmidbaur, R. Herr and G. Müller, *Z. Chem.*, 1984, **24**, 378.
- R. Usón, A. Laguna, M. Laguna and J. A. Abad, *J. Organomet. Chem.*, 1983, **249**, 437.
- R. Usón, A. Laguna and M. L. Arrese, *Synth. React. Inorg. Metal-Organ. Chem.*, 1984, **14**, 557.
- G. M. Sheldrick, SHELXL 93, A program for crystal structure refinement, University of Göttingen, 1993.

Received 15th May 1997; Paper 7/03357E