

Enhancement of *threo*-Selectivity in the Reaction of But-2-enyl-lithium with Aldehydes *via* Allylic Boronate Complexes


By YOSHINORI YAMAMOTO,* HIDETAKA YATAGAI, and KAZUHIRO MARUYAMA

(Department of Chemistry, Faculty of Science, Kyoto University, Kyoto 606, Japan)

Summary The *threo*-selectivity of the reaction of but-2-enyl-lithium with aldehydes in the presence of certain trialkylboranes is enhanced, the corresponding allylic boronate complexes are presumably involved as intermediates.

RECENTLY, attention has been focused on the diastereoselective synthesis of β -hydroxylcarbonyl-compounds which may be of use in the synthesis of macrolide antibiotics¹ The stereoselective cross-aldol condensation is a very useful direct method for this purpose The indirect method, whereby


allylic organometallic compounds are treated with aldehydes followed by a Wacker-type oxidation, has recently been introduced by us² and others³ (Scheme 1). Notably, a



SCHEME 1. M' = Al, B, Li, Mg, Si, Sn, Zn, Zr. M = Al, B, Li, Mg, Si, Sn. i, Wacker oxidation.

completely *erythro*-selective and regioselective cross-aldol condensation is achieved *via* the (*Z*)-alken-2-yltin route.² Unfortunately, however, *threo*-selectivity is quite low or even negligible when but-2-enyl-lithium or -magnesium derivatives are used. We report that the *threo*-selectivity in the reaction of but-2-enyl-lithium (1) with aldehydes is remarkably enhanced *via* allylic boronate complexes (Scheme 2). The results are summarized in the Table.

sociation of the complex into (1) or exert an influence on the rigidity of the cyclic transition-state.⁵ The importance of the metal ions is indicated by the reaction of but-2-enyl magnesium chloride (reactions 2 and 5).



SCHEME 2.

The reaction presumably proceeds *via* the allylic boronate complexes (2).⁶ Migration of the R' group is not detected and BR'₃ is recovered after completion of the reaction. The coupling always occurs at the γ -position. Thus, the behaviour of (2) resembles that of the but-2-enyl magnesium

TABLE. Enhancement of *threo*-selectivity in the reaction of (1) with aldehydes in the presence of trialkylboranes.^a

Reaction	Aldehyde	Additive	Solvent	Product ratio/% ^b		Total yield/% ^c
				<i>threo</i> [(3)]	<i>erythro</i>	
1	PhCHO	None	Ether	50	50	52 ^d
2	PhCHO ^e	None	Ether	50	50	(92)
3	PhCHO	Et ₃ Al	Ether	56	44	93
4	PhCHO	Et ₃ B	Ether	82	18	90
5	PhCHO ^e	Et ₃ B	Ether	50	50	95
6	PhCHO	Et ₃ B	Tetrahydrofuran	83	17	91
7	PhCHO	Et ₃ B	Ether-HMPA ^f	83	17	83
8	PhCHO	Bu ⁿ ₃ B	Ether	83	17	89
9	PhCHO	Bu ^s ₃ B	Ether	61	39	85
10	PhCHO	1-Bu ⁿ -3,6-Me ₂ -boraepan	Ether	80	20	85
11	MeCHO	Et ₃ B	Ether	85	15	(78)
12	Me ₂ CHCHO	Et ₃ B	Ether	68	32	(82)

^a All reactions were carried out on a 1 mmol scale. An equivalent amount of the Lewis acids (boranes or alane) was added to a solution of (1) at -70°C under N₂ and then the aldehydes were added. After 1 h, the reaction was quenched with methanol at -70°C . The products were identified by comparison with authentic materials. 'None' indicates the normal reaction without the additives. ^b Determined by ¹H n.m.r. spectra for reactions 1–10 and by g.l.c. (CW 6000, 2m) for reactions 11 and 12. ^c G.l.c. yield (isolated yield). ^d The straight-chain product arising from the coupling at the α position of (1) was also detected. ^e But-2-enyl magnesium chloride was used instead of but-2-enyl-lithium. ^f HMPA = hexamethylphosphoric triamide.

As apparent, the condensation with benzaldehyde in the presence of triethylborane, tri-*n*-butylborane, or 1-*n*-butyl-3,6-dimethylboraepan produces predominantly the *threo*-derivative (reactions 4, 6–8, and 10). Interestingly, the very sterically hindered tri-*s*-butylborane is not so effective, presumably because of the formation of a weak complex with (1) (reaction 9). The lack of effectiveness of Et₃Al may also be ascribed to a weak complex, since the longer C–Al bond and its more ionic nature⁴ should facilitate the dis-

derivative except for its high *threo*-selectivity.[†] Perhaps the most remarkable feature of the present procedure for the stereochemical control is its simplicity; merely the addition of BR'₃ before addition of the aldehydes produces the *threo*-selectivity. Furthermore, the present development suggests that trialkylboranes may be useful for the stereochemical control of certain carbanions.

(Received, 4th August 1980; Com. 853.)

[†] The *threo*-selectivity of trivalent but-2-enyl boranes, such as 9-but-2-enyl-9-borabicyclo[3.3.1]nonene, towards aldehydes is *ca.* 60%.

¹ P. A. Bartlett, *Tetrahedron*, 1980, **36**, 3.

² H. Yatagai, Y. Yamamoto, and K. Maruyama, *J. Am. Chem. Soc.*, 1980, **102**, 4548.

³ R. W. Hoffman and W. Ladner, *Tetrahedron Lett.*, 1979, 4653; D. B. Collum, J. H. McDonald III, and W. C. Still, *J. Am. Chem. Soc.*, 1980, **102**, 2118.

⁴ Y. Yamamoto, H. Yatagai, and K. Maruyama, *J. Org. Chem.*, 1980, **45**, 195.

⁵ D. A. Evans, E. Vogel, and J. V. Nelson, *J. Am. Chem. Soc.*, 1979, **101**, 6120.

⁶ Y. Yamamoto and K. Maruyama, *J. Am. Chem. Soc.*, 1978, **100**, 6282.