This article was downloaded by: [McMaster University] On: 22 October 2014, At: 06:45 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/gpss20

SYNTHESIS AND SPECTROSCOPIC CHARACTERIZATION OF HETEROCYCLIC DITHIOCARBAMATE COMPLEXES OF TRIORGANOTIN (IV)

Jyoti Sharma^a, Yashpal Singh^a & Audhesh Kumar Rai^a ^a Department of Chemistry, University of Rajasthan, Jaipur, 302 004, India Published online: 04 Oct 2006.

To cite this article: Jyoti Sharma , Yashpal Singh & Audhesh Kumar Rai (1996) SYNTHESIS AND SPECTROSCOPIC CHARACTERIZATION OF HETEROCYCLIC DITHIOCARBAMATE COMPLEXES OF TRIORGANOTIN (IV), Phosphorus, Sulfur, and Silicon and the Related Elements, 112:1-4, 19-26, DOI: 10.1080/10426509608046344

To link to this article: http://dx.doi.org/10.1080/10426509608046344

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

Phosphorus, Sulfur, and Silicon, 1996, Vol. 112, pp. 19-26 Reprints available directly from the publisher Photocopying permitted by license only © 1996 OPA (Overseas Publishers Association) Amsterdam B.V. Published in The Netherlands under license by Gordon and Breach Science Publishers SA Printed in Malaysia

SYNTHESIS AND SPECTROSCOPIC CHARACTERIZATION OF HETEROCYCLIC DITHIOCARBAMATE COMPLEXES OF TRIORGANOTIN (IV)

JYOTI SHARMA, YASHPAL SINGH and AUDHESH KUMAR RAI*

Department of Chemistry, University of Rajasthan, Jaipur 302 004, India

(Received March 7, 1995; in final form July 22, 1995)

Heterocyclic dithiocarbamate complexes of triorganotin (IV) having general formula,

 $CH_2CH_2XCH_2CH_2NCS_2SnR_3$ (where X = >CH₂, >CH—CH₃, >N—CH₃, >O; R = —CH₃ and —C₆H₃)

have been synthesized by the reaction of triorganotin chloride with the sodium salt of the corresponding ligand in 1:1 stoichiometry. These complexes have been characterized by elemental analysis, molecular weight measurements, IR and NMR (¹H, ¹³C and ¹¹⁹Sn) spectroscopy, suggesting that the methyl tin complexes are probably four coordinate, while the phenyl tin ones are probably five-coordinate with bidentate dithiocarbamate.

INTRODUCTION

The geometry and coordination number of the central tin atom in triorganotin (IV) derivatives of various ligands can mainly depend on the nature of organo groups attached to the central tin atom.^{1,2} Studies on $R_nSn(S_2CNR'R'')_{4-n}$ reveal that the dithiocarbamate ring can act both as monodentate³ ($R = -C_6H_5$, $n-C_4H_9$ and $-C_6H_5CH_2$; n = 3), bidentate^{4,5} ($R = CH_3$, $n-C_4H_9$, $-C_6H_5$; n = 1 or 2) and chelating uninegative⁶ ($R = -CH_3$; n = 3) ligand. Although the bidentate^{4,5} as well as chelating uninegative⁶ nature of dithiocarbamates are well established, the reported monodentate behaviour of dithiocarbamates is yet to be confirmed. Recently, ¹J (¹¹⁹Sn, ¹³C) coupling constants^{4,5,7} along with the ¹¹⁹Sn NMR data have been used to elucidate the structures of a variety of organotin (IV) compounds,⁸ specially those of diorganotin dithiocarbamates.⁹ No such report is yet available on the triorganotin (IV) derivatives of heterocyclic dithiocarbamate complexes. We now report the synthesis and characterization of triorganotin (IV) derivatives of some heterocyclic dithiocarbamates.

RESULTS AND DISCUSSION

The reactions of triorganotin chloride with the sodium salt of heterocyclic dithiocarbamates in 1:1 molar ratio in refluxing benzene solution yield the corresponding triorganotin (IV) derivatives.

CH₂CH₂XCH₂CH₂NCS₂Na

+
$$R_3SnCl \xrightarrow{\text{benzene}} CH_2CH_2XCH_2CH_2NCS_2SnR_3 + NaCl \downarrow$$

(where X = >CH₂, >CH-CH₃, >N-CH₃, >O; R = --CH₃ and --C₆H₅)

Complexes where $R = --CH_3$ are viscous liquids and those having $R = --C_6H_5$ are crystalline solids with sharp melting points. Both are soluble in common organic solvents like benzene, chloroform, dichloromethane and CS₂. Vapour pressure os-mometric molecular weight measurements in chloroform solution at 45°C confirm their monomeric composition.

IR Spectra

The IR spectra of these derivatives exhibit a strong band in the region 1470 ± 30 cm⁻¹ which may be attributed to the $\nu(C - N)$ mode. The presence of one intense absorption band at 1000 ± 10 cm⁻¹ for $\nu(C - S)$ mode, with a weak shoulder at 1010 ± 20 cm⁻¹ reveals that the dithiocarbamate ligand are monodentate³ moieties in trimethyltin(IV) complexes. However, the appearance of only one strong band at 1000 ± 10 cm⁻¹, without a shoulder in the triphenyltin(IV) analogs indicates the bidentate behaviour by the ligand. Analysis of the position of (C - N) mode further support such behaviour of ligands. This band, in fact, undergoes a shift towards a lower region or remains unchanged,¹⁰ while for bidentate coordination the frequency of their stretching mode would undergo upward shift when compared with its position in the corresponding dithiocarbamate sodium salt. The absorption band observed at 370 ± 30 cm⁻¹ has been assigned to ν Sn—S¹¹ mode. The bands observed at 520 ± 10 cm⁻¹ and 560 ± 30 cm⁻¹ are due to $\nu_{sym}(Sn-C)$ and $\nu_{asym}(Sn-C)$, respectively.^{12,13}

NMR Spectra

The ¹H NMR spectra of the complexes (Tables I) display the characteristic signals due to —CH₃, ring-CH₂, >N-CH₃, >CH-CH₃, $O_{\backslash CH_2}^{/CH_2}$, —N $_{\backslash CH_2}^{/CH_2}$, protons. A singlet at δ 0.56 to 0.67 ppm in trimethyltin(IV) complexes may be assigned to methyl protons attached to tin. The observed tin proton coupling constants ²J (¹¹⁹Sn, ¹H) are about 28.12 Hz. This suggests that the tin atom in these complexes is four-coordinate.¹⁴ A correlation between ²J (¹¹⁹Sn, ¹H) values and C—Sn—C bond angle θ is given by the relation⁷:

$$\theta = 0.0161[^{2}J]^{2} - 1.32[^{2}J] + 133.4$$

Substituting the relevant coupling constant data in the above equation, the C—Sn —C angles are 109°, indicating a tetrahedral environment around the central tin atom. Phenyl protons are observed as a complex multiplet in the range δ 7.23–7.97 ppm, and no ¹¹⁹Sn-¹H coupling was observed.

A comparison of the ¹³C NMR spectra of the ligands¹⁵ with the corresponding triphenyltin(IV) complexes (Tables II) shows an upfield shift in the position of CS_2 carbon signal indicating the chelating tendency of the dithiocarbamate moiety. How-

Downloaded by [McMaster University] at 06:45 22 October 2014

TABLE I

¹H NMR spectral data of heterocyclic dithiocarbamate complexes of triorganotin(IV) (values indicated in the table represents the protons in the bold letters) (δ) ppm

				•	•					
Complex	-CH,	CH ₂ CH ₂ CH ₂	>CH-Me	Me-CH ^{CH7} CH7	Mc—NCH ⁷ CH ⁷	$c_{-N} \stackrel{CH_{\overline{2}}}{\swarrow} CH_{\overline{2}}$	0 CH₂	Sn—R	² <i>J</i> (¹¹⁹ Sn, ¹ H)	Me—Sn—Me Angle <i>θ</i> °
(CH ₁) ₃ Sn(Pipdtc)		1.69(s)				4.15(t)		0.62(s)	28.12	109
(CH ₃),Sn(4-MePipdtc)	0.96(d) (3H)	(H0)	0.96–1.68(m) (1H)	3.13(t)(4H)		(4H) 5.22(d) (AH)	I	(9H) 0.56(s) (9H)	l	ļ
(CH ₃) ₃ Sn(N-MePzdtc)	() 2.33(s)	Ι	ĴI	Ι	2.49(t)(4H)	4.18(t)	I	0.56(s)	Į	l
(CH ₃) ₃ Sn(Morphdtc)	(HS)	ļ	ł	ļ	I	(4H) 3.84(t)	4.24(t)	(9H) 0.67(s)	28.12	601
(C ₆ H ₅) ₃ Sn(Pipdtc)	[1.68(s)	ł	I	I	(4H) 4.02(d)	(4H) 	(9H) 7.31–7.95(m)	l	ι
(C,H ₅) ₃ Sn(4-MePipdtc)	(b)96(d) /3H)	(H0)	1.20-2.25(m)	3.13(t)(4H)	ł	(4H) 4.90(d)	I	(HCI) 7.23-7.87(m) (H21)	ļ	l
(C ₆ H ₅) ₃ Sn(N-MePzdtc)	(JU) 2.25(s)	ŀ		ł	2.41(t)(4H)	(411) 4.10(t)	-	(1101) 7.23–7.87(m)	ł	ļ
(C ₆ H ₅) ₃ Sn(Morphdtc)	(uc)	I	ł	I	1	(4n) 2.41(t) (4H)	4.10(t) (4H)	(11C1) (m)7.31 – 7.87(m) (H51)	l	l
CH ₂ CH ₂ XCH ₂ CH ₂ N	CS ₂ SnR ₃ :	where X = >	CH ₂ (Pipdtc); >CF	H-CH ₃ (4-MePipd	tc); >N—CH ₃ (N-	MePzdtc); >O(N	Aorphdtc), 1	espectively.		

TRIORGANOTIN (IV) COMPLEXES

~+
-
0
C)
ч
Ō
<u> </u>
3
Ö
0
5
8
C 4
Ń.
ন
Ö
Ō
÷
а
_
Ŀ
÷
5
ē.
.≥
Ξ
5
-
5
ž
ä
÷
4
<u> </u>
Σ
>
ف.
ă
Ť
ē
2
D_
No.
5
ň.

:				TAB	LE II					
¹³ C and ¹¹ Sn NMR	k spectral	data of	heteroc	yclic dit	hiocarba	amate complexes o	f triorganotin(IV) (ð) ppm		
Compounds	C.	ů	c	Сď	C	Sn-R	'J(¹¹⁹ Sn, ¹³ Cl) Hz	MeSnMe $\theta(^{\circ}C)$ Angle	CN†	nS ⁰¹¹
d c b a CH ₂ CH ₂ (CH ₂)CH ₂ CH ₂ NCS ₂ Sn(CH ₃),	211.08	52.60	25.35	23.24	I	-1.078	373.89	106.66	4	19.74
d c b a CH2CH2(CH2)CH2CH2NCS5Sn(C6H5)	194.21	53.85	25.62	22.97	I	142.58; 136.57*; 128.34; 128.93	1	1	I	(-)190.89
d e b a CH2CH2(CH-CH3)CH2CH2NCS2Sn(CH3)	211.10	51.62	33.26	29.52	20.75	-1.241	372.34	106.53	4	17.39
d c b a CH ₂ CH ₂ (CH—CH ₃)CH ₂ CH ₂ NCS ₂ Sn(C ₆ H ₃)	194.48	53.20	33.69	29.63	21.02	142.64;136,62 * 128.39;128.93	1	ļ	1	(–)190.60
CH ₂ CH ₃ (N—CH ₃)CH ₂ CH ₂ NCS ₂ Sn(CH ₃),	212.20	44.80	50.49	l	53.52	-1.403	375.02	106.75	4	22.65
CH2CH2(N-CH3)CH2CH2NCS2Sn(C6H3)	195.89	45.34	52.06	l	54.06	142.10;136.57 128.39;129.04	1	I	I	(-)175.54
$\left \begin{array}{c} c & b \\ cH_2CH_2(O)CH_2CH_2NCS_2Sn(CH_3)_{3} \end{array} \right $	212.32	65.87	51.30	l	Ι	-1.132	373.64	106.64	4	34.48
CH ₂ CH ₂ (O)CH ₂ CH ₂ NCS ₂ Sn(C ₄ H ₅)	196.54	65.93	52.11	l	I	141.83;136.57* 128.50;129.15	ł			(-)181.83

J. SHARMA, Y. SINGH and A. K. RAI

*Tin-phenyl values are given in the order C(i), C(o), C(m) and C(p), respectively. \dagger Coordination number.

22



ever, in trimethyltin(IV) complexes this signal does not show any shift in its position when compared with the spectra of the corresponding ligands. This may be interpretated for the monodentate behaviour of ligand. In trimethyltin(IV) complexes, a singlet for methyl carbon occurred in the range δ 1.07–1.40 ppm. The tin-carbon coupling constants, ¹J (¹¹⁹Sn, ¹³C), were observed in the range of 372–375 Hz, which indicates that the central tin atom is four-coordinate.¹⁶ A correlation between ¹J (¹¹⁹Sn, ¹³C) values and C—Sn—C bond angle (θ) in trimethyltin(IV) complexes is given by⁷:

$$^{1}J = 11.4\theta - 875$$

or

$$\theta = 0.0877[^{1}J] + 76.7543$$

Substituting the coupling constant data in the above equation, the C—Sn—C angles are found to be in the range of $106.5^{\circ}-106.8^{\circ}$, suggesting a distorted tetrahedral environment around the tin atom.⁷

¹¹⁹Sn chemical shifts for trimethyltin(IV) complexes are in the range δ (-) 17– 35 ppm (Table II). The appearance of a ¹¹⁹Sn signal in this region is consistent with tetrahedral bonding⁸ around the central tin atom (Figure 1).

In all the organotin(IV) dithiocarbamates the geometries are distorted. In diorganotin(IV) bisdithiocarbamates, the ligand behave as bidentate moiety and Me—Sn —Me bond angle in trans geometry are found^{4.5} to be in the range of $136-144^{\circ}$. The two modifications of Me₃Sn S₂CNMe₂ show the average Me—Sn—Me bond angles of 111° (orthorhombic)¹⁷ and 113.7° (monoclinic)⁶ in distorted tetrahedral geometry. The behaviour of ligands may be is best described as chelating uninegative



(where $X = >CH_2$, $>CH-CH_3$, $>N-CH_3$, >O)

FIGURE 2

4
Ξ
0
\sim
H
ğ
2
3
Ö
\circ
<u> </u>
0
\sim
5
÷.
9
0
ц
co
_
\mathbf{E}
·Ħ
S
Ð
>
· =
T.
ъ
e
st
a
Z
÷.
Ц
\geq
>
6.
Ŗ
Fe Fe
2
č
ĭ
ц
WD
own
Jown

TABLE III Analytical and physical data of triorganotin(IV) complexes

Reactants in g		Molecular Formula	NaCl g	M.P.	Analy	sis	MW	V(C N)	V(C S)
NaS ₁ CNCH ₁ CH ₁ X CH ₁ CH ₁	R _, SnCl	(% Yield)	Found (Calc.)	(±1°C)	Sn% Found (Calc.)	S% Found (Calc.)	Found (Calc.)	cm. ¹	CHI-
X=>CH ₂ 2.22	R=-CH ₃ 2.02	C ₉ H ₉ NS ₂ Sn (92)	0.58 (0.59)		36.00 (36.62)	19.32 (19.78)	322.12 (324.06)	1450	1010 1020
X = >CH-CH, 2.32	R = -CH ₃ 2.10	C ₁₀ H ₃₁ NS ₂ Sn (94)	0.60 (0.61)	ı	35.08 (35.10)	18.52 (18.96)	335.42 (338.09)	1470	990 1010
X = >N-CH ₃ 2.09	R = -CH ₃ 1.93	C ₉ H ₂₀ N ₂ S ₂ Sn (85)	0.54 (0.56)		34 96 (35.00)	18.62 (18.90)	338.74 (339.08)	1440	1000 980
X = >O 2.06	R = -CH ₃ 1.86	C _e H ₁ ,NOS ₂ Sn (90)	0.53 (0.54)		36.32 (36.40)	19.28 (19.66)	325.92 (326.03)	1470	1010 1000
X = >CH ₁ 1.29	R = -C ₆ H ₅ 2.27	C ₂₄ H ₂₅ NS ₂ Sn (92)	0.33 (0.34)	150	23.20 (23.26)	12.38 (12.56)	510 (510.27)	1490	1010
X = >CH-CH ₃ 1.32	R = -C ₆ H ₅ 2.36	C ₂₃ H ₇ ,NS ₂ Sn (86)	0.33 (0.35)	140	22.58 (22.63)	12.18 (12.22)	510 (524.30)	1500	1010
X = > N-CH ₃ 1.42	R = -C _. H, 2.53	C24H26N2S2Sn (89)	0.37 (0.38)	164	22.52 (22.59)	12.16 (12.20)	525 (525.29)	1480	066
X = >0 1.30	R = -C ₆ H, 2.28	C ₁₃ H ₂₃ NOS ₂ Sn (90)	0.33 (0.34)	168	23.12 (23.17)	12.44 (12.51)	512.16 (512.25)	1470	1000

J. SHARMA, Y. SINGH and A. K. RAI

moiety. The central tin atom acquire 5-coordination through weak Sn—S dative bond.^{6,17} The average bond angle of 107° in the present case however indicates further weakening or absence of Sn—S(2) interaction (Figure 1) in solution. But it has to be borne in mind that the structure in solution and solid state may be different, and in organotin derivatives the bond angles in solution are generally less than in solid state as they are free of solid state packing constraints.⁴ Thus, the possibility of weak interaction between Sn and S(2) atoms can not be completely ruled out in the absence of X-ray structural data in solid state.

In contrast to the above, the ¹¹⁹Sn NMR signals of triphenyltin(IV) complexes are in the range δ (-)175-191 ppm. The appearance of a ¹¹⁹Sn signal in this region is consistent with 5-coordination around the central tin atom⁸ (Figure 2).

These results are not surprising in view of the enhanced electronegativity of R group in phenyl derivatives.¹⁸ A five coordination around central tin atom in compounds where $R = -C_6H_5$ is likely to be favoured with trigonal bipyramidal geometry with ligand acquiring axial-equatorial positions in a highly distorted structure.¹⁸

Although we conclude that the dithiocarbamates behave as monodentate ligands when three methyl groups are attached to the central tin atom and as bidentate ligands in presence of three phenyl groups, we recognize that X-ray diffraction studies are essential to confirm bonding modes and geometries in the solid state.

EXPERIMENTAL

The ligands used in this investigation were prepared by literature method.¹⁹ Me₃SnCl was distilled at 153° C and Ph₃SnCl was recrystallised from benzene-hexane solution. Tin and sulfur were estimated gravimetrically as tin oxide and barium sulphate (Messenger's Method), respectively.²⁰ Molecular weights were determined with a "Knauer vapour pressure osmometer," using chloroform solutions at 45° C.

Infrared spectra were recorded in the range 4000–400 cm⁻¹ on a Perkin Elmer 577 spectrophotometer using KBr pellets. ¹H and ¹³C NMR spectra were recorded for CDCl₃ and CHCl₃ solutions respectively on a 90 MHz JEOL FX 90 Q spectrometer using TMS as an internal reference. ¹¹⁹Sn NMR spectra of complexes were recorded in CHCl₃ solution using Me₄Sn as an external reference. ¹³C NMR spectra of four samples in CDCl₃ solution were recorded on a Bruker WH 270 MHz spectrometer using TMS as an internal reference.

Since all complexes were synthesized in the same way, experimental details of a representative complex are given below. Analytical results are summarised in Table III.

Synthesis of $(CH_3)_3Sn(Pipdtc)$; $(C_{19}H_{19}NS_2Sn)$

A benzene solution of Me₃SnCl (2.02 g, 10.2 mmol) was added drop by drop to the benzene suspension of the sodium salt of piperadine dithiocarbamate (2.22 g, 10.2 mmol). The reaction mixture was stirred for ~4 hours at room temperature and then refluxed for ~6 hours to ensure complete reaction. The NaCl formed was filtered off and excess solvent was removed from the filtrate under reduced pressure. The crude product was recrystallized from chloroform-hexane.

ACKNOWLEDGEMENT

One of the authors Jyoti Sharma is thankful to the UGC, New Delhi and the State Government of Rajasthan for providing financial assistance under special assistance programme. Prof. C. L. Khetrapal of Indian Institute of Science Bangalore is thanked for recording 270 MHz ¹³C NMR spectra.

REFERENCES

- 1. H. C. Clark, V. K. Jain, I. J. McMahon and R. C. Mehrotra, J. Organomet Chem., 243, 299 (1983).
- J. Holeček, A. Lyčka, K. Handlier and M. Nadvornik, Collect Czech Chem Commun., 55, 1193 (1990).
- 3. T. N. Srivastava and V. Kumar, J. Organomet. Chem., 107, 55 (1976).
- T. P. Lockhart, W. F. Manders, E. O. Schlemper and J. J. Zuckerman, J. Am. Chem. Soc., 108, 4074 (1986).
- 5. T. P. Lockhart, W. F. Manders and E. O. Schlemper, J. Am. Chem. Soc., 107, 7451 (1985).
- 6. G. M. Sheldrick, W. S. Sheldrick, R. F. Dalton and K. Jones, J. Chem. Soc. (A)., 493 (1970).
- 7. T. P. Lockhart and W. F. Manders, Inorg. Chem., 25, 892 (1986).
- 8. J. Holeček, M. Nadvornik and K. Handlier, J. Organomet. Chem., 315, 299 (1986).
- 9. A. G. Davies, P. G. Harrison, J. D. Kennedy, T. N. Mitchell, R. J. Puddephatt and W. McFarlane, J. Chem. Soc., C, 1136 (1969).
- 10. G. St. Nikoloves, N. Jordanov and I. Hanezov, J. Inorg. Nucl. Chem., 33, 1055 (1971).
- 11. M. Schmidt and H. Schumann, Z. Anorg. Allg. Chem., 325, 130 (1963).
- 12. V. Kawasaki, T. Tanaka and R. Okawara, Bull. Chem. Soc. Japan., 903 (1964).
- 13. M. M. Mc Grady and R. S. Tobias, J. Am. Chem. Soc., 87, 1909 (1965).
- 14. J. Otera, T. Hinoishi, Y. Kawabe and R. Okawara, Chem. Lett., 273 (1981).
- 15. J. Sharma, Y. P. Singh and A. K. Rai, Phosphorous, Sulphur and Silicon., 86, 197 (1994).
- 16. T. N. Mitchell, J. Organomet. Chem., 59, 189 (1973).
- 17. G. M. Sheldrick, W. S. Sheldrick, R. F. Dalton and K. Jones, J. Chem. Soc., (A), 490 (1970).
- 18. K. C. Molloy, S. J. Blunden and R. Hill, J. Chem. Soc. Dalton Trans., 1259 (1988).
- 19. G. Marcotrigiano, G. C. Pellacani and C. Preti, J. Inorg. Nucl. Chem., 36, 3709 (1974).
- 20. A. I. Vogel, "Quantitative Inorg. Analysis," The ELBS Longmans Green and Co. Ltd., 1978.