Research Paper



Applications of hypervalent iodine(III) reagents in constructing ortho-iodo aromatic ethers

Journal of Chemical Research 1–5 © The Author(s) 2021 Article reuse guidelines: sagepub.com/journals-permissions DOI: 10.1177/17475198211005385 journals.sagepub.com/home/chl



Zhai Yang-Yang, Wang Zhen-Hui, Gong Xue-Yun, Hou Bao-Hua, Cui Xiao-Rui, Sun Yan-Feng, Wang Ke-Yang, Liu Peng-Wei, Li Xue-Yan and Zhou De-Jun

Abstract

A one-pot method for the synthesis of aromatic ethers using hypervalent iodine(III) reagents obtained from the corresponding iodoaryl compounds is developed. In this concise method, six diaryl ethers and three heterocyclic aromatic ethers are synthesized in good yields. Furthermore, possible mechanisms for the syntheses of the hypervalent iodine reagents and construction of the aromatic ethers are proposed.

Keywords

diaryl ethers, heterocyclic aromatic ethers, hypervalent iodine(III) compounds, mechanisms

Date received: 3 March 2021; accepted: 8 March 2021



Introduction

Hypervalent iodine(III) reagents have been known for over a century, but these compounds have only recently been applied extensively in many important organic transformations.^{1–3} Diaryl ethers are common structural features in numerous biologically active compounds and natural products, some of which are potential drugs. Numerous methods have been developed to synthesize diaryl ethers, and new routes for a wide range of biologically active compounds have been devised.^{4–8} Heterocyclic aromatic ethers are likely to exhibit higher potent biological activity than the corresponding diphenyl ethers. However, many of the reported methods for the synthesis of heterocyclic aromatic ethers have limited efficiency.⁹ In this study, we report a concise and efficient method for the preparation of trivalent iodine compounds **4–6**, and the application of these compounds for the construction of aromatic ethers.

Results and discussion

In our laboratory, we developed an efficient and inexpensive method for preparing diacetoxyiodoarenes^{10–12} in ideal yields from corresponding iodoarenes with sodium perborate

Medical School, Henan Polytechnic University, Jiaozuo, P.R. China

Corresponding author: Zhou De-Jun, Medical School, Henan Polytechnic University, Jiaozuo 454000, P.R. China. Email: zhoudj@hpu.edu.cn

Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (https://creativecommons.org/licenses/by-nc/4.0/) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (https://us.sagepub.com/en-us/nam/open-access-at-sage).



Scheme I. Preparation of diacetoxyiodoarenes.



Scheme 2. Construction of aromatic ethers.

tetrahydrate.¹³ The optimized method for the preparation of diacetoxyiodoarenes from iodoarenes is described in section "Experimental" (Scheme 1).

The *ortho*-iodo aromatic ethers can be synthesized smoothly from the corresponding phenols with the diacetoxyiodoarenes. In particular, the 4-pyridone also could be oxidized to *ortho*-iodo aromatic ethers via diacetoxyiodoarenes. The detail experiment of the preparation of aromatic ethers is described in section "Experimental" (Scheme 2).

A possible mechanism for the construction of diacetoxyiodoarenes has been proposed based on the experimental



Scheme 3. Possible mechanism for the construction of the diacetoxyiodoarenes.



Scheme 4. A possible mechanism for the formation of heterocyclic aromatic ethers.

results (Scheme 3). The starting material is oxidized to an unstable intermediate iodosyl benzene **a**, which then reacts with acetic anhydride to form intermediate **b**. Intramolecular rearrangement then yields the diacetoxyiodoarenes. Infrared spectroscopy and UV-Vis light spectroscopy data showed that all of the diacetoxyiodoarenes existed in virtually ionic form.¹³

Based on the experiment results and previous reports,¹⁴ a possible mechanism for the formation of heterocyclic aromatic ethers is shown in Scheme 4. Initially, the lone pair of the hydroxy group attacks the iodine of the diacetoxyiodoarene, which loses an acetic acid to form intermediate **c**. Intramolecular electrophilic aromatic substitution with loss of an acetate anion then yields cation **d**. Next, aromatization forms the relatively stable intermediate **e**, and a subsequent shift of the phenyl group from iodine to oxygen then yields the heterocyclic aromatic ether.

Conclusion

In this work, we report an efficient method for synthesis of diacetoxyiodoarenes via intramolecular rearrangement, and three diacetoxyiodoarenes were prepared smoothly via this method. Then, the front freshly prepared compounds are used to oxidize some phenols, and the corresponding six diaryl ethers and three heterocyclic aromatic ethers are synthesized in good yields. Both of the possible mechanisms of preparing diacetoxyiodoarenes and constructing diaryl ethers are proposed.

Experimental

All chemical reagents were obtained from commercial suppliers (Aldrich, Tokyo Chemical Industry (TCI), Aladdin, Macklin, and Bidepharm Pure Chemical Industries) and

were used without further purification. Anhydrous solvents were obtained via standard protocols. All non-aqueous reactions were carried out under an Ar atmosphere. Thinlayer chromatography (TLC) was performed on silica gel 60 F₂₅₄ glass plates precoated with a 0.25-mm thickness of silica gel (Yantai Jiangyou). Column chromatography was carried out on Cica Silica Gel 60N (spherical, neutral, 40-50 µm or 63-210 µm). ¹H and ¹³C NMR spectra were obtained on a Varian UNITY plus 300 (300 MHz for ¹H and 75 MHz for ¹³C) instrument, with CDCl3 as the solvent an d internal reference. IR spectra were measured on a JNM FTIR-460 Plus spectrometer. Mass spectra were recorded on JEOL D-200, JEOL JMS-GCmate II, SHIMAZU GC-MS-QP 500, or JEOL AX 505 spectrometers. Melting points were recorded with a Yanagimoto micro melting point apparatus and are uncorrected.

Optimized method for the preparation of diacetoxyiodoarenes from iodoarenes

 $NaBO_3 \cdot 4H_2O$ (10 mmol) was slowly added portionwise over 30 min to a stirred solution of iodoarene 1 (1 mmol) in Ac_2O (1.5 mL) and AcOH (99.5%, 1.5 mL) under Ar at 55 °C. The mixture was stirred at 55 °C until the starting material had disappeared completely (as confirmed by TLC analysis). Water (20 mL) was added and the reaction mixture was stirred at room temperature for 30 min. A considerable amount of solid precipitates was observed. The solid was separated by filtration, washed with water (25 mL), and dried under air at room atmosphere. The filtrate was extracted with EtOAc (3 10 mL). And the combined extracts were dried over anhydrous MgSO₄ and filtered. The solvent was evaporated under reduced pressure. The combined crude products were purified by recrystallization from a mixture of AcOH/H₂O (1:1).

General procedure for the preparation of aromatic ethers

A mixture of 4-pyridone or substrate **10–12** (100 mg) and freshly prepared diacetoxyiodoarene (5 equiv.) in MeOH (5mL) was refluxed for 4–8h. The reaction mixture was quenched with saturated NaCl (10mL) and extracted with ethyl acetate (3×10 mL). The organic layers were combined, dried over MgSO₄, and concentrated under vacuum. The residue was purified by column chromatography on silica gel to afford the desired product.

Diacetoxyiodobenzene (4). Yield: 91%; white solid; m.p. $161-163 \,^{\circ}$ C (lit.¹⁵161.2–162.2 °C). ¹H NMR (300 MHz, CDCl₃): δ 1.98 (6 H, s), 7.46 (2 H, d, *J*=7.6 Hz), 7.58 (1 H, td, *J*=7.6 Hz, *J*=1.1 Hz), 8.08 (2 H, dd, *J*=7.6 Hz, *J*=1.1 Hz); ¹³C NMR (75 MHz, CDCl₃): δ 20.18 (q), 20.18 (q), 121.38 (s), 130.77 (d), 131.59 (d), 134.76 (d), 176.19 (s).

p-(Diacetoxyiodo)toluene (5). Yield: 89%; white solid; m.p. 109–110 °C. ¹H NMR (300 MHz, CDCl₃): δ 1.97 (6 H, s), 2.40 (s, 3 H), 7.27 (2 H, d, *J*=8.1 Hz), 7.95 (2 H, d, *J*=8.1 Hz); ¹³C NMR (75 MHz, CDCl₃): δ 20.7 (q), 21.6 (q), 119.2 (s), 131.8 (d), 134.5 (d), 142.8 (s), 176.3 (s). IR (KBr, cm⁻¹): 3385, 3321, 2261, 1631, 1441, 1385, 1193; MS-ES: m/z=336 (M⁺); HRMS-ES: m/z [M]⁺ calcd for C₁₁H₁₃IO₄: 335.9859; found: 335.9849.

o-(Diacetoxyiodo)toluene (**6**). Yield: 92%; white solid; m.p. 100–102 °C. ¹H NMR (300 MHz, CDCl₃): δ 1.98 (6 H, s), 2.72 (3 H, s), 7.22-7.28 (1 H, m), 7.50-7.52 (2 H, m), 8.16 (1 H, d, *J*=7.42 Hz); ¹³C NMR (75 MHz, CDCl₃): δ 20.13 (q), 25.39 (q), 126.83 (s), 128.05 (d), 130.49 (d), 132.37 (d), 136.82 (d), 140.20 (s), 175.94 (s). IR (neat, cm⁻¹) 3435, 1649, 1614, 1367, 1293, 1011, 764, 669; MS-ES: *m/z*=336 (M⁺); HRMS-ES: *m/z* [M]⁺ calcd for C₁₁H₁₃IO₄: 335.9859; found: 335.9851.

lodo-4-phenoxypyridine (7). Yield: 94%; colorless oil. ¹H NMR (300 MHz, CDCl₃): δ 6.56 (1H, d, J=5.77 Hz), 7.09-7.12 (2 H, m), 7.25-7.31 (1 H, m), 7.41-7.48 (2 H, m), 8.29 (1 H, d, J=5.77 Hz), 8.86 (1 H, s); ¹³C NMR (75 MHz, CDCl₃): δ 85.38 (s), 111.01 (d), 120.65 (d), 125.73 (d), 130.18 (d), 150.36 (d), 153.66 (s), 158.43 (d), 163.74 (s); IR (neat, cm⁻¹): 3038, 2360, 1562, 1464, 1397, 1270, 1199, 882; MS-ES: m/z=297 (M⁺); HRMS-ES: m/z [M]⁺ calcd for C₁₁H₈INO: 296.9651; found: 296.9652.

3-lodo-4-(*p*-tolyloxy)pyridine (**8**). Yield: 91%; colorless oil. ¹H NMR (300 MHz, CDCl₃): δ 2.37 (3 H, s), 6.53 (1 H, d, *J*=5.77 Hz), 6.97 (2 H, d, *J*=9.52 Hz), 7.20 (2 H, d, *J*=9.52 Hz), 8.25 (1 H, d, *J*=5.77 Hz), 8.83 (1 H, s); ¹³C NMR (75 MHz, CDCl₃): δ 20.92 (q), 85.15 (s), 110.69 (d), 120.46 (d), 130.44 (d), 135.44 (s), 150.25 (d), 151.28 (s), 158.25 (d), 163.93 (s); IR (neat, cm⁻¹): 3034, 1567, 1504, 1462, 1271, 1201, 1016, 885; MS-ES: *m*/*z*=311 (M⁺); HRMS-ES: *m*/*z* [M]⁺ calcd for C₁₂H₁₀INO: 310.9807; found: 310.9799.

3-lodo-4-(*o*-tolyloxy)pyridine (**9**). Yield: 95%; colorless oil. ¹H NMR (300 MHz, CDCl₃): δ 2.16 (3 H, s), 6.41 (1 H, d, *J*=5.49 Hz), 7.03 (1 H, d, *J*=7.69 Hz), 7.19-7.32 (4 H, m), 8.28 (1 H, d, *J*=5.77 Hz), 8.86 (1 H, s); ¹³C NMR (75 MHz, CDCl₃): δ 16.07 (q), 84.64 (s), 109.89 (d), 121.14 (d), 126.19 (d), 127.55 (d), 130.26 (s), 131.87 (d), 150.12 (d), 151.54 (s), 157.96 (d), 163.42 (s); IR (neat, cm⁻¹): 3035, 2360, 1563, 1464, 1271, 1181, 888; MS-ES: *m/z*=311 (M⁺); HRMS-ES: *m/z* [M]⁺ calcd for C₁₂H₁₀INO: 310.9807; found: 310.9791.

Methyl 3-iodo-4-(*p*-tolyloxy)benzoate (**13**). Yield: 85%; colorless oil. ¹H NMR (300 MHz, CDCl₃): δ 2.36 (3 H, s), 3.90 (3 H, s), 6.72 (1 H, d, *J*=8.52 Hz), 6.95 (2 H, d, *J*=7.97 Hz), 7.19 (2 H, d, *J*=7.97 Hz), 7.89 (1 H, dd, *J*=8.52, 1.92 Hz), 8.52 (1 H, d, *J*=1.92 Hz); ¹³C NMR (75 MHz, CDCl₃): δ 20.91 (q), 52.26 (q), 86.48 (s), 115.82 (d), 119.80 (d), 125.85 (s), 130.47 (d), 131.06 (d), 134.49 (s), 141.25 (d), 152.91 (s), 161.12 (s), 165.13 (s); IR (neat, cm⁻¹) 2863, 1720, 1588, 1505, 1477, 1433, 1253; MS-ES: *m*/*z*=368 (M⁺); HRMS-ES: *m*/*z* [M]⁺ calcd for C₁₅H₁₃IO₃: 367.9909; found: 367.9909.

Methyl 5-iodo-2-methoxy-4-(p-tolyloxy)benzoate (14). Yield: 91%; white solid; m.p. 77–79 °C. ¹H NMR (300 MHz, CDCl₃): δ 2.36 (3 H, s), 3.69 (3 H, s), 3.87 (3 H, s), 6.35 (1 H, s), 6.94 (2 H, d, *J*=8.52 Hz), 7.19 (2 H, d, *J*=8.52 Hz), 8.30 (1 H, s); ¹³C NMR (75 MHz, CDCl₃): δ 20.91 (q), 52.08 (q), 56.18 (q), 75.09 (s), 101.54 (d), 116.12 (s), 119.36 (d), 130.46 (d), 134.28 (s), 142.43 (d), 152.98 (s), 161.10 (s), 161.31 (s), 164.50 (s); IR (neat, cm⁻¹): 2948, 1687, 1588, 1505, 1438, 1276, 1103, 834; MS-ES: *m*/*z*=398 (M⁺); HRMS-ES: *m*/*z* [M]⁺ calcd for C₁₆H₁₅IO₄: 398.0015; found: 398.0010.

2-iodo-4-nitro-1-(p-tolyloxy)benzene (**15**). Yield: 79%; colorless oil. ¹H NMR (300 MHz, CDCl₃): δ 2.39 (3 H, s), 6.71 (1 H, d, *J*=9.07 Hz), 6.98 (2 H, d, *J*=8.52 Hz), 7.24 (2 H, d, *J*=8.52 Hz), 8.10 (1 H, dd, *J*=9.07, 2.75 Hz), 8.73 (1 H, d, *J*=2.75 Hz); ¹³C NMR (75 MHz, CDCl₃): δ 20.97 (q), 85.66 (s), 114.63 (d), 120.22 (d), 125.13 (d), 130.77 (d), 135.33 (d), 135.51 (s), 142.66 (s), 152.17 (s), 162.78 (s); IR (neat, cm⁻¹): 3093, 2923, 1576, 1518, 1465, 1342, 1260893; MS-ES: *m/z*=355 (M⁺); HRMS-ES: *m/z* [M]⁺ calcd for C₁₃H₁₀INO₃: 354.9706; found: 354.9713.

Methyl 3-iodo-4-(o-tolyloxy)benzoate (16). Yield: 89%; colorless oil. ¹H NMR (300 MHz, CDCl₃): δ 2.19 (3 H, s), 3.90 (3 H, s), 6.54 (1H, d, *J*=8.79 Hz), 6.97 (1 H, d, *J*=7.69 Hz), 7.14-7.31 (3 H, m), 7.87 (1 H, dd, *J*=8.79, 1.92 Hz), 8.54 (1 H, d, *J*=1.92 Hz); ¹³C NMR (75 MHz, CDCl₃): δ 16.25 (q), 52.25 (q), 85.67 (s), 114.38 (d), 120.56 (d), 125.38 (d), 125.54 (s), 127.38 (d), 130.10 (s), 131.15 (d), 131.74 (d), 141.30 (d), 152.90 (s), 160.57 (s), 165.15 (s); IR (neat, cm⁻¹): 2950, 2360, 1720, 1581, 1478, 1433, 1283, 1252, 1111; MS-ES: *m/z*=368 (M⁺); HRMS-ES: *m/z* [M]⁺ calcd for C₁₅H₁₃IO₃: 367.9909; found: 367.9910.

Methyl 5-iodo-2-methoxy-4-(o-tolyloxy)benzoate (**17**). Yield: 86%; colorless oil. ¹H NMR (300 MHz, CDCl₃): δ 2.21 (3 H, s), 3.65 (3 H, s), 3.87 (3 H, s), 6.18 (1 H, s), 6.94 (1 H, d, *J*=7.97 Hz), 7.12-7.31 (3 H, m), 8.32 (1 H, s); ¹³C NMR (75 MHz, CDCl₃): δ 16.28 (q), 52.05 (q), 56.11 (q), 74.27 (s), 100.07 (d), 115.65 (s), 119.99 (d), 125.23 (d), 127.32 (d), 129.86 (s), 131.72 (d), 142.50 (d), 152.87 (s), 160.91 (s), 161.25 (s), 164.48 (s); IR (neat, cm⁻¹): 2949, 2360, 1731, 1592, 1488, 1244, 1093, 781; MS-ES: *m/z*=398 (M⁺); HRMS-ES: *m/z* [M]⁺ calcd for C₁₆H₁₅IO₄: 398.0015; found: 398.0010.

2-iodo-4-nitro-1-(o-tolyloxy)benzene (**18**). Yield: 85%; colorless oil. ¹H NMR (300 MHz, CDCl₃): δ 2.17 (3 H, s), 6.54 (1 H, d, *J*=9.07 Hz), 7.01 (1 H, d, *J*=7.69 Hz), 7.19-7.34 (3 H, m), 8.09 (1 H, dd, *J*=9.07, 2.75 Hz), 8.75 (1 H, d, *J*=2.75 Hz); ¹³C NMR (75 MHz, CDCl₃): δ 16.13 (q), 85.03 (s), 113.45 (d), 120.90 (d), 125.25 (d), 126.15 (d),

127.67 (d), 130.13 (s), 131.98 (d), 135.39 (d), 142.56 (s), 152.32 (s), 162.06 (s); IR (neat, cm⁻¹): 3093, 2359, 1574, 1518, 1462, 1341, 1258, 745; MS-ES: m/z=355 (M⁺); HRMS-ES: m/z [M]⁺ calcd for C₁₃H₁₀INO₃: 354.9705; found: 354.9713.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: The authors are grateful to the National Natural Science Foundation of China (no. 81803399), the Science and Technology Research Project of Henan Province (nos 182102310300 and 192102310144), and the Colleges and Universities in Henan Province Key Scientific Research Project (nos 18A150028 and 19B350003) for the financial support.

ORCID iD

Zhou De-Jun (D) https://orcid.org/0000-0002-9577-306X

References

- Qin Y, Zhu LH and Luo SZ. Chem Rev 2017; 117: 9433– 9520.
- 2. Nadia O, Ilchenko MH and Kalman JS. *Chem Sci* 2017; 8: 1056–1061.
- 3. Erwann G and Jerome W. Org Lett 2018; 20: 1473-1476.
- Kise N, Hamada Y and Sakurai T. J Org Chem 2016; 81: 11043–11056.
- Saper NI and Hartwig JF. J Am Chem Soc 2017; 139: 17667– 17676.
- Evans DA, Wood MR, Trotter BW, et al. Angew Chem Int Ed 1998; 37: 2700–2704.
- Nicolaou KC, Natarajan S, Li H, et al. *Angew Chem Int Ed* 1998; 37: 2708–2714.
- Boger DL, Miyazaki S, Kim SH, et al. J Am Chem Soc 1999; 121: 10004–10011.
- Wang D, Zhang HJ, Fan Y, et al. *Sci. Total Environ.* 2020; 721: 137563–137569.
- 10. Kazmierczak P and Skulski L. Synthesis 1998; 24: 1721-1723.
- 11. Alcock NW and Waddington TD. *J Am Chem Soc* 1963; 85: 4103–4109.
- 12. Hossain MD and Kitamura T. J Org Chem 2005; 70: 6984–6986.
- Zhou DJ, Zhai YY, Meng LP, et al. *Lett Organ Chem* 2019; 16: 911–914.
- 14. Zhou DJ, Yin SQ, Fan YC, et al. *Resea Chem Int* 2016; 42: 5387–5394.
- 15. Leffler JE and Story LJ. J Am Chem Soc 1967; 89: 2333– 2338.