Acknowledgment. This work was supported by a grant from the National Institutes of Health, for which we are grateful. We thank Dr. G. Wallraff of IBM Almaden Research Center for his help with the molecular mechanics calculations.

## Synthesis of Simple CrL<sub>4</sub>(alkyne) Complexes by Displacement of a Labile Cyclooctadiene Ligand

Donald J. Wink\* and B. Timothy Creagan

Department of Chemistry, New York University New York, New York 10003 Received June 27, 1990

We have shown that, in contrast to the simple tetracarbonyl system  $M(CO)_4$ (diene)<sup>1,2</sup> (M = Cr, Mo, W), 1,5-cyclooctadiene (1,5-COD) is displaced from  $M(CO)_3(P(OMe)_3)(\eta^4-1,5-COD)$ by conjugated dienes.<sup>3</sup> As reported here, alkynes, which are similar to 1.3-dienes in their frontier orbital characteristics,<sup>4</sup> also displace the COD in  $Cr(CO)_3(P(OMe)_3)(\eta^4-1,5-COD)$  to afford unprecedented  $CrL_4(PhC \equiv CR)$  complexes.

Diphenylacetylene and phenylacetylene react in a few hours with mer-Cr(CO)<sub>3</sub>(P(OMe)<sub>3</sub>)( $\eta^4$ -1,5-COD) to give products of COD substitution and phosphite and carbonyl ligand redistribution (eq 1).<sup>7,8</sup> The PhC=CPh complex 1a<sup>9</sup> forms in good yield as black crystals directly from the reaction mixture in pentane while the PhC=CH complex 1b is crystallized separately in lower yield. A superior route to 1b is alkyne exchange with 1a (eq 2).<sup>10</sup>

$$mer-Cr(CO)_{3}P(OMe)_{3}(\eta^{4}-1,5-COD) + RC \equiv CPh \xrightarrow{C_{3}(\eta_{2})} Cr(CO)_{2}(P(OMe)_{3})_{2}(\eta^{2}-RC \equiv CPh) + COD + 1a, R = Ph; 1b, R = H Cr(CO)_{4}(1,5-COD) (1)$$

 $Cr(CO)_2(P(OMe)_3)_2(\eta^2 PhC = CPh) + HC = CPh - C_3H_{12}$  $Cr(CO)_{2}(P(OMe)_{1})_{2}(\eta^{2}-HC\equiv CPh) + PhC\equiv CPh$  (2)

Integration of the <sup>1</sup>H NMR spectra supports a 2:1 phosphite:alkyne stoichiometry, and a pair of bands in the IR spectra for  $\nu(CO)$  indicates a *cis*-dicarbonyl structure. The signals for the alkyne carbons for 1a appear at  $\delta$  180.2 and for 1b at  $\delta$  174.3 (=CPh) and  $\delta$  163.7 (=CH). These are downfield from the region generally cited for the carbons of a four-electron alkyne,<sup>11</sup> but of course the metal in this case is quite electron rich. Resonances for the alkyne and carbonyl carbons for both complexes and for the alkyne hydrogen in 1b appear as triplets due to coupling to two phosphorus ligands. This is consistent with either structure 2, with magnetically equivalent phosphorus atoms, or 3, in which the phosphorus atoms would be rendered equivalent only by rotation of the alkyne. Variable-temperature NMR experiments with 1b establish that the ground-state structure is

(1) Kreiter, C. G. Adv. Organomet. Chem. 1986, 26, 297

- (2) Dixon, D. T.; Burkinshaw, P. M.; Howell, J. A. S. J. Chem. Soc., Dalton Trans. 1980, 2237.
- (3) (a) Wink, D. J.; Oh, C. K. Organometallics 1990, 9, 2605. (b) Wink, D. J.; Wang, N.-F.; Creagan, B. T. Organometallics 1989, 8, 561.

(4) The isolobal character of alkynes and other four-electron donors has (4) The isobal end of the interview of a large state of the interview of the i

Vol. 3

(6) Wink, D. J. Organometallics, in press.
(7) Michael, G.; Kreiter, C. G. Z. Naturforsch. 1984, 39B, 1738.
(8) No intermediates are detected by <sup>1</sup>H NMR spectroscopy when the phenylacetylene reaction is performed in CD<sub>2</sub>Cl<sub>2</sub>.
(9) Complex 1a is prepared in 58% yield (29% based on Cr). <sup>13</sup>C NMR:

d 180.2 (t,  $J_{PC} = 15$  Hz,  $C_{alkyne}$ ). IR:  $\nu(CO) = 1932$ , 1850 cm<sup>-1</sup>. Complete details are provided in the supplementary material. (10) Complex **1b** is prepared from **1a** in 57% yield. <sup>13</sup>C NMR: 175.5 (t,  $^{2}J_{PC} = 9$  Hz,  $\equiv CPh$ ), 163.7 (t,  $^{2}J_{PC} = 18$  Hz,  $\equiv CH$ ). IR:  $\nu(CO) = 1934$ , 1840 cm<sup>-1</sup>

1840 cm<sup>-1</sup>

(11) Templeton, J. L.; Ward, B. C. J. Am. Chem. Soc. 1980, 102, 3288.

P1 СЗ C15 C16 С9 C10 C26 C25 C 68 06 08 C2 C6 Figure 1. ORTEP diagram of  $Cr(CO)_2(P(OMe)_3)_2(\eta^2-PhC=CPh)$ . Im-

portant bond lengths (Å) and angles (deg): Cr1-P1, 2.252 (2); Cr1-P2, 2.264 (2); Cr1-C1, 1.857 (7); Cr1-C2, 1.827 (7); Cr1-C9, 1.959 (6); Cr1-C10, 1.958 (5); C1-O1, 1.144 (6); C2-O2, 1.166 (7); C9-C10, 1.311 (7); P1-Cr1-P2, 153.94 (7); C1-Cr1-C2, 107.0 (3); C10-C9-C11, 135.0 (5); C9-C10-C21, 136.0 (5).

3; the triplet with  $J_{PH} = 20$  Hz for the acetylenic proton "loses" the central resonance and becomes a doublet with  $J_{PH} = 40 \text{ Hz}$ upon cooling.<sup>12</sup> A barrier to formal rotation of the alkyne of 13.3 (3) kcal mol<sup>-1</sup> is calculated from the decoalescence of the  $^{13}C$ NMR resonance for the phosphite methyl carbons at ca. -5 °C  $(\Delta = 51 \text{ Hz}).^{13}$ 



An X-ray diffraction study<sup>14</sup> of **1a** confirms that in the solid state the alkyne ligand lies almost perpendicular<sup>15</sup> with respect to the P-Cr-P plane in a distorted trigonal-bipyramidal geometry  $(P-Cr-P, 153.94 (7)^{\circ}; OC-Cr-CO, 107.0 (3)^{\circ})$  (Figure 1). Distances and angles involving the alkyne are consistent with those in other chromium-diphenylacetylene complexes.<sup>16</sup>

The alkyne ligands in these complexes are labile to displacement by a variety of donor ligands, including carbon monoxide and trimethyl phosphite, to apparently give other CrL<sub>6</sub> complexes. Decomposition occurs, with liberation of free alkyne, in the presence of thf and ethyl diazoacetate. There is no reaction with conjugated or nonconjugated dienes.

Molecular orbital calculations at the extended Hückel level performed on a  $Cr(CO)_4(\eta^2-HC==CH)$  model show that the conformation with the alkyne perpendicular to the P-Cr-P plane is preferred by 15 kcal mol<sup>-1</sup>. This orientation maximizes the

<sup>(12)</sup> Honeychuck, R. V.; Hersh, W. J. Am. Chem. Soc. 1989, 111, 6056.

<sup>(12)</sup> Honeychuck, R. V.; Hersh, W. J. Am. Chem. Soc. 1989, 111, 6056. (13) Binsch, G. In Dynamic Nuclear Magnetic Resonance Spectroscopy; Jackman, L. M., Cotton, F. A., Eds.; Academic: New York, 1975; pp 45-82. (14) Crystals were grown from ether at -20 °C. The space group is  $P_{ben}$ with a = 19.064 (3) Å, b = 14.070 (2) Å, c = 18.979 (2) Å, and V = 5091(2) Å<sup>3</sup>. Refinement of 298 least-squares parameters using 2205 reflections with  $I > 2.0\sigma(I)$  gave R = 4.6% and  $R_w = 5.6\%$ . (15) There is a slight tilt (12°) from the perpendicular. (16) (a) Koldova, N. E.; Zhvanko, O. S.; Andreanov, V. G.; Karapyetyan, A. A.; Struchkov, Y. T. Koord. Khim. 1980, 6, 1407. (b) Wink, D. J.; Fox, J. R.; Cooper, N. J. J. Am. Chem. Soc. 1985, 107, 5012. (c) Salt, J. E.; Girolami, G. S.; Wilkinson, G.; Motevalli, M.; Thornton-Pett, M.; Hursthouse, M. B. J. Chem. Soc., Dalton Trans. 1985, 685. M. B. J. Chem. Soc., Dalton Trans. 1985, 685.

 $\pi$ -donation with a vacant metal fragment orbital in the equatorial plane and  $\pi$ -back-bonding with an orbital oriented in the axial direction.<sup>17</sup> When the system is converted to a simple square pyramid, the calculated barrier to formal rotation becomes 12 kcal mol<sup>-1</sup>, reminiscent of change in the barrier for olefin rotation in d<sup>8</sup> Fe(CO)<sub>4</sub>(olefin).<sup>18</sup>

We are aware of no reports concerning the synthesis of a simple four-electron monoalkyne complex of a  $d^6$  ML<sub>4</sub> fragment.<sup>19</sup> This is remarkable considering the extensive group 6 alkyne chemistry that began with the synthesis of W(alkyne)<sub>3</sub>(CO) complexes over 25 years ago.<sup>21</sup> The availability of reagents that are both electron rich and labile, such as the 1,5-COD complexes utilized here, is critical to the synthetic strategy for introducing excellent  $\pi$ -acceptor ligands, such as alkynes and 1,3-dienes.

Acknowledgment. We are grateful to Drs. James Springer and John Dewan for helpful discussions about the crystal structure determination. We are grateful to the donors of the Petroleum Research Fund, administered by the American Chemical Society, for financial support and to the National Science Foundation for funds to purchase an X-ray diffractometer.

Supplementary Material Available: Synthetic and spectroscopic data for 1a and 1b, details of extended Hückel calculations, and methods and results (listings) of the crystal structure study of 1a (11 pages); listing of observed and calculated structure factor amplitudes (16 pages). Ordering information is given on any current masthead page.

(19) Four-electron mono(alkyne) complexes of formally d<sup>2</sup> or d<sup>4</sup> group 6 metal centers are well-known,<sup>20</sup> and some d<sup>6</sup> bis(alkyne) containing four-electron alkynes are also known: refs 16b,c. Dötz, K. H.; Mühlmeier, J. Angew. Chem., Int. Ed. Engl. 1982, 21, 929.
(20) Templeton, J. L. Adv. Organomet. Chem. 1989, 29, 1.

21) Tate, D. P.; Augl, J. M.; Ritchey, W. M.; Ross, B. L.; Grasselli, J. G. J. Am. Chem. Soc. 1964, 86, 3261.

## An Unusual, Electronically Unsaturated Dirhenium Polyhydride Complex with No Metal-Metal Bond

Karen E. Meyer, Phillip E. Fanwick, and Richard A. Walton\*

> Department of Chemistry, Purdue University West Lafayette, Indiana 47907 Received July 20, 1990

Many mononuclear and dinuclear mixed polyhydride-phosphine (or arsine or stibine) complexes of rhenium are known, in a variety far greater than that for any other transition element. In the case of the dirhenium species, these are of the generic types  $[Re_2H_9(PR_3)_4]^+$ ,  $Re_2H_8(PR_3)_4$ ,  $[Re_2H_7(PR_3)_5]^+$ ,  $Re_2H_6(PR_3)_5$ ,  $[Re_2H_5(PR_3)_6]^+$ , and  $Re_2H_4(PR_3)_6$ , where PR<sub>3</sub> represents a monodentate or one-half a bidentate phosphine.<sup>1-14</sup> With one

- Chatt, J.; Coffey, R. S. J. Chem. Soc. A 1969, 1963.
   Bau, R.; Carroll, W. E.; Teller, R. G.; Koetzle, T. F. J. Am. Chem. Soc. 1977, 99, 3872.
- (1) Allison, J. D.; Walton, R. A. J. Am. Chem. Soc. 1984, 106, 163.
   (4) Moehring, G. A.; Fanwick, P. E.; Walton, R. A. Inorg. Chem. 1987, 26 1861
- (5) Cotton, F. A.; Luck, R. L. Inorg. Chem. 1989, 28, 4522.
  (6) Cotton, F. A.; Luck, R. L.; Root, D. R.; Walton, R. A. Inorg. Chem.
  1990, 29, 43.
  (7) Costello, M. T.; Moehring, G. A.; Walton, R. A. Inorg. Chem. 1990, 29, 1578.
- (8) Green, M. A.; Huffman, J. C.; Caulton, K. G. J. Am. Chem. Soc. 1981, 103, 695.



Figure 1. ORTEP view of the structure of the [Re<sub>2</sub>H<sub>5</sub>(dmpm)<sub>3</sub>]<sup>+</sup> cation with the hydrogen atoms of the dmpm ligands omitted. The thermal ellipsoids are drawn at the 50% probability level, except for the hydride ligands, which are circles of arbitrary radius. Some representative and important bond distances (Å) and angles (deg) and torsional angles (deg) are as follows: Re(1)-P(11) = 2.408(2), Re(1)-P(12) = 2.365(2),Re(1)-P(13) = 2.371(2), Re(1)-H(B) = 1.53(7), Re(1)-H(11) = 1.64(8),  $\operatorname{Re}(1) - H(12) = 1.7(1)$ ,  $\operatorname{Re}(2) - H(B) = 2.06(7)$ ;  $\operatorname{Re}(2) - \operatorname{Re}(1) - P$ -(11) = 85.48 (4), Re(2)-Re(1)-H(11) = 76 (3), Re(2)-Re(1)-H(12)= 135 (3), P(11)-Re(1)-H(12) = 139 (3), P(11)-Re(1)-P(12) = 102.73(7), P(12)-Re(1)-P(13) = 151.07 (6); Re(1)-H(B)-Re(2) = 156 (5); P(11)-Re(1)-Re(2)-P(21) = 0.30 (7), P(12)-Re(1)-Re(2)-P(22) =2.34 (7), P(13)-Re(1)-Re(2)-P(23) = 1.69 (7). Distances involving Re(2) are similar to those for Re(1) and, therefore, are not listed here.

exception, the structural characterizations of representative examples of these compounds have revealed the presence of three or four bridging hydrido ligands and a short Re-Re distance (2.51-2.61 Å).<sup>2,5,8-11</sup> The latter observation has led to the conclusion that some degree of multiple metal-metal bonding is present, represented in terms of direct Re-Re bonding and/or three-center, two-electron Re-H-Re bonds.<sup>15,16</sup> The single exception is the complex  $Re_2(\mu-H)_2H_6(\mu-dppm)_2$  (dppm = Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub>), which contains only two bridging hydrido ligands and possesses the much longer Re-Re bond distance of 2.9335 (9) Å. We now report the isolation of a molecule, viz.,  $[Re_2H_5(dmpm)_3]PF_6(dmpm = Me_2PCH_2PMe_2)$ , that takes this trend to its extreme, namely, no direct Re-Re interaction whatsoever and metal centers that are electronically unsaturated. This may presage the discovery of other compounds of this type with metals that normally show a propensity to form strong metal-metal interactions.1

The reaction of the triply bonded dirhenium(II) complex  $\operatorname{Re}_{2}\operatorname{Cl}_{4}(\mu\operatorname{-dmpm})_{3}^{18}$  (0.453 g, 0.49 mmol) with LiAlH<sub>4</sub> (0.32 g, 8.4 mmol) in THF (20 mL) at room temperature afforded a green solution. The mixture was stirred for 2 h, hydrolyzed with 20 mL of a  $H_2O/THF$  (5:15 mL) mixture, and then heated gently. After

- (9) Green, M. A.; Huffman, J. C.; Caulton, K. G. J. Am. Chem. Soc. 1982, 104, 2319.
- (10) Allison, J. D.; Cotton, F. A.; Powell, G. L.; Walton, R. A. Inorg. Chem. 1984, 23, 159
- (11) Fanwick, P. E.; Root, D. R.; Walton, R. A. Inorg. Chem. 1989, 28, 2239.
- (12) Root, D. R.; Meyer, K. E.; Walton, R. A. Inorg. Chem. 1989, 28, 2503.
- (13) Costello, M. T.; Fanwick, P. E.; Meyer, K. E.; Walton, R. A. Inorg. Chem., in press.
- (14) An interesting example of a complex that contains a tridentate phosphine ligand has been encountered in the case of the unsymmetrical anionic species  $[H_6Re(\mu-H)_3Re[MeC(CH_2PPh_2)_3]^-$ : Abrahams, S. C.; Ginsberg, A. P.; Koetzle, T. F.; Marsh, P.; Sprinkle, C. R. Inorg. Chem. 1986, 25, 250Ō.
- (15) Dedieu, A.; Albright, T. A.; Hoffmann, R. J. Am. Chem. Soc. 1979, 101, 3141.
- (16) Dahl, L. F. Ann. N.Y. Acad. Sci. 1983, 415, 1.
- (17) Cotton, F. A.; Walton, R. A. Multiple Bonds Between Metal Atoms; Wiley: New York, 1982. (18) Anderson, L. B.; Cotton, F. A.; Falvello, L. R.; Harwood, W. S.;
- Lewis, D.; Walton, R. A. Inorg. Chem. 1986, 25, 3637.

<sup>(17) (</sup>a) Wink, D. J.; Creagan, B. T. Organometallics 1990, 9, 328. (b) Kamata, M.; Hirotsu, K.; Higuchi, T.; Kido, M.; Tatsumi, K.; Yoshida, T.; Otsuka, S. Inorg. Chem. 1983, 22, 2416. (c) Tatsumi, K.; Hoffmann, R.; Templeton, J. L. Inorg. Chem. 1982, 21, 466. (d) Templeton, J. L.; Winston, P. B.; Ward, B. C. J. Am. Chem. Soc. 1981, 103, 7713. (18) Albright, T. A.; Hoffmann, R.; Thibeault, J. C.; Thorn, D. L. J. Am. Chem. Soc. 1979, 101, 3801.
(19) Four-electron mono(alkune) complexes of formally d<sup>2</sup> or d<sup>4</sup> aroup 6