## Enantioselective Total Synthesis of (+)-Isolaurepinnacin

Daniel Berger,<sup>1</sup> Larry E. Overman,<sup>\*</sup> and Paul A. Renhowe

Department of Chemistry University of California Irvine, California 92717-2025

Received June 23, 1993

Red seaweeds of the genus Laurencia produce a wide variety of halogenated C15 acetogenins, which characteristically contain terminal envne or bromoallene functionality.<sup>2</sup> The majority of these metabolites are oxygen heterocycles, with five- and eightmembered oxacycles occurring particularly widely.<sup>2</sup> Isolaurepinnacin  $(1)^3$  and rogioloxepane A  $(2)^4$  are examples of the rarer class of C15 Laurencia metabolites that contain seven-membered oxacyclic (oxepane) rings.<sup>5</sup> In this communication, we report



the first total synthesis of a Laurencia acetogenin of the oxepane group.<sup>6</sup> Our strategy was to form the *cis*-2,7-disubstituted  $\Delta^4$ oxepene ring of 1 by an acetal-alkene cyclization.<sup>7</sup> The defining reaction of the total synthesis of (+)-isolaurepinnacin recorded herein is the selective conversion of a  $\beta$ -chloro mixed acetal (3) to an oxepene (4) (eq 1), a conversion that effectively deals with all of the structural and stereochemical issues posed by this total synthesis target.6d



The syntheses of the (3S,4S)-vinylsilane alcohol 9 and the (R)- $\alpha$ -chloroacetal 13 precursors of the mixed acetal cyclization substrate 15 are summarized in Scheme I. Phenylsulfonylation of enantiopure epoxy alcohol 5,8-10 followed by regioselective

(1) Current address: American Cyanamid Co., Lederle Laboratories Division, Middletown Rd., Pearl River, NY 10965.

(2) Moore, R. E. In Marine Natural Products; Scheuer, P. J., Ed.; Academic: New York, 1978; Vol. 1, pp 43-121. Erickson, K. L. In Marine Natural Products; Scheuer, P. J., Ed.; Academic: New York, 1983; Vol. 5, pp 131-257.

(3) Fukuzawa, A.; Masamune, T. Tetrahedron Lett. 1981, 41, 4081

(4) Guella, G.; Mancini, I.; Chasera, G.; Pietra, F. Helv. Chim. Acta 1992, 75, 310.

(5) For the isolation of other Laurencia oxepanes, see: (a) Kurosawa, A.; Irie, T. Tetrahedron Lett. 1973, 4135. (b) Wratten, S. J.; Faulkner, D. J. J. Org. Chem. 1977, 42, 3343. (c) Suzuki, M.; Kurata, K.; Suzuki, T.; Kurosawa, E. Bull. Chem. Soc. Jpn. 1986, 59, 2953. (d) Guella, G.; Pietra, F. Helv. Chim. Acta 1991, 74, 47. (e) Guella, G.; Mancini, I.; Chiasera, G.; Pietra, F. Ibid. 1992, 75, 303.

(6) For previous model studies in this area, see: (a) Kadota, I.; Kotsuki, H.; Ochi, M.; Ushio, Y. J. Org. Chem. 1989, 54, 5153. (b) Davies, M. J.; Moody, C. J. Synlett 1990, 95. (c) Davies, M. J.; Moody, C. J. J. Chem. Soc., Perkin Trans. 1 1991, 9. (d) Berger, D.; Overman. L. E. Synlett 1992, 811. (e) Carling, R. W.; Clark, J. S.; Holmes, A. B. J. Chem. Soc., Perkin Trans. ì 1992, 83

(7) Castañeda, A.; Kucera, D. J.; Overman, L. E. J. Org. Chem. 1989, 54, 5695

(8) Sharpless epoxidation<sup>9</sup> of commercially available (Z)-2-pentenol gives 5 in 78% yield and 80% ee. Enantiopure 5 is obtained by recrystallization (and subsequent hydrolysis) of the 3,5-dinitrobenzoate derivative.<sup>10</sup>

(9) Rossiter, B. E.; Katsuki, T.; Sharpless, K. B. J. Am. Chem. Soc. 1981, 103, 464. Evans, D. A.; Bender, S. L.; Morris, J. Ibid. 1988, 110, 2506. (10) Chong, J. M. Tetrahedron 1989, 45, 623.

Scheme I



opening of the epoxy sulfonate derivative according to the procedure of Murai,<sup>11</sup> provided bromohydrin 6 in 94% yield.<sup>12</sup> Conversion of 6 to the volatile bromoepoxide 7 was best accomplished in ether by treatment at low temperature with MeLi. Although this epoxide does not react cleanly with 2-(trimethylsilyl)-2-propenyl cuprates or silanes, it does afford the desired vinylsilane alcohol 9 in high yield upon reaction at low temperature with the allyltin reagent 8 in the presence of  $EtAlCl_2$ .<sup>13,14</sup>

The (R)- $\alpha$ -chloroacetal 13 was prepared from commercially available 2,2-dimethoxyethanol (10) by Swern oxidation followed by in situ reaction of glyoxal dimethyl acetal with 3-(triisopropylsiloxy)propynyllithium to provide propargylic alcohol 11 in 88% yield.<sup>15</sup> Jones oxidation of this intermediate followed by enantioselective reduction<sup>16</sup> of the derived ketone provided (S)-11 (80% ee).<sup>17-19</sup> Enantiopure (S)-11 could be obtained from this material by chromatographic separation (and subsequent

(12) All intermediates were fully characterized by <sup>1</sup>H and <sup>13</sup>C NMR, IR, and MS analysis. The elemental composition of analytical samples of new compounds was confirmed by combustion analysis or high-resolution mass spectrometry. Unless noted otherwise, yields refer to purified products.

(13) Allylstannane 8 was prepared in 54% overall yield from 1-(bromovinyl) trimethylsilane by the following sequence: (a) Mg, THF; HCHO; (b) n-BuLi, THF, -60 °C; MsCl; (c) LiSnBu, THF, -78 °C. For a related sequence, see: Lee, E.; Yu, S.-G.; Hur, C.-U.; Yang, S.-M. Tetrahedron Lett. 1988, 29, 6969

(14) Our investigations of the somewhat idiosyncratic reaction of  $\alpha$ -halo epoxides with 2-silyl-2-propenyl organometallics will be published shortly: Overman, L. E.; Renhowe, P. A., manuscript in preparation.

(15) Bernard, D.; Doutheau, A.; Gore, J. Synth. Commun. 1987, 17, 1807.
(16) Noyori, R.; Tomino, I.; Tanimoto, Y.; Nishizawa, M. J. Am. Chem. Soc. 1984, 106, 6709.

(17) The  $\alpha$ -acetal group apparently partially erodes enantioselection. Other catalytic and stoichiometric reductants (commonly used for the enantioselective reduction of ynones) that we screened gave lower enantioselectivities.

(18) The absolute configuration of (S)-11 (which is opposite to that predicted by the typical model for stereoselective ketone reductions with BINAL-H<sup>16</sup> and undoubtedly results from the presence of the  $\alpha$ -acetal functionality) was determined by chemical correlation with 1-methoxy-5-(triisopropylsiloxy)-2-pentanol. The comparison sample of (2R)-1-methoxy-5-(triisopropylsiloxy)-2-pentanol was prepared from 1-(triisopropylsiloxy)-4-pentene by the following sequence: (a) Sharpless asymmetic dihydroxylation (AD-mix- $\beta$ ),<sup>19</sup> t-BuOH, H2O, 0 °C; (b) MeI, NaH, THF, 0 °C.
 (19) Jacobsen, E. N.; Markó, I.; Mungall, W. S.; Schröder, G.; Sharpless,

K. B. J. Am. Chem. Soc. 1988, 110, 1968.

<sup>(11)</sup> Gao, L.; Saitoh, H.; Feng, F.; Murai, A. Chem. Lett. 1991, 1787.

Scheme II



cleavage) of the diastereomeric carbamates derived from (S)-(-)- $\alpha$ -methylbenzyl isocyanate.<sup>20</sup> Catalytic hydrogenation of enantiomerically pure (S)-11 then afforded enantiopure 12 in 44% overall yield from racemic propargylic alcohol 11. Conversion of 12 to the triflate derivative followed by reaction of this intermediate with (n-Bu)<sub>4</sub>NCl provided the (R)- $\alpha$ -chloroacetal 13 in 71% yield.<sup>21</sup>

The total synthesis of (+)-isolaurepinnacin was efficiently completed from the optically active fragments 9 and 13 as summarized in Scheme II. Treatment of acetal 13 at -78 °C with Me<sub>2</sub>BBr, followed by removal of Me<sub>2</sub>BOMe under reduced pressure, provided the corresponding  $\alpha$ -bromo ether 14.<sup>22</sup> This crude intermediate was directly combined with alcohol 9 in the presence of AgOTf to afford the mixed acetal 15 in high yield.<sup>6d,23</sup> Cyclization of this polyfunctional intermediate was best accomplished in CH<sub>2</sub>Cl<sub>2</sub> by reaction with 1.4 equiv of BCl<sub>3</sub> at -78  $\rightarrow$ 0 °C.<sup>6d</sup> After desilylation of the crude cyclization product, *oxepene 17 was isolated in 90% overall yield from the*  $\beta$ -*chloroacetal*. This remarkably selective conversion proceeds by way of the  $\alpha$ -chloro ether intermediate 16,<sup>24</sup> which can be detected (and isolated) at short reaction times. The (E)-enyne side chain

was then developed by initial oxidation of 17 with the Dess-Martin reagent,<sup>25</sup> followed by direct reaction of the resulting aldehyde with ethynylmagnesium bromide to afford 18. Dehydration of this intermediate under a variety of standard conditions was not stereoselective. However, dehydration of the hexacarbonyldicobalt complex of 18 proceeded with high (>24:1) (E) stereoselection.<sup>26</sup> The optimal procedure was to initially convert 18 to the corresponding hexacarbonyldicobalt complex, which upon subsequent reaction in CH<sub>2</sub>Cl<sub>2</sub> with Tf<sub>2</sub>O afforded the hexacarbonyldicobalt complex of isolaurepinnacin. Decomplexation of this intermediate with ceric ammonium nitrate provided (+)-isolaurepinnacin (1), contaminated with less than 4% of its (Z) stereoisomer, in 65% overall yield from 18. Synthetic 1 displays <sup>1</sup>H and <sup>13</sup>C NMR and IR spectra that are indistinguishable from those of the natural isolate and showed the following optical properties:  $[\alpha]^{25}_{D} + 0.6^{\circ}, [\alpha]^{25}_{546} + 1.2^{\circ}, [\alpha]^{25}_{405}$ +3.2° (c = 1.4, CHCl<sub>3</sub>).<sup>27</sup>

The first total synthesis of (+)-isolaurepinnacin has been achieved with high stereoselectivity in 12 steps and 15% overall yield from *cis*-2-penten-1-ol. This synthesis rigorously establishes the S configuration of 1 at C(13), which had previously been assigned on biosynthetic grounds, and corrects the rotation of 1 to be dextrorotatory.<sup>28,29</sup> Of particular note is the integrity of both bromine and chlorine functionalities during the Lewis acidpromoted cyclization of mixed acetal 15; this conversion highlights the extraordinary selectivity that can be realized in acetal-alkene cyclizations. The formation of a single stereoisomer in this key conversion moreover demonstrates that chiral  $\alpha$ -chlorooxocarbenium ions are sufficiently configurationally stable to not epimerize during a favorable acetal-alkene cyclization. This observation suggests that other oxacyclic marine natural products containing common 1-haloalkyl side chains can be accessed in asymmetric fashion by Prins-type cyclizations of  $\beta$ -haloacetals.

Acknowledgment. This investigation was supported by NSF Grant CHE-90-03812, and NMR and mass spectra were determined with spectrometers purchased with the assistance of NSF Shared Instrumentation Grants. We particularly thank Professor Akio Fukuzawa (Department of Engineering, Hokkaido Tokai University) for copies of spectra for natural isolaurepinnacin and for providing us with, and allowing us to report, unpublished information on the rotation of isolaurepinnacin and (3Z)isolaurepinnacin isolated from Laurencia pinnata Yamada.

Supplementary Material Available: Listings of spectroscopic and analytical data for new compounds and copies of <sup>1</sup>H NMR spectra of synthetic isolaurepinnacin (5 pages). Ordering information is given on any current masthead page.

<sup>(20)</sup> Pirkle, W. H.; Hauske, J. R. J. Org. Chem. 1977, 42, 1839. Pirkle, W. H.; Hauske, J. R. Ibid. 1977, 42, 2781.

<sup>(21)</sup> We have no direct measure of the enantiomeric purity of this intermediate. However, the high yield of a single stereoisomer observed in the key cyclization confirms that the enantiopurity of 13 was high.<sup>27</sup>

<sup>(22)</sup> Guindon, Y.; Yoakim, C.; Morton, H. E. J. Org. Chem. 1984, 49, 3912.

<sup>(23)</sup> Kronzer, F. J.; Schuerch, C. Carbohydr. Res. 1973, 27, 379.

<sup>(24)</sup> Goff, D. A.; Harris, R. N., III; Bottaro, J. C.; Bedford, C. D. J. Org. Chem. 1986, 51, 4711. Overman, L. E.; Thompson, A. S. J. Am. Chem. Soc. 1988, 110, 2248.

<sup>(25)</sup> Dess, D. B.; Martin, J. C. J. Am. Chem. Soc. 1991, 113, 7277.

<sup>(26)</sup> Nicholas, K. M.; Pettit, R. J. Organomet. Chem. 1972, 44, C21. Saksena, A. K.; Green, M. J.; Mangiaracina, P.; Wong, J. K.; Kreutner, W.; Gulbenkian, A. R. Tetrahedron Lett. 1985, 26, 6423. Melikyan, G. C.; Mineif, A.; Vostrowsky, O.; Bestmann, H. J. Synthesis 1991, 633.

<sup>(27)</sup> The C(6) epimer of 1, which is available from cyclization of the mixed acetal formed from 9 and rac-13, is distinguishable from 1 by  $^{13}$ C NMR and TLC comparisons.

<sup>(28)</sup> The small negative rotation,  $[\alpha]^{25}_{D}$  -6.2° (CHCl<sub>3</sub>), reported for 1 isolated from *Laurencia pinnata* Yamada is believed to be due to contamination from laurepinnacin,  $[\alpha]^{25}_{D}$  -35.3° (c = 1.1, CHCl<sub>3</sub>).<sup>29</sup> Co-occurring (3Z)-isolaurepinnacin showed a rotation of  $[\alpha]^{25}_{D}$  +2.0° (c = 1.0, CHCl<sub>3</sub>).<sup>29</sup> Desbromodeschlorooctahydroisolaurepinnacin prepared from synthetic (+)-1 was dextrorotatory and, thus, in accord with the absolute configuration assignments of Kotsuki for this isolaurepinnacin degradation product.<sup>6a</sup>

<sup>(29)</sup> Fukuzawa, A., personal communication to L.E.O. of June 10, 1993.