

## THE CHEMISTRY OF HETERO-ALLENE AND -ALLYLIC DERIVATIVES WITH RHODIUM AND IRIIDIUM

### II \*. RHODIUM(I)- AND IRIIDIUM(I)-PHOSPHINE COMPLEXES OF HETERO-ALLYLIC LIGANDS OF THE TYPE $[\text{Ph}_2\text{PC}(\text{X})\text{NR}]^-$ ( $\text{X} = \text{S}, \text{NR}, \text{O}$ ) AND $[\text{Ph}_2\text{P}(\text{Q})\text{C}(\text{S})\text{NR}]^-$ ( $\text{Q} = \text{S}, \text{O}$ ). SYNTHESIS AND $^{31}\text{P}$ -NMR

D.H.M.W. THEWISSEN \*\*, H.P.M.M. AMBROSIUS, H.L.M. VAN GAAL and J.J. STEGGERDA

*Department of Inorganic Chemistry, Catholic University, Toernooiveld, 6525 ED Nijmegen (The Netherlands)*

(Received November 12th, 1979)

#### Summary

Some known and new hetero-allylic derivatives  $[\text{Ph}_2\text{PC}(\text{X})\text{NR}]^-$  ( $\text{X} = \text{S}, \text{NR}, \text{O}$ ) and  $[\text{Ph}_2\text{P}(\text{Q})\text{C}(\text{S})\text{NR}]^-$  ( $\text{Q} = \text{S}, \text{O}$ ) have been shown to react with rhodium(I)- and iridium(I)-phosphine compounds to give stable complexes  $\text{M}(\text{PPh}_3)_2[\text{Ph}_2\text{PC}(\text{X})\text{NR}]$  (A) and  $\text{M}(\text{PPh}_3)_2[\text{Ph}_2\text{P}(\text{Q})\text{C}(\text{S})\text{NR}]$  (B) ( $\text{M} = \text{Rh}, \text{Ir}$ ) in which the ligands are bidentate. P, S(P) and O(P) are always coordinated and the coordination preference of the other atoms is:  $\text{S} > \text{NR} > \text{O}$ .

A and B react with carbon monoxide to give  $\text{M}(\text{PPh}_3)(\text{CO})[\text{Ph}_2\text{PC}(\text{X})\text{NR}]$  and  $\text{M}(\text{PPh}_3)(\text{CO})[\text{Ph}_2\text{P}(\text{Q})\text{C}(\text{S})\text{NR}]$ , respectively, in which CO is preferentially found *trans* to the chelate donor atom with the lowest *trans*-influence.

A correlation between  $\sqrt{{}^1J(\text{Rh}-\text{P}_1) \times {}^1J(\text{Rh}-\text{P}_2)}$  and  ${}^2J(\text{P}_1-\text{P}_2)_{\text{cis}}$  of 0.946 was found; an even better correlation with  $r = 0.988$  gave the equation:  ${}^2J(\text{P}_1-\text{P}_2)_{\text{cis}} = 0.43 \sqrt{{}^1J(\text{Rh}-\text{P}_1) \times {}^1J(\text{Rh}-\text{P}_2)} - 0.072 |{}^1J(\text{Rh}-\text{P}_1) - {}^1J(\text{Rh}-\text{P}_2)| - 30.8 \text{ Hz}$ . A similar correlation was shown by a number of asymmetric *cis*-Pt<sup>0</sup>-bis-phosphine ( $\eta^2$ -ligand) complexes and *trans*- and *cis*-WLL'-(CO)<sub>4</sub> complexes reported in the literature.

#### Introduction

By addition of  $\text{Ph}_2\text{P}-\text{H}$  or  $\text{Ph}_2\text{P}(\text{Q})-\text{H}$  ( $\text{Q} = \text{S}, \text{O}$ ) to the hetero-allene molecules  $\text{S}=\text{C}=\text{NR}$  ( $\text{R} = \text{Ph}, \text{Me}$ ), *p*-tolN=C=N*p*-tol and  $\text{O}=\text{C}=\text{NPh}$ , hetero-allylic

\* For part I see ref. 1.

\*\* Present address: O.C.I., T.N.O., Croesestraat 79, Utrecht, The Netherlands.




Fig. 1. The ambidentate hetero-allylic chelating ligands  $[X-C(Z)-Y]^-$ , used in this paper; LI, *P,P*-diphenyl-*N*-phenyl-phosphinothioformamido; LII, *P,P*-diphenyl-*N*-methyl-phosphinothioformamido; LIII, *P,P*-diphenyl-*N,N*-di-*p*-tolyl-phosphinoformamidinato; LIV, *P,P*-diphenyl-*N*-phenyl-phosphinoformamido; LV, *P,P*-diphenyl-*N*-phenyl-thiophosphinylthioformamido; LVI, *P,P*-diphenyl-*N*-phenyl-phosphinylthioformamido.

derivatives result with the general formula  $Ph_2PC(X)NHR$  ( $X = S, NR, O$ ) and  $Ph_2P(Q)C(S)NHR$  ( $Q = S, O$ ). The synthesised ligands are shown in Fig. 1 in their deprotonated forms.

These ligands are ambidentate, because of the presence of three hetero atoms with coordinating properties. The mode of coordination of the ambidentate ligands can be determined by means of IR and  $^{31}P$  NMR spectral parameters.

Gal et al. [2,3] reported the complexes  $Rh(PPh_3)_2(LI)$  and  $Rh(PPh_3)_2(LVI)$ . LI was found to coordinate to rhodium through P and S, giving a four-membered chelate ring complex, whereas LVI coordinated through O(P) and S, resulting in a five-membered chelate ring complex.

## Experimental

IR spectra were measured on a Perkin-Elmer 283 spectrophotometer ( $4000-200\text{ cm}^{-1}$ ), in CsI pellets.  $^{31}P\{^1H\}$  NMR spectra were recorded on a Varian XL-1000 FT spectrometer at 40.5 MHz, using the deuterated solvent as internal lock. Solutions for NMR measurements were prepared in a glove-box.

C, H and N analyses were carried out at the microanalytical department of this university. Analytical data are given in Table 1.

Reactions were carried out at room-temperature in analytical grade solvents under nitrogen.

$RhCl(PPh_3)_3$  [4],  $[IrCl(C_8H_{14})_2]_2$  [5],  $Ph_2PC(S)NHR$  ( $R = Ph, Me$ ) [6],  $Ph_2PC-(Np\text{-}tol)NHp\text{-}tol$  [7],  $Ph_2PC(O)NHPh$  [8],  $Ph_2P(S)C(S)NHPh$  and  $Ph_2P(O)C(S)NHPh$  [9] were prepared according to literature procedures.

### $Rh(PPh_3)_2[X-C(Z)-Y]$ (IIa, IVa, Va)

0.3 mmol of the protonated ligand was added to a solution of 0.3 mmol  $RhCl-(PPh_3)_3$  in 20 ml benzene. After 20 minutes a small excess of  $Et_3N$  was added and after 12 hours the  $Et_2N \cdot HCl$  was filtered off. The complex was precipitated with *n*-hexane, filtered off, washed with ethanol and diethyl ether, and dried in vacuo. [ $^1H$  NMR IIa:  $\tau(Me)$  7.00 ppm (s), 7.23 ppm (s)].

*Rh(PPh<sub>3</sub>)<sub>2</sub>[Ph<sub>2</sub>PC(Np-tol)Np-tol] (IIIa)*

The initial red solution from 0.3 mmol RhCl(PPh<sub>3</sub>)<sub>3</sub> and 0.3 mmol Ph<sub>2</sub>PC-(Np-tol)NHp-tol in 20 ml benzene changed to orange-red upon addition of an equimolar amount n-BuLi. The complex was precipitated with n-hexane, filtered off, washed with ethanol and diethylether and dried in vacuo. [<sup>1</sup>H NMR:  $\tau$ (Me) 8.16 ppm (s), 8.22 ppm (s)].

*Ir(PPh<sub>3</sub>)<sub>2</sub>[X-C(Z)-Y] (VIIa, IXa, Xa, XIa)*

To a solution of 0.15 mmol [IrCl(C<sub>8</sub>H<sub>14</sub>)<sub>2</sub>]<sub>2</sub> and 0.6 mmol PPh<sub>3</sub> in 30 ml benzene 0.3 mmole of the protonated ligand was added. After 20 minutes a small excess Et<sub>3</sub>N was added. The initial red colour changed in three hours to yellow. After removal Et<sub>3</sub>N · HCl by filtration the complex was precipitated with n-hexane, filtered off, washed with small amounts ethanol and diethylether, and dried in vacuo.

*Ir(PPh<sub>3</sub>)<sub>2</sub>[Ph<sub>2</sub>PC(Np-tol)Np-tol] (VIIIa)*

This complex could not be isolated. On addition of an equimolar amount n-BuLi to a solution of 0.15 mmol [IrCl(C<sub>8</sub>H<sub>14</sub>)<sub>2</sub>]<sub>2</sub>, 0.6 mmol PPh<sub>3</sub> and 0.3 mmol Ph<sub>2</sub>PC(Np-tol)NHp-tol in 30 ml benzene, VIIIa was formed, as could be proved by IR measurements of the reaction-mixture. However, VIIIa was not stable and a subsequent elimination of *p*-tolN=C=Np-tol occurred, as will be described elsewhere [10].

TABLE 1  
ANALYTICAL DATA  
(Theoretical values in parentheses)

No.	Compound	colour	Found (calcd.) (%)			yield (%)
			C	H	N	
IIa	Rh(PPh <sub>3</sub> ) <sub>2</sub> [Ph <sub>2</sub> PC(S)NMe]	yellow	67.9 (67.4)	5.2 (5.0)	1.7 (1.6)	75
IIIa	Rh(PPh <sub>3</sub> ) <sub>2</sub> [Ph <sub>2</sub> PC(Np-tol)Np-tol]	yellow-orange	72.1 (73.1)	5.4 (5.3)	2.5 (2.7)	70
IVa	Rh(PPh <sub>3</sub> ) <sub>2</sub> [Ph <sub>2</sub> PC(NPh)O]	yellow-orange	69.0 (68.5)	5.2 (5.3)	1.5 (1.6)	70
Va	Rh(PPh <sub>3</sub> ) <sub>2</sub> [Ph <sub>2</sub> P(S)C(S)NPh]	yellow	68.2 (67.4)	5.0 (4.6)	1.3 (1.4)	70
Vb	Rh(PPh <sub>3</sub> )(CO)[Ph <sub>2</sub> P(S)C(S)NPh]	orange	60.4 (61.2)	4.1 (4.1)	2.0 (1.9)	85
VIIa	Ir(PPh <sub>3</sub> ) <sub>2</sub> [Ph <sub>2</sub> PC(S)NPh]	yellow-brown	64.0 (63.7)	4.7 (4.4)	1.3 (1.4)	70
VIIb	Ir(PPh <sub>3</sub> )(CO)[Ph <sub>2</sub> PC(S)NPh]	orange	55.7 (56.9)	4.1 (3.8)	1.8 (1.7)	75
IXa	Ir(PPh <sub>3</sub> ) <sub>2</sub> [Ph <sub>2</sub> PC(NPh)O]	yellow-brown	62.1 (64.7)	4.9 (4.4)	1.5 (1.4)	45
Xa	Ir(PPh <sub>3</sub> ) <sub>2</sub> [Ph <sub>2</sub> P(S)C(S)NPh]	yellow-orange	60.2 (61.8)	4.5 (4.2)	1.4 (1.3)	40
Xb	Ir(PPh <sub>3</sub> )(CO)[Ph <sub>2</sub> P(S)C(S)NPh]	yellow	57.0 (54.7)	3.9 (3.6)	1.5 (1.7)	35
XIa	Ir(PPh <sub>3</sub> ) <sub>2</sub> [Ph <sub>2</sub> P(O)C(S)NPh]	yellow	62.6 (62.7)	4.3 (4.3)	1.2 (1.3)	40
XIb	Ir(PPh <sub>3</sub> )(CO)[Ph <sub>2</sub> P(O)C(S)NPh]	yellow	56.0 (55.7)	4.0 (3.7)	1.5 (1.7)	35

$M(PPh_3)(CO)[X-C(Z)-Y]$  (Vb, VIIb, Xb, XIb)

On passing CO through a solution of 0.3 mmol  $M(PPh_3)_2[X-C(Z)-Y]$  in 30 ml benzene the colour of the reaction mixture rapidly changed from yellow to orange-yellow. Dilution with n-hexane and cooling gave an orange or yellow precipitate.

The analogous complexes IIIb, IVb, VIIIb and IXb could not be prepared. IIIa, IVa and IXa reacted differently with CO [10]. Yields are given in Table 1. The Ir complexes were more difficult to obtain and were much less stable than the Rh analogs.

## Results and discussion

### Reactions and products

The  $Rh^I$  and  $Ir^I$  complexes  $M(PPh_3)_2[X-C(Z)-Y]$  can be prepared either by a metathetical substitution using the Li salts of the ligands or by means of a N—H bond breaking reaction activated by a base, e.g.  $Et_3N$ .

The complex  $Ir^{III}(H)(Cl)(PPh_3)_2[Ph_2PC(S)NPh]$  is one of the products of the reaction between  $[IrCl(C_8H_{14})_2]_2$ ,  $PPh_3$  and  $Ph_2PC(S)NPh$  without addition of  $Et_3N$ . This indicates that the N—H bond breaking is an oxidative addition reaction. The N—H bond breaking does not occur with  $Ph_2PC(Np-tol)NHp-tol$ . Probably this ligand is not sufficiently acidic to give a facile N—H bond breaking.  $[IrCl(C_8H_{14})_2]_2$  reacts with  $[Ph_2PC(Np-tol)NHp-tol]^-Li^+$  to give the intermediate  $\{Ir(PPh_3)_2[Ph_2PC(Np-tol)NHp-tol]\}$ , as can be observed from the IR spectrum of the reaction mixture. A subsequent elimination reaction occurs, and this will be discussed later [10].

The carbonyl complexes of the type  $M(PPh_3)(CO)[X-C(Z)-Y]$  can be prepared by passing CO through a solution of  $M(PPh_3)_2[X-C(Z)-Y]$  one phosphine being replaced by CO. In the case of  $Rh(PPh_3)_2[Ph_2PC(Np-tol)NHp-tol]$  and  $M(PPh_3)_2[Ph_2PC(O)NPh]$ , however, a second reaction takes place [10]. This subsequent elimination also occurs to some extent in the reaction of  $Ir(PPh_3)_2[Ph_2P(S)C(S)NPh]$  and  $Ir(PPh_3)_2[Ph_2P(O)C(S)NPh]$  with CO. The occurrence of this second reaction may explain the fairly low yields in the syntheses of both Ir complexes.




Fig. 2. Formation of  $Ir(PPh_3)_2[Ph_2PC(S)NPh]$  from  $IrCl(PPh_3)_2(C_8H_{14})$  and  $Ph_2PC(S)NPh$  via an oxidative addition reaction.

## IR spectra and mode of coordination

Table 2 shows the relevant absorptions in the IR spectra between 2000 and 500  $\text{cm}^{-1}$ . In these complexes a strong absorption can be attributed to a normal mode having  $\nu(\text{C}\equiv\text{E})$  of the exocyclic  $\text{C}=\text{E}$  band as the main component [11–14]. Two regions are found for this absorption: 1500–1575  $\text{cm}^{-1}$ , attributed to  $\nu(\text{C}\equiv\text{N})$ , and 1620–1640  $\text{cm}^{-1}$ , assigned to  $\nu(\text{C}\equiv\text{O})$ . This establishes for all complexes the mode of bidentate coordination of the ambidentate ligands.

Coordination by P and S is observed for the complexes of  $[\text{Ph}_2\text{PC}(\text{S})\text{NR}]^-$  ( $\text{R} = \text{Ph}, \text{Me}$ ). The ligand  $[\text{Ph}_2\text{PC}(\text{Np-tol})\text{Np-tol}]^-$  is obviously coordinated by P and N. For the complexes of  $[\text{Ph}_2\text{PC}(\text{X})\text{NR}]^-$  ( $\text{X} = \text{S}, \text{NR}$ )  $\nu(\text{C}\equiv\text{N})$  is found in the region 1547–1575  $\text{cm}^{-1}$ , whereas the complexes of  $[\text{Ph}_2\text{PC}(\text{NPh})\text{O}]^-$  have  $\nu(\text{C}\equiv\text{N})$  at about 1620  $\text{cm}^{-1}$ , which is assigned to  $\nu(\text{C}\equiv\text{O})$ . Therefore  $[\text{Ph}_2\text{PC}(\text{NPh})\text{O}]^-$  is coordinated by P and N.

The ligands  $[\text{Ph}_2\text{P}(\text{Q})\text{C}(\text{S})\text{NPh}]^-$  ( $\text{Q} = \text{S}, \text{O}$ ) have  $\nu(\text{C}\equiv\text{N})$  in the region 1509–1533  $\text{cm}^{-1}$  and form five-membered chelate rings as is also indicated by the values of  $\nu(\text{P}=\text{Q})$ . In the spectra of  $\text{M}(\text{PPh}_3)_2(\text{L})[\text{Ph}_2\text{P}(\text{S})\text{C}(\text{S})\text{NPh}]$  ( $\text{M} = \text{Rh}, \text{Ir}$ ;  $\text{L} = \text{PPh}_3, \text{CO}$ ) new absorptions of medium intensity are observed in the region 602–625  $\text{cm}^{-1}$  and of strong intensity between 512–520  $\text{cm}^{-1}$ . These absorptions are assigned to  $\nu(\text{P}=\text{S})$ . For complexes of the ligand  $\text{R}_2\text{P}(\text{S})\text{CSS}^-$   $\nu(\text{P}=\text{S})$  was reported in the range 580–610  $\text{cm}^{-1}$ ; for  $\text{Pd}[\text{Ph}_2\text{P}(\text{S})\text{C}(\text{S})\text{S}]_2$   $\nu(\text{P}=\text{S})$  was assigned at 601  $\text{cm}^{-1}$  [15], and  $\text{Zn}[\text{Cy}_2\text{P}(\text{S})\text{C}(\text{S})\text{S}]_2$  at 597  $\text{cm}^{-1}$  [16]. In complexes of the ligand  $\text{R}_2\text{PS}^-$  and  $\text{R}_2\text{PHS}$   $\nu(\text{P}=\text{S})$  was assigned at 520  $\text{cm}^{-1}$  for  $[\text{Mn}(\text{CO})_4(\mu\text{-Me}_2\text{PS})]_2$  [22], and at 495  $\text{cm}^{-1}$  for  $\text{Mo}(\text{CO})_5(\text{SPHMe}_2)$  [23].  $\nu(\text{P}=\text{O})$  decreases 45–50  $\text{cm}^{-1}$  on complexation. The mode of coordination to form a five membered chelate ring complex was already established for  $[\text{Ph}_2\text{P}(\text{O})\text{C}(\text{S})\text{NPh}]^-$  [2].

$\nu(\text{PCS})$  is observed from 920–930  $\text{cm}^{-1}$  for the four ring complexes, whereas the five ring compounds exhibit  $\nu(\text{PCS})$  in the region 952–970  $\text{cm}^{-1}$ .

In the carbonyl complexes the  $\text{C}\equiv\text{O}$  absorptions are observed from 1950–1987  $\text{cm}^{-1}$ , which is normal for four coordinate Rh and Ir complexes. The value of  $\nu(\text{CO})$  is somewhat higher for the Rh complex than for the analogous Ir compound.

Summarising, we have found that P, S(P) and O(P) are always coordinated and that the coordination preference of the other groups is  $\text{S} > \text{NR} > \text{O}$ .

## $^{31}\text{P}$ NMR spectra and correlation between coupling constants

The measured  $^{31}\text{P}\{^1\text{H}\}$  NMR parameters are given in Table 3. The values are not corrected for higher order effects. The spectra of  $\text{Rh}(\text{PPh}_3)_2[\text{Ph}_2\text{PC}(\text{Np-tol})\text{Np-tol}]$  and  $\text{Rh}(\text{PPh}_3)_2[\text{Ph}_2\text{P}(\text{S})\text{C}(\text{S})\text{NPh}]$  are shown in Figs. 3 and 4, respectively.

Assignments are as given in Table 3. All the complexes contain two  $\text{PPh}_3$  ligands *cis* to each other. As observed by Gal et al. for five of the complexes [2], the  $\text{PPh}_3$  group with the highest value of  $^1J(\text{Rh}-\text{P})$ , i.e. the phosphine *trans* to the hetero atom with the lowest *trans*-influence, is displaced by a  $\pi$ -acceptor ligand such as CO. In the four-membered chelate ring complexes the phosphine *trans* to the hetero atom S or NR is displaced by CO, whereas in Va and VIa the

TABLE 2

IR ABSORPTIONS (cm<sup>-1</sup>)

No.	Complex	$\nu(\text{C}\equiv\text{O})$	$\nu(\text{C}=\text{E})$	$\nu(\text{P}=\text{Z})$	$\nu(\text{P}-\text{C}-\text{S})$	$\nu(\text{N}-\text{Ph})$
Ia	Rh(PPh <sub>3</sub> ) <sub>2</sub> [Ph <sub>2</sub> PC(S)NPh]		1547vs ( $\nu(\text{C}=\text{N})$ )		925w	803w
Ib	Rh(PPh <sub>3</sub> ) <sub>2</sub> (CO)[Ph <sub>2</sub> PC(S)NPh]	1975vs	1572s ( $\nu(\text{C}=\text{N})$ )		930m	803w
Ila	Rh(PPh <sub>3</sub> ) <sub>2</sub> [Ph <sub>2</sub> PC(S)NMe]		1575vs ( $\nu(\text{C}=\text{N})$ )		927m	
IIla	Rh(PPh <sub>3</sub> ) <sub>2</sub> [Ph <sub>2</sub> PC(Np-tol)Np-tol]		1552vs ( $\nu(\text{C}=\text{N})$ )			
IVa	Rh(PPh <sub>3</sub> ) <sub>2</sub> [Ph <sub>2</sub> PC(NPh)O]		1624vs ( $\nu(\text{C}=\text{O})$ )			808w
Va	Rh(PPh <sub>3</sub> ) <sub>2</sub> [Ph <sub>2</sub> P(S)C(S)NPh]		1520s ( $\nu(\text{C}=\text{N})$ )	625m ( $\nu(\text{P}=\text{S})$ )	960m	798w
Vb	Rh(PPh <sub>3</sub> ) <sub>2</sub> (CO)[Ph <sub>2</sub> P(S)C(S)NPh]	1987vs	1524s ( $\nu(\text{C}=\text{N})$ )	612m ( $\nu(\text{P}=\text{S})$ )	952m	800w
VIa	Rh(PPh <sub>3</sub> ) <sub>2</sub> [Ph <sub>2</sub> P(O)C(S)NPh]		1511s ( $\nu(\text{C}=\text{N})$ )		968m	802m
VIIa	Rh(PPh <sub>3</sub> ) <sub>2</sub> (CO)[Ph <sub>2</sub> P(O)C(S)NPh]	1980vs	1533s ( $\nu(\text{C}=\text{N})$ )	1135s ( $\nu(\text{P}=\text{O})$ )	968m	802w
VIIb	Ir(PPh <sub>3</sub> ) <sub>2</sub> [Ph <sub>2</sub> PC(S)NPh]		1552vs ( $\nu(\text{C}=\text{N})$ )	1130s ( $\nu(\text{P}=\text{O})$ )	922m	801m
VIIIa	Ir(PPh <sub>3</sub> ) <sub>2</sub> (CO)[Ph <sub>2</sub> PC(S)NPh]	1955vs	1555vs ( $\nu(\text{C}=\text{N})$ )		920m	804w
IXa	Ir(PPh <sub>3</sub> ) <sub>2</sub> [Ph <sub>2</sub> PC(Np-tol)Np-tol] <sup>a</sup>		1562vs ( $\nu(\text{C}=\text{N})$ )			
Xa	Ir(PPh <sub>3</sub> ) <sub>2</sub> [Ph <sub>2</sub> PC(NPh)O]		1621vs ( $\nu(\text{C}=\text{O})$ )			806w
Xb	Ir(PPh <sub>3</sub> ) <sub>2</sub> [Ph <sub>2</sub> P(S)C(S)NPh]		1509s ( $\nu(\text{C}=\text{N})$ )	610m ( $\nu(\text{P}=\text{S})$ )	957m	795w
XIa	Ir(PPh <sub>3</sub> ) <sub>2</sub> (CO)[Ph <sub>2</sub> P(S)C(S)NPh]	1951vs	1529s ( $\nu(\text{C}=\text{N})$ )	602m ( $\nu(\text{P}=\text{S})$ )	951m	800w
XIb	Ir(PPh <sub>3</sub> ) <sub>2</sub> [Ph <sub>2</sub> P(O)C(S)NPh]		1512s ( $\nu(\text{C}=\text{N})$ )	1132s ( $\nu(\text{P}=\text{O})$ )	970m	802m
XIIb	Ir(PPh <sub>3</sub> ) <sub>2</sub> (CO)[Ph <sub>2</sub> P(O)C(S)NPh]	1947vs	1517s ( $\nu(\text{C}=\text{N})$ )	1140s ( $\nu(\text{P}=\text{O})$ )	965m	803w

Spectra measured in Cal pellets (<sup>a</sup> measured in benzene solution).




Fig. 3. The  $^{31}\text{P}$   $\{^1\text{H}\}$  NMR spectrum of  $\text{Rh}(\text{PPh}_3)_2[\text{Ph}_2\text{PC}(\text{Np-tol})\text{Np-tol}]$ .

$\text{PPh}_3$  ligand *trans* to S(P) and O(P) is replaced. It is noteworthy that the spectrum of Va differs slightly from that of VIa. The  $^2J(\text{Rh-Q-P}_3)$  coupling is 3 Hz in VIa and unobserved in Va, whereas the  $^3J(\text{P}_1\text{-Rh-Q-P}_3)$  coupling is unobserved in VIa and 3 Hz in Va.

Summarised,  $^1J(\text{Rh-P}_i)$  ( $i = 1, 2$ ) is observed in the range 164–202 Hz for the  $\text{PPh}_3$  ligand that is substituted by CO, and 148–170 Hz for the phosphine which is not displaced, while  $^1J(\text{Rh-P}_3)$  for the ligand  $\text{PPh}_2$ -group lies in the range 102–124 Hz.  $^2J(\text{P}_1\text{-P}_2)_{\text{cis}}$  varies from 35 to 47 Hz in the Rh complexes and from 14 to 22 in the Ir analogues, whereas  $^2J(\text{P}_2\text{-P}_3)_{\text{cis}}$  is observed from 24 to 36 Hz.  $^2J(\text{P}_2\text{-P}_3)_{\text{trans}}$  in the four membered chelate ring complexes is found from 300–333 Hz for the Rh complexes and amounts to 340 Hz for  $\text{Ir}(\text{PPh}_3)_2[\text{Ph}_2\text{PC}(\text{S})\text{NPh}]$ .

There is an approximate linear relationship between  $^1J(\text{Rh-P}_i)$  and  $\delta(\text{P}_i)$ , the correlation coefficient being 0.844. Gal et al. have attributed such a relationship to the *trans*-influence of the various hetero atoms.

It has been mentioned that variations in  $^1J(\text{M-P})$  and  $^2J(\text{P-P})$  are expected to parallel each other [17–19]. Indeed  $^1J(\text{Rh-P}_i)$  and  $^2J(\text{P}_1\text{-P}_2)_{\text{cis}}$  exhibit similar





Fig. 4. The  $^{31}\text{P}$   $\{^1\text{H}\}$  NMR spectrum of  $\text{Rh}(\text{PPh}_3)_2[\text{Ph}_2\text{P}(\text{S})\text{C}(\text{S})\text{NPh}]$ .

trends. Therefore, we have examined the correlations for the set of asymmetric complexes of the type  $\text{Rh}(\text{PPh}_3)_2[\text{X}-\text{C}(\text{Z})-\text{Y}]$  ( $\text{X} \neq \text{Y}$ ) described in this paper and the related compounds reported by Gal et al. [2], assuming the same sign for all complexes (positive).



$^1J(\text{Rh}-\text{P}_1)$  and  $^2J(\text{P}_1-\text{P}_2)_{\text{cis}}$  gave a low correlation coefficient of 0.45. However, the correlation between the geometric mean  $\sqrt{^1J(\text{Rh}-\text{P}_1) \times ^1J(\text{Rh}-\text{P}_2)}$  and  $^2J(\text{P}_1-\text{P}_2)_{\text{cis}}$  was 0.946, implying that  $^2J(\text{P}_1-\text{P}_2)$  is related to both of the  $^1J$  coupling constants. The relationship between  $^2J(\text{P}_1-\text{P}_2)_{\text{cis}}$  and  $\sqrt{^1J(\text{Rh}-\text{P}_1) \times ^1J(\text{Rh}-\text{P}_2)}$  is plotted in figure 5 [x]. The plot reveals that  $^2J(\text{P}_1-\text{P}_2)_{\text{cis}}$  is also a






Fig. 5. Plot of  ${}^2J(P_1-P_2)_{cis}$  versus  $\sqrt{{}^1J(Rh-P_1) \times {}^1J(Rh-P_2)}$  [X] and of  ${}^2J(P_1-P_2)_{cis} + 0.072|{}^1J(Rh-P_1) - {}^1J(Rh-P_2)|$  versus  $\sqrt{{}^1J(Rh-P_1) \times {}^1J(Rh-P_2)}$  [O]. Solid line calculated, using equation 1 in the form:

$${}^2J(P_1-P_2)_{cis} + \frac{b}{\mu}|{}^1J(Rh-P_1) - {}^1J(Rh-P_2)| = \frac{a}{\mu}\sqrt{{}^1J(Rh-P_1) \times {}^1J(Rh-P_2)} - c$$

with  $\frac{a}{\mu} = 0.43$ ,  $\frac{b}{\mu} = 0.072$  and  $c = 30.8$  Hz. Complexes: 1,  $Rh(PPh_3)_2[Me_2NC(S)NPh]$ ; 2,  $Rh(PPh_3)_2[Ph_2PC(S)S]$ ; 3,  $Rh(PPh_3)_2[Ph_2PC(S)NPh]$ ; 4,  $Rh(PPh_3)_2[Ph_2PC(S)NMe]$ ; 5,  $Rh(PPh_3)_2[Ph_2P(O)C(S)NPh]$ ; 6,  $Rh(PPh_3)_2[Ph_2P(S)C(S)NPh]$ ; 7,  $Rh(PPh_3)_2[Ph_2PC(Np-tol)Np-tol]$ ; 8,  $Rh(PPh_3)_2[Ph_2PC(NPh)O]$ ; 9,  $RhCl(PPh_3)_3$  [24]; 10,  $RhBr(PPh_3)_3$  [24]; 11,  $RhI(PPh_3)_3$  [24].

function of the absolute value of the difference of the two coupling constants  ${}^1J(Rh-P_1)$  and  ${}^1J(Rh-P_2)$ , which is a measure of the asymmetry within the complex, so that an empirical relation between  ${}^1J(M-P_1)$ ,  ${}^1J(M-P_2)$  and  ${}^2J(P_1-P_2)$  may be given:

$${}^2J(P_1-P_2)_{cis} = \frac{1}{\mu}[\alpha\sqrt{{}^1J(M-P_1) \times {}^1J(M-P_2)} - b|{}^1J(M-P_1) - {}^1J(M-P_2)|] - c \quad (1)$$

$\mu$  represents the nuclear magnetic moment of the metal in nuclear magnetons;  $a$ ,  $b$  and  $c$  are empirical constants. For the set of asymmetric *cis*- $Rh^I$ -bisphosphine complexes we computed:  $a/\mu = 0.43$ ,  $b/\mu = 0.072$  and  $c = 30.8$  Hz, with a correlation-coefficient  $r = 0.988$ .

The existence of such a correlation is confirmed by the similar behaviour of ten iso-structural asymmetric *cis*- $Pt^0$ -bisphosphine- $(\eta^2\text{-ligand})$  complexes, the NMR parameters of which are reported by Meij et al. [20–21]. In Fig. 6 the plot of  ${}^2J(P_1-P_2)_{cis}$  versus  $\sqrt{{}^1J(Pt-P_1) \times {}^1J(Pt-P_2)}$  is shown. Again signs are taken as positive, except  ${}^2J(P_1-P_2)_{cis}$  of compounds 3 and 4. If the signs for  ${}^2J(P_1-P_2)_{cis}$  of compounds 3 and 4 are not taken as negative, no acceptable correlation results.

The correlation between  ${}^2J(P_1-P_2)_{cis}$  and  ${}^1J(Pt-P_i)$  ( $i = 1, 2$ ) has a correlation coefficient of 0.46; the value for the correlation between  ${}^2J(P_1-P_2)_{cis}$  and

TABLE 3  
 $^{31}\text{P}\{^1\text{H}\}$  NMR PARAMETERS

Complex	Number	L	$\delta_1$	$^1J(\text{Rh}-\text{P}_1)$	group <i>trans</i>	$\delta_2$
	Ia	PPh <sub>3</sub>	-41.5	164	S	-32.6
	Ib	CO				-31.2
	IIa	PPh <sub>3</sub>	-41.5	164	S	-33.6
	IIIa	PPh <sub>3</sub>	-48.0	166	N	-32.7
	IVa	PPh <sub>3</sub>	-47.7	169	N	-32.4
	Va	PPh <sub>3</sub>	-43.4	186	S(P)	-39.3
	Vb	CO				-41.4
	VIa	PPh <sub>3</sub>	-54.8	202	O(P)	-40.9
	VIB	CO				-35.6
	VIIa	PPh <sub>3</sub>	-19.4		S	-21.3
	Xa	PPh <sub>3</sub>	-33.1		S(P)	-19.7
	XIa	PPh <sub>3</sub>	-38.4		O(P)	-24.2

In ppm relative to  $\text{O}=\text{P}(\text{OMe})_3$  (TMP), internal reference; upfield = +;  $J$  in Hz. Measured in  $\text{CD}_2\text{Cl}_2$  solution.

<sup>a</sup> Decomposes slowly in  $\text{CD}_2\text{Cl}_2$ .

$\sqrt{{}^1J(\text{Pt}-\text{P}_1) \times {}^1J(\text{Pt}-\text{P}_2)}$  is 0.949. Using expression 1 we calculated a correlation coefficient of 0.990 with  $a/\mu = 0.077$   $b/\mu = 0.0097$  and  $c = 285$  Hz.

Although no theoretical interpretation is available a relationship such as expressed in equation 1 may have a fundamental basis.

The term  $\sqrt{{}^1J(\text{M}-\text{P}_1) \times {}^1J(\text{M}-\text{P}_2)}$  is a measure for the contribution of the s

$^1J(\text{Rh}-\text{P}_2)$	group <i>trans</i>	$\delta_3$	$^1J(\text{Rh}-\text{P}_3)$	group <i>trans</i>	$^nJ(\text{P}_1-\text{P}_2)$	$^nJ(\text{P}_1-\text{P}_3)$	$^nJ(\text{P}_2-\text{P}_3)$
148	-PPh <sub>2</sub>	+6.0	119	PPh <sub>3</sub>	35 <i>cis</i>	24 <i>cis</i>	330 <i>trans</i>
136	-PPh <sub>2</sub>	+7.8	102	PPh <sub>3</sub>			332 <i>trans</i>
147	-PPh <sub>2</sub>	+8.2	120	PPh <sub>3</sub>	35 <i>cis</i>	25 <i>cis</i>	333 <i>trans</i>
158	-PPh <sub>2</sub>	+6.6	119	PPh <sub>3</sub>	40 <i>cis</i>	32 <i>cis</i>	300 <i>trans</i>
160	-PPh <sub>2</sub>	+0.8	124	PPh <sub>3</sub>	40 <i>cis</i>	36 <i>cis</i>	302 <i>trans</i>
161	S	-34.4	0	PPh <sub>3</sub>	43 <i>cis</i>	3 <i>trans</i>	36 <i>cis</i>
175	S	-40.7	5	PPh <sub>3</sub>			24 <i>cis</i>
170	S	-28.6	3	PPh <sub>3</sub>	47 <i>cis</i>	0 <i>trans</i>	18 <i>cis</i>
152	S	-33.1	4	CO			17 <i>cis</i>
	-PPh <sub>2</sub>	+11.1		PPh <sub>3</sub>	20 <i>cis</i>	20 <i>cis</i>	342 <i>trans</i>
	S	-17.4		PPh <sub>3</sub>	30 <i>cis</i>	6 <i>trans</i>	26 <i>cis</i>
	S	-27.6		PPh <sub>3</sub>	19 <i>cis</i>		12 <i>cis</i>

orbitals to the two metal-ligand bonds, as  $^1J(\text{M}-\text{P}_1)$  and  $^1J(\text{M}-\text{P}_2)$  are mainly determined by the *s* character of the M-P bonds [25-27].  $^2J(\text{P}_1-\text{P}_2)_{\text{cis}}$  is determined by the metal *s* and *d* character of the M-P bonds in the case of  $\text{Rh}^{\text{I}}$  and by the *s*, *p* and *d* character in the case of  $\text{Pt}^0$ , because this coupling takes place via the metal-centre. The metal *s*- and *d*-orbitals have opposite contributions to




Fig. 6. Plot of  ${}^2J(P_1-P_2)_{cis}$  versus  $\sqrt{{}^1J(Pt-P_1) \times {}^1J(Pt-P_2)}$  [X] and of  ${}^2J(P_1-P_2)_{cis} + 0.0097 |{}^1J(Pt-P_1) - {}^1J(Pt-P_2)|$  versus  $\sqrt{{}^1J(Pt-P_1) \times {}^1J(Pt-P_2)}$  [O]. Solid line calculated, using equation 1 in the form:

$${}^2J(P_1-P_2)_{cis} + \frac{b}{\mu} |{}^1J(Pt-P_1) - {}^1J(Pt-P_2)| = \frac{a}{\mu} \sqrt{{}^1J(Pt-P_1) \times {}^1J(Pt-P_2)} - c$$

with  $\frac{a}{\mu} = 0.077$ ,  $\frac{b}{\mu} = 0.0097$  and  $c = 285$  Hz. Complexes: 1,  $Pt(PPh_3)_2(p\text{-tolN}=\text{S}=\text{N}p\text{-tol})$ ; 2,  $Pt(PPh_3)_2(p\text{-Cl-phenylN}=\text{S}=\text{N}p\text{-Cl-phenyl})$ ; 3,  $Pt(PPh_3)_2(3,5\text{-xylylN}=\text{S}=\text{N-3,5-xylyl})$ ; 4,  $Pt(PPh_3)_2CS_2$ ; 5,  $Pt(PPh_3)_2(PhN=\text{S}=\text{O})$ ; 6,  $Pt(PPh_3)_2(o\text{-tolN}=\text{S}=\text{O})$ ; 7,  $Pt(PPh_3)_2(o\text{-tolN}=\text{S}=\text{O})$ ; 8,  $Pt(PPh_3)_2(3,5\text{-xylylN}=\text{S}=\text{O})$ ; 9,  $Pt(PPh_3)_2(2,4,6\text{-mesitylN}=\text{S}=\text{O})$ ; 10,  $Pt(PPh_3)_2(PhN=\text{S}=\text{NPh})$  [20–21].

${}^2J(P_1-P_2)_{cis}$ , because of the node of the  $d$ -orbital (i.e. the  $d$ -orbital changes sign over  $90^\circ$ ).

The values of  $a$  (0.038) and  $b$  (0.0064) of the  $Rh^I$  series are comparable with the values of  $a$  (0.047) and  $b$  (0.0059) of the  $Pt^0$  series; the difference between  $a(Rh)$  and  $a(Pt)$  can be related to the different coordination number, which is four for  $Rh^I$  and three for  $Pt^0$ .

That  ${}^2J(P_1-P_2)_{cis}$  is close to zero indicates that the  $s$  and  $d$  contributions cancel each other. Such a cancelling is obviously less present in the *trans*-coupling, which is determined by the difference between the contribution of the metal  $p_\sigma$ -orbital and the metal  $s$ - and  $d_\sigma$ -orbitals. The number of compounds in our set is too small to allow a statistical treatment of the  ${}^2J(P_i-P_j)_{trans}$  data.

Dr. W.A. Schenk of the University of Würzburg has kindly provided us with the coupling constant data for the complexes *cis*- and *trans*- $LL'W(CO)_4$  ( $L, L' =$  tertiary phosphine, aminophosphine or phosphite) [28]. In these complexes the substituents on phosphorus were varied but not those on the other ligands on the metal. Nonetheless, for the 17 *trans*-complexes an expression for  ${}^2J(P_1-P_2)_{trans}$  similar to equation 1 gives a good fit with a correlation coefficient  $r$  of 0.990 and  $a_{trans}/\mu_W = 0.759$ ,  $b_{trans}/\mu_W = 0.067$  and  $c = 152$  Hz. Clearly, similar

rules govern the  $^1J(M-P)$  and  $^2J(P_1-P_2)$  relationship in *cis* and *trans* complexes.

For the 10 *cis* LL'W(CO)<sub>4</sub> complexes, the correlation according to equation 1 is poorer. These complexes are found in groups which are apparently determined by the number of alkoxy groups present on the phosphorus. The statistical constants are:  $a_{cis}/\mu_W = 0.039$ ,  $b_{cis}/\mu_W = 0.0057$  and  $c = -2.4$  Hz, with a correlation coefficient of 0.912. The  $a$  values of the tungsten complexes cannot be compared with the values of the rhodium and platinum complexes because different phosphines are involved.

## Acknowledgment

The authors wish to thank Dr. W.A. Schenk for providing the data for the tungsten compounds, Dr. F.W. Pijpers of the department of Analytical Chemistry for the statistical treatment of the NMR data, Mr. J. van Kessel for recording the  $^{31}P$  NMR spectra, and Mr. P.J.J. Koonen for performing C, H and N analyses.

## References

- 1 D.H.M.W. Thewissen and A.W. Gal, Proceedings of the Conference on Rhodium in Homogeneous Catalysis, Veszprém, 1978, pp. 6-13; Part IA: D.H.M.W. Thewissen and H.L.M. van Gaal, *J. Organometal. Chem.* 172 (1979) 69; Part IB: D.H.M.W. Thewissen, *J. Organometal. Chem.*, 188 (1980) 211.
- 2 A.W. Gal, J.W. Gosselink and F.A. Vollenbroek, *J. Organometal. Chem.*, 142 (1977) 357.
- 3 A.W. Gal and F.H.A. Bolder, *J. Organometal. Chem.*, 142 (1977) 375.
- 4 J.A. Osborn and G. Wilkinson, *Inorg. Synth.*, 10 (1967) 67.
- 5 A. van der Ent and A.L. Onderdelinden, *Inorg. Synth.*, 14 (1973) 94.
- 6 K. Issleib and G.H. Harzfeld, *Chem. Ber.*, 97 (1964) 3430; K. Issleib and G. Harzfeld, *Z. Anorg. Allg. Chem.*, 351 (1967) 18.
- 7 A. van der Gen, Belg. Patent, C.A. 65 (1966) 8960h.
- 8 D.H.M.W. Thewissen and H.P.M.M. Ambrosius Rec. Trav. Chim. Pays-Bas, in press.
- 9 I. Ojima, K. Akiba and N. Inamoto, *Bull. Chem. Soc. Japan*, 42 (1969) 2975.
- 10 D.H.M.W. Thewissen, *J. Organometal. Chem.*, 192 (1980) 115.
- 11 A. Cormier, K. Nakamoto, P. Crystophliempk and A. Muller, *Spectrochim. Acta A*, 31 (1974) 1059.
- 12 P. Lakshimi, R. Bhaskera and U. Agarwala, *Inorg. Chim. Acta*, 5 (1971) 354.
- 13 K.A. Jensen, B.M. Dahl, P.H. Nielsen and G. Borch, *Acta Chem. Scand.*, 26 (1972) 2241.
- 14 R. Mattes and G. Pauleickhoff, *Spectrochim. Acta A*, 30 (1974) 386.
- 15 S.N. Olafsson, Ph.D. Thesis Hamburg, 1973.
- 16 R. Lenck, Ph. D. Thesis, Hamburg, 1973.
- 17 E.G. Finer and R.K. Harris, *Progress in NMR Spectroscopy* Vol. 6, Pergamon Press, London, 1971, p61.
- 18 R.D. Bertrand, F.B. Ogilvie and J.G. Verkade, *J. Amer. Chem. Soc.*, 92 (1970) 1908.
- 19 F.B. Ogilvie, J.M. Jenkins and J.G. Verkade, *J. Amer. Chem. Soc.*, 92 (1970) 1916.
- 20 R. Meij, D.J. Stufkens and K. Vrieze, *J. Organometal. Chem.*, 144 (1978) 239.
- 21 R. Meij, Ph. D. Thesis, Amsterdam, 1978.
- 22 E. Lindner and B. Schilling, *Chem. Ber.*, 110 (1977) 3889.
- 23 E. Lindner and W.P. Meyer, *J. Organometal. Chem.*, 67 (1974) 277.
- 24 T.H. Brown and P.J. Green, *J. Amer. Chem. Soc.*, 92 (1970) 2359.
- 25 J.F. Nixon and A. Pidcock, *Ann. Rev. NMR Spectra*, 2 (1969) 346.
- 26 T.G. Appleton, H.C. Clark and L.E. Manzer, *Coord. Chem. Rev.*, 10 (1973) 335.
- 27 F.H. Allen, A. Pidcock and C.R. Waterhouse, *J. Chem. Soc. A*, (1970) 2087.
- 28 W.A. Schenk, *Habilitationsschrift, Universität Würzburg*, 1979; *Proceedings IXth I.C.O.M.C., Dijon*, 1979, P13F and D3.