

dergoing very rapid exchange in a process which does not permute CF_3 groups. This result is confirmed by the averaged A_3B_3 pattern [$\phi = -74.72$ and -75.51 , ($J = 9.2$ Hz)] observed in the ^{19}F NMR spectrum (CF_3 region) of an equimolar mixture of **2/4** (-25°C). The trifluoromethyl groups in **2** exchange at elevated temperatures, with an estimated activation energy $\Delta G^{\ddagger}_{54} = 17.0$ kcal/mol. Noticeable curvature appeared in the plot of $\ln k$ vs. $1/T$, and the accuracy of derived activation parameters is thereby reduced. We interpret the nonlinearity to indicate that at least two enantiomerization mechanisms are operating.¹⁵ In contrast to the behavior of **1**, spectra of **2** are solvent dependent, and the exchange rate is sensitive to impurities. Addition of small amounts of HMDS⁶ to propionitrile solutions of **2** reduced the CF_3 exchange rate and raised ΔG^{\ddagger}_{94} to ca. 19 kcal/mol.

The observation that the exchange rate of **1** is independent of solvent or added nucleophilic solvent militates against involvement of hexacoordinate silicon.¹⁷ While enantiomerization of **2** cannot be accounted for by simple silicon-oxygen bond breaking steps, rapid exchange of the fluoride ligand may play a role in the itinerary for CF_3 exchange.

Acknowledgment. We thank Dr. G. S. Reddy and W. John for ^{19}F NMR data. Professor J. D. Roberts kindly supplied a copy of DNMR 3 which was adapted for use on our computer by Dr. R. M. Hilmer.

Supplementary Material Available: Tables of atomic coordinates, thermal parameters, and structure amplitudes (47 pages). Ordering information is given on any current masthead page.

(15) Many alternate mechanisms are possible and some may act in concert with pseudorotation. We note two which appear most reasonable: (1) An HF-catalyzed axial-equatorial exchange of the fluorine ligand; (2) axial loss coupled with edge attack on **4** (or equatorial loss coupled with face attack) of the rapidly exchanging fluorine ligand.¹⁶

(16) The lowest energy pathway for fluoride exchange is believed to involve axial loss and face attack. For calculations on model systems, see Wilhite, D. L.; Spialter, L. *J. Am. Chem. Soc.* **1973**, *95*, 2100. Paybutt, P. *Mol. Phys.* **1975**, *29*, 389.

(17) Such species have been implicated in the racemization of tetracoordinate silicon halides (Corriu, R. J. P.; Henner, M. *J. Organomet. Chem.* **1974**, *74*, 1.) They were believed responsible for axial/equatorial site exchange in pentacoordinate silicon fluorides (Marat, R. K.; Janzen, A. F. *Can. J. Chem.* **1977**, *55*, 3845.)


Alkylation and Oxidative Dimerization of Enolate Anions by Radical Chain Processes¹

Glen A. Russell,* Boguslaw Mudryk, and Mikolaj Jawdosiuik

Department of Chemistry, Iowa State University
Ames, Iowa 50011

Received March 20, 1981

The radical chain involving reactions 1-3^{2,3} has been termed



$\text{S}_{\text{RN}}1$.⁴ In such substitutions the possibility exists that reactions 1 and 2 may merge and that R^\cdot may not be an intermediate. A distinction between a concerted process and the consecutive reactions 1 and 2 can be made on the basis of competitive experiments. When two anions are allowed to compete for a series of substrates containing a common R moiety, the relative reactivities


(1) Electron Transfer Processes. 27. This work was supported by Grant CHE-7823866 from the National Science Foundation.

(2) Russell, G. A.; Danen, W. C. *J. Am. Chem. Soc.* **1966**, *88*, 5663; **1961**, *90*, 347. Russell, G. A.; Norris, R. K.; Panek, E. J. *Ibid.* **1971**, *93*, 5839.

(3) Kornblum, N.; Michael, R. E.; Kerber, R. C. *J. Am. Chem. Soc.* **1966**, *88*, 5660, 5662. Kornblum, N.; Boyd, S. R.; Stuchal, F. W. *Ibid.* **1970**, *92*, 5783. Kornblum, N. K.; Boyd, S. D. *Ibid.* **1970**, *92*, 5784.



(4) Bunnett, J. F. *Acc. Chem. Res.* **1978**, *11*, 413.

Scheme I. Bimolecular Substitution and Oxidative Dimerization





of the anions should be independent of the leaving groups only if free R^\cdot is an intermediate. This technique has been employed in aromatic $\text{S}_{\text{RN}}1$ reactions,⁵ and for the reaction of XCMe_2NO_2 ($\text{X} = \text{Cl}, \text{NO}_2$, or $p\text{-MePhSO}_2$) with $\text{Me}_2\text{C}=\text{NO}_2^-$ and $\text{MeC}(\text{CO}_2\text{Et})_2^-$ in Me_2SO .⁶

We now report a second type of competition in the reaction of easily oxidized monoenolate anions ($\text{E} = \text{RC}(\text{O}^-)=\text{CHR}'$) with XCMe_2NO_2 , leading to coupling (**1** and **2**) and symmetrical dimerization (**3**) products. Our results require that both processes **4** and **5** proceed by free radical chains and that the competition




between these processes is determined by bimolecular reactions of $\text{XCMe}_2\text{NO}_2^\cdot$ and not by reactions of free $\text{O}_2\text{NCMe}_2^\cdot$.

The competition between (4) and (5) for $\text{E}^- = \text{PhC}(\text{O}^-)=\text{CHR}'$ with 2-chloro-2-nitropropane leads only to **1** and **2** with $\text{R}' = \text{H}$ and only to **3** with $\text{R}' = \text{Ph}$. With $\text{R}' = \text{Me}, \text{Et}$, or $i\text{-Pr}$ both products **1** and **3** are observed. Increasing the steric bulk of R' or the resonance stabilization of E^- favors process 5. In all cases the presence of 5-10 mol % of $(t\text{-Bu})_2\text{NO}^\cdot$ prevents the formation of **1-3**, and the starting phenone can be recovered. With $\text{R}' = \text{Me}$ or $i\text{-Pr}$ the ratio **1/3** is unaffected by the presence of $\text{Me}_2\text{C}=\text{NO}_2\text{Li}$ in solvent mixtures of THF-hexane- Me_2SO (or HMPA). For these systems E^- is trapped more readily by E^- than by $\text{Me}_2\text{C}=\text{NO}_2^-$. This is surprising since the addition step of reaction 6 must be more exothermic than the corresponding step of reaction 7. Apparently E^- preferentially reacts with the anion which is the stronger base.⁷



The ratio **1/3** from $\text{E}^- = \text{PhC}(\text{O}^-)=\text{CHR}'$ or $\text{Me}_3\text{CC}(\text{O}^-)=\text{CH}_2$ and XCMe_2NO_2 is independent of the concentrations of E^- or XCMe_2NO_2 but depends strongly on the nature of X with the ratio **1/3** decreasing from $\text{X} = \text{Cl}$ to $\text{X} = p\text{-MePhSO}_2$ or NO_2 (Table I). A similar effect is observed for cyclohexanone enolate anion where $\text{ClCMe}_2\text{NO}_2$ yields >80% of the C-alkylation products, but $\text{Me}_2\text{C}(\text{NO}_2)_2$ yields mainly **3**.^{8,9}

It is impossible to explain the variation in the ratio **1/3** from competition between reactions 8 and 9 or by competition between



(5) Galli, C.; Bunnett, J. F. *J. Am. Chem. Soc.* **1979**, *102*, 6137.

(6) Russell, G. A.; Ros, F.; Mudryk, B. *J. Am. Chem. Soc.* **1980**, *102*, 7601.

(7) Tolbert, L. M. *J. Am. Chem. Soc.* **1980**, *102*, 3531.

(8) Russell, G. A.; Jawdosiuik, M.; Ros, F. *J. Am. Chem. Soc.* **1979**, *101*, 3379.

(9) In THF-hexane-HMPA (53:34:13) at -35°C for 1 h, cyclohexanone enolate and $\text{ClCMe}_2\text{NO}_2$ yield 67% of **1**, 15% of **2**, and 11% of **3**.


Table I. Yields of RCOCH(R')CMe₂NO₂ (1), RCOC(R')=CMe₂ (2), [RCOCH(R')]₂ (3), and RCOCH(R')X (4) from the Reaction of RC(OLi)=CHR' and XCMe₂NO₂ in THF-Hexane (60:40)^a

R, R'	X	conditions ^b	yield, %				
			1 ^c	2 ^c	3 ^c	4 ^c	(1 + 2)/3
<i>t</i> -Bu, H	Cl ^d	3 h	0, 0 ^e	72, 6 ^e	<5, 2 ^e	0, 0 ^e	>14
<i>t</i> -Bu, H	NO ₂ ^d	3 h	0, 0 ^e , 0 ^f	20, 7, 25 ^f	30, 9, 27 ^f	15, 16, 13 ^f	0.7, 0.9 ^f
<i>t</i> -Bu, H	<i>p</i> -MePhSO ₂ ^d	3 h	0	28	38	0	1.4
<i>t</i> -Bu, H	<i>p</i> -MePhSO ₂ ^d	1.5 h, 13 vol % Me ₂ SO	0, 0 ^f	45, 29 ^f	10, 6 ^f	0	4.5
<i>t</i> -Bu, H	<i>p</i> -MePhSO ₂ ^d	1.5 h, 0 °C	25	10	48	0	0.8
Ph, H	Cl ^d	1 h, -20 °C, 13 vol % HMPA	0	97	0	0	∞
Ph, Me	Cl	1 h, 0-10 °C	70, 78, 66 ^h	0, 0, 0 ^h	23, 19, 22 ^h	0, 0, 0 ^h	3.0, 4.1, 3.0 ^h
Ph, Me	Cl	1 h	48, 0 ^e	0, 0 ^e	37, 0 ^e	0, 0 ^e	1.3
Ph, Me	Cl	1 h, 100% THF	50	0	28	0	1.8
Ph, Me	NO ₂	1 h	10	0	17	24	0.6
Ph, Me	<i>p</i> -MePhSO ₂	1 h	21, 26, 32 ^{g, h}	0, 0, 0 ^{g, h}	52, 47, 54 ^{g, h}	15, 5, 5 ^{g, h}	0.6, 0.4, 0.55 ^{g, h}
Ph, Me	Cl	1 h, 0 °C, 13 vol % HMPA	76, 52, 0 ^e	4, 11, 0 ^e	12, 14, 0 ^e	0, 0, 0 ^e	6.6, 4.0 ^f
Ph, Me	Cl	15 min, 0-10 °C, 13 vol % Me ₂ SO	58	7	11	0	5.0
Ph, Me	Cl	1 h, 35 °C, 13 vol % Me ₂ SO	28	21	13	0	3.8
Ph, Me	Cl	8 min, 30 °C, K ⁺ in Me ₂ SO	0	7	31	0	0.2
Ph, Me	Cl	1 h, 5 °C, HMPA	0	5	31	0	0.2
Ph, Et	Cl	1 h, -10-0 °C, 13 vol % HMPA	73	0	13	0	5.6
Ph, <i>i</i> -Pr	Cl	1 h	4, 8, 0 ^e	0, 0, 0 ^e	66, 69, 0 ^e	0, 0, 0 ^e	0.06, 0.1 ⁱ
Ph, <i>i</i> -Pr	Cl	1 h, 0-10 °C, 13 vol % HMPA	35, 29, 25 ^h	0, 0, 0 ^h	42, 42, 37 ^h	1.5, 4, 7 ^h	0.8, 0.7, 0.7 ^h
Ph, <i>i</i> -Pr	Cl	1 h, 0 °C, 13 vol % Me ₂ SO	24, 0 ^e	0, 0 ^e	32, 0 ^e	8.5, 29 ^e	0.8
Ph, <i>i</i> -Pr	Cl	1 h, 10 °C, 38 vol % Me ₂ SO	16, 0 ^e	0, 0 ^e	23, 0 ^e	20, 44 ^e	0.7
Ph, <i>i</i> -Pr	Cl	1 h, 35 °C, K ⁺ in Me ₂ SO	2, 0 ^e	0, 0 ^e	25, 0 ^e	18, 30 ^e	0.05
Ph, <i>i</i> -Pr	Cl	1 h, 5 °C, HMPA	0	0	0	44	-
Ph, Ph	Cl	3 h, 35 °C	<2, 0 ^e	0, 0 ^e	66, 0 ^e	0, 0 ^e	<0.02

^a Satisfactory elemental analysis, ¹H NMR spectra and high resolution MS were obtained for all new compounds. The known dimers 3 isolated as the *meso*, *dl* mixtures, or as the pure isomers were demonstrated to be symmetrical dimers by ¹H NMR and my comparison with literature data. ^b Five millimoles of (*i*-Pr)₂NH and 5 mL of THF were added to 5 mmol of *n*-BuLi in hexane (3.2 mL) at -40 °C. The solution was warmed to 0 °C, cooled to -20 °C, and the ketone added dropwise to give a solution ~0.5 M in E⁻. Additional reagents were added at 0 or 25 °C and the solutions irradiated with a 275-W sunlamp which in the absence of cooling maintained a reaction temperature of 35 °C. Unless otherwise indicated the molar ratio of E⁻ to XCMe₂NO₂ was 1:1. ^c Based on ¹H NMR. For reactions with E⁻/XCMe₂NO₂ = 2, yields are based on the theoretical formation of 1 mol of product per mol of XCMe₂NO₂. With E⁻/XCMe₂NO₂ ≤ 1, yields are based on the theoretical formation of 1 mol of 1, 2, or 4 and 0.5 mol of 3 per mol of E⁻. ^d Molar ratio E⁻/XCMe₂NO₂ = 2:1. ^e 0.05 M (*t*-Bu)₂NO₂, 25 °C, no irradiation. ^f 0.15 M Me₂C=NO₂Li. ^g Diluted 5-fold with THF. ^h Molar ratio E⁻/XCMe₂NO₂ = 1:2. ⁱ 0.5 M 12-crown-4-ether.

unimolecular and bimolecular reactions of XCMe₂NO₂⁻. However, the effect of X on the ratio 1/3 is easily explained if reactions (1) and (2) have merged and competition occurs as shown in Scheme I. Reaction of XCMe₂NO₂⁻ with E⁻ to form 1⁻ or E⁻ predicts that the ratio 1/3 should depend on the structure of X. However, the mechanism by which E⁻ replaces X at a tetrasubstituted carbon atom is puzzling. Perhaps the formation of 1 and 3 involves a common intermediate distinct from free O₂NCMe₂. Electron transfer from E⁻ to XCMe₂NO₂⁻ would almost certainly be dissociative and the cage intermediate, [E⁻ X⁻ Me₂C=NO₂], is a possibility. Escape of E⁻ from the cage would lead to 3 via reaction 7, while coupling of E⁻ and Me₂C=NO₂⁻ in the cage would lead to 1⁻. It appears that as the ease of one-electron donation from the anion increases [e.g., from MeC(CO₂Et)₂⁻ to PhC(O⁻)=CHR'], the chain reaction involved shifts from one involving discrete step, (1)-(3), to a process in which reactions (1) and (2) are no longer distinct.

The competition between reactions 4 and 5 depends upon counterion and solvent (see Table I). For PhC(OLi)=CH-*i*-Pr the ratio 1/3 passes through a maximum as Me₂SO or HMPA is added to the THF-hexane solvent. In 100% Me₂SO or HMPA the ratio 1/3 is quite low for secondary enolate anions, perhaps reflecting that free monoenolate anions react with XCMe₂NO₂⁻ or O₂NCMe₂ to yield mainly E⁻. As Me₂SO or HMPA is added to the THF-hexane solvent system, the E₂ elimination of HNO₂ from 1 becomes important, particularly at higher temperatures (Table I). With primary enolate anions it is difficult to prevent this reaction, and experiments with these anions were performed by using 2 equiv of E⁻ to maximize the yield of 2. In solvents containing Me₂SO or HMPA, chlorine atom or nitro group transfer occurs (reaction 10). Reaction 10 was observed in the



presence or absence of radical chain inhibitors, and from inhibited

experiments it is clear that 4 is not converted to 3 by an S_N2 displacement.¹⁰

(10) The source of PhCOCH(Me)SO₂PhMe-*p* mentioned in Table I is not clear, but it may involve the attack of E⁻ on *p*-MePhSO₂⁻.


Bis Heteroannulation. 3. Facile Syntheses of (±)-Ligularone and (±)-Petasalbine

Peter A. Jacobi* and Donald G. Walker

Hall-Atwater Laboratories, Wesleyan University
Middletown, Connecticut 06457

Received March 23, 1981

Ligularone (1) and petasalbine (2) are furanoeremophilanes isolated from the rhizomes of the *Ligularia* and *Petasites* genus of plants.¹ These same species produce at least 10 other com-



pounds having the basic skeleton of the furanoeremophilanes,^{1b}

(1) (a) Novotný, L.; Herout, V.; Sörn, F. *Tetrahedron Lett.* **1961**, 697. (b) Naya, K.; Nakagawa, M.; Hayashi, M.; Hayashi, M.; Tsuji, K.; Naito, M. *Ibid.* **1971**, 2961. (c) Ishii, H.; Tozoy, T.; Minato, H. *Tetrahedron* **1965**, 21, 2605.