

# Synthesis of the bicyclic core of tagetitoxin†

Julien R. H. Plet and Michael J. Porter\*

Received (in Cambridge, UK) 19th January 2006, Accepted 31st January 2006

First published as an Advance Article on the web 14th February 2006

DOI: 10.1039/b600819d

A synthesis of the 9-oxa-3-thiabicyclo[3.3.1]nonane ring system, which constitutes the core of the RNA polymerase inhibitor tagetitoxin, has been achieved through cyclisation of a thiol onto an electrophilic ketone.

Tagetitoxin is a phytotoxin which was isolated from the plant pathogenic bacterium *Pseudomonas syringae* pv. *tagetis* in 1981.<sup>1</sup> The compound induces chlorosis in the apex of infected plants, an effect which has been traced to inhibition of RNA polymerase in chloroplasts.<sup>2</sup> Furthermore, tagetitoxin inhibits bacterial RNA polymerase,<sup>2</sup> and is the only natural product known to inhibit eukaryotic RNA polymerase III in a specific manner.<sup>3</sup> Recently, a crystal structure of the RNA polymerase from *Thermus thermophilus* with tagetitoxin bound to the active site was published;<sup>4</sup> consideration of this structure led the authors to postulate that tagetitoxin acts by stabilisation of an inactive intermediate in the transcription process.

The structure which was assigned to the compound shortly after its isolation<sup>5</sup> was later rejected in favour of a bicyclic structure based on the 9-oxa-3-thiabicyclo[3.3.1]nonane ring system. Structure **1** was favoured, although the spectroscopic data did not rule out the closely related structure **2** (Fig. 1).<sup>6</sup>

To date, limited work has been published detailing synthetic approaches to tagetitoxin and its analogues,<sup>7</sup> and none has yet been successful in generating the bicyclic ring system. In this communication we describe a synthesis of this bicyclic core structure, starting from a carbohydrate precursor.

Our initial strategy focused on the use of a carbene-mediated ring expansion of 1,3-oxathiolanes.<sup>8</sup> D-Glucose was converted to the bicyclic monothioacetal **3**<sup>9</sup> through displacement of an anomeric bromide and a 6-tosylate with potassium *O*-ethylxanthate (Scheme 1).<sup>10</sup> Ring expansion was then attempted using ethyl diazo(triethylsilyl)acetate and catalytic rhodium(II) acetate;<sup>8</sup> this led not to the anticipated bridged bicycle, but instead to glycol **5**. This product may arise through sulfur ylid formation





Fig. 1 Proposed structures of tagetitoxin.

and heterolytic C–S bond cleavage to give zwitterion **4**; rather than the desired C–C bond formation, this intermediate is presumed to undergo a ring-flip to the more stable conformer followed by proton transfer to afford the observed product.

To circumvent this difficulty, a conformationally constrained substrate was designed whose derived zwitterion (analogous to **4**) would be incapable of ring-flipping. 3-Methyl-D-glucose was converted to bicycle **6**, then the acetate groups were cleaved and a bridging di-*tert*-butylsilylene protecting group installed. Treatment of the resulting tricycle **7** with ethyl diazo(triethylsilyl)acetate in the presence of rhodium(II) heptafluorobutyrate gave a low yield of primary alcohol **9** as the only isolable product. In this case, the sulfur ylid **8** forms as required but this, rather than undergoing C–S bond heterolysis and ring expansion, is trapped by adventitious water to give the bicyclic alcohol **9**.


Due to the failure of our initial approach to the tagetitoxin skeleton, a new strategy was adopted in which the 1,4-oxathiane ring of the natural product would be formed by cyclisation of a thiol onto an electron-deficient ketone, rather than through a carbene-mediated ring expansion.



**Scheme 1** Reagents and conditions: i. TsCl, pyridine; Ac<sub>2</sub>O; ii. HBr, AcOH; iii. KSCSOEt, DMF, 50 °C (46% **3** over 3 steps) or KSCSOEt, acetone, reflux (47% **6** over 3 steps); iv. Et<sub>3</sub>SiC(N<sub>2</sub>)CO<sub>2</sub>Et, Rh<sub>2</sub>(OAc)<sub>4</sub>, benzene, reflux, 34%; v. NH<sub>3</sub>, MeOH, H<sub>2</sub>O, 50%; vi. <sup>t</sup>Bu<sub>2</sub>SiCl<sub>2</sub>, Et<sub>3</sub>N, CH<sub>2</sub>Cl<sub>2</sub>, 86%; vii. Et<sub>3</sub>SiC(N<sub>2</sub>)CO<sub>2</sub>Et, Rh<sub>2</sub>(O<sub>2</sub>CC<sub>3</sub>F<sub>7</sub>)<sub>4</sub>, benzene, reflux, 21%.

Department of Chemistry, University College London, Christopher Ingold Building, 20 Gordon Street, London, UK WC1H 0AJ.  
E-mail: m.j.porter@ucl.ac.uk; Fax: +44 20 7679 7463;  
Tel: +44 20 7679 4710

† Electronic supplementary information (ESI) available: HMQC and HMBC spectra for **21**. See DOI: 10.1039/b600819d




**Scheme 2** Reagents and conditions: i. TBDPS-Cl, imidazole, DMF, 99%; ii. BnBr, NaH, DMF, 87%; iii. NBS, aq. acetone, 95%; iv. Dess–Martin periodinane, pyridine, CH<sub>2</sub>Cl<sub>2</sub>, 69%; v. TMSCH=CH, *n*-BuLi, CeCl<sub>3</sub>·7H<sub>2</sub>O, THF, −78 °C to rt, 96%; vi. Et<sub>3</sub>SiH, TMSOTf, CH<sub>2</sub>Cl<sub>2</sub>, 74%; vii. NaOH, MeOH, CH<sub>2</sub>Cl<sub>2</sub>, 100%; viii. NBS, AgNO<sub>3</sub>, acetone, 98%; ix. KMnO<sub>4</sub>, NaHCO<sub>3</sub>, MgSO<sub>4</sub>, aq. MeOH, 84%; x. HF·py, THF, −78 °C to rt, 77%.

Phenyl 1-thio-β-D-glucopyranoside (**10**) was converted to the fully protected analogue **11**<sup>11</sup> before NBS-promoted hydrolysis of the thioglycoside linkage (Scheme 2). Oxidation to the δ-lactone **12** was accomplished using Dess–Martin periodinane. Cerium-mediated addition of trimethylsilylacetylene followed by deoxygenation and desilylation then afforded terminal alkyne **14**. Bromination to give **15** was followed by oxidation with potassium permanganate in aqueous methanol<sup>12</sup> to yield α-ketoester **16**.

On cleavage of the silyl ether with TBAF, concomitant elimination of the 2-benzoyloxy group (glucose numbering) to form an enol ether was observed. However, when silyl ether **16** was treated with HF–pyridine, the sole product was tricyclic acetal **17**, in which not only the silyl ether but also the 3- and 4-benzyl ethers had been cleaved, and an acetal had formed between the ketone and the 3- and 6-OH groups.

Formation of the tricyclic acetal **17** precluded introduction of a sulfur atom at C6, but alteration of the order of steps allowed completion of the synthesis of the tagetitoxin skeleton; thus double desilylation of **13** could be achieved with TBAF to give primary alcohol **18** (Scheme 3). Activation as the mesylate was followed by displacement with potassium thioacetate, and bromination of the alkyne afforded **19**. Oxidation as for compound **15** gave the α-ketoester **20**, and removal of the *S*-acetyl protecting group using hydrazine hydrate in methanol led directly to the bicyclic hemithioacetal **21**.

The structure of **21** was confirmed by mass spectrometry and NMR;† in particular, an HMBC correlation was observed between the hemithioacetal carbon at 71.9 ppm and one of the CH<sub>2</sub>S protons at 1.57 ppm. Vicinal coupling constants of 9.3 and 9.6 Hz between the pairs of CHOBn protons indicated a boat conformation for the tetrahydropyran ring, as depicted in **21**. The hemithioacetal was obtained as a single stereoisomer, although the configuration of this centre was not determined.



**Scheme 3** Reagents and conditions: i. TBAF, THF, 99%; ii. MsCl, Et<sub>3</sub>N, DMAP, CH<sub>2</sub>Cl<sub>2</sub>, 95%; iii. KSac, DMF, 99%; iv. NBS, AgNO<sub>3</sub>, acetone, 98%; v. KMnO<sub>4</sub>, NaHCO<sub>3</sub>, MgSO<sub>4</sub>, aq. MeOH, 71%; vi. N<sub>2</sub>H<sub>4</sub>·H<sub>2</sub>O, MeOH, 88%.

In conclusion, we have carried out the first synthesis of the tagetitoxin skeleton, by unmasking of a thiol in the presence of an electrophilic α-ketoester, triggering spontaneous cyclisation to a hemithioacetal. Efforts towards synthesis of the fully functionalised tagetitoxin structure are under way and will be reported in due course.

We thank EPSRC and UCL for financial support of this work.

## Notes and references

† Data for **21**:  $[\alpha]_D^{25} +3.1$  (*c* 1.07 in EtOH);  $\nu_{\max}/\text{cm}^{-1}$  (film) 3445 (OH), 2928 (CH), 1736 (C=O);  $\delta_{\text{H}}$  (500 MHz, C<sub>6</sub>D<sub>6</sub>) 7.38–6.99 (15H, m, ArH), 5.01 (1H, d, *J* 11.3), 4.92 (1H, d, *J* 11.3), 4.82 (1H, d, *J* 11.6), 4.75 (1H, d, *J* 11.6) and 4.65 (1H, d, *J* 12.0, 5 of PhCH<sub>2</sub>), 4.43 (1H, dd, *J* 9.3, 2.8, *H*-2), 4.39 (1H, *J* 12.0, 1 of PhCH<sub>2</sub>), 4.36 (1H, br d, *J* 2.8, *H*-1), 4.22 (1H, t, *J* 9.4, *H*-3), 4.19 (1H, td, *J* 3.6, 1.9, *H*-5), 4.11 (1H, s, OH), 4.03 (1H, dd, *J* 9.6, 3.7, *H*-4), 3.30 (1H, dd, *J* 13.4, 3.6, 1 of CH<sub>2</sub>S), 3.24 (3H, s, CH<sub>3</sub>), 1.57 (1H, br d, *J* 13.4, 1 of CH<sub>2</sub>S);  $\delta_{\text{C}}$  (125 MHz, C<sub>6</sub>D<sub>6</sub>) 173.4 (C=O), 139.1, 138.9 and 138.6 (3 × aromatic C), 82.2 (C-3), 80.0 (C-4), 79.8 (C-2), 79.5 (C-1), 75.0 (PhCH<sub>2</sub>), 73.3 (C-5), 73.2 (PhCH<sub>2</sub>), 72.4 (PhCH<sub>2</sub>), 71.9 (SCOCH), 52.5 (OCH<sub>3</sub>), 40.9 (C-6), other aromatic carbons obscured by solvent; *m/z* (FAB+) 559 (MNa<sup>+</sup>, 2%), 326 (21), 199 (26), 176 (100); HRMS (FAB+) found 559.1784; C<sub>30</sub>H<sub>32</sub>O<sub>6</sub>SNa (MNa<sup>+</sup>) requires 559.1766.

- R. E. Mitchell and R. D. Durbin, *Physiol. Plant Pathol.*, 1981, **18**, 157–168.
- D. E. Mathews and R. D. Durbin, *J. Biol. Chem.*, 1990, **265**, 493–498.
- T. H. Steinberg, D. E. Mathews, R. D. Durbin and R. R. Burgess, *J. Biol. Chem.*, 1990, **265**, 499–505; for a recently identified specific inhibitor of RNA polymerase III, see L. P. Wu, J. Pan, V. Thoroddsen, D. R. Wyong, R. K. Blackman, C. E. Bulawa, A. E. Gould, T. D. Ocain, L. R. Dick, P. Errada, P. K. Dorris, T. Parkinson, T. Wood, D. Kornitzer, Z. Weissman, I. M. Willis and K. McGovern, *Eukaryotic Cell*, 2003, **2**, 256–264.
- D. G. Vassilyev, V. Svetlov, M. N. Vassilyeva, A. Perederina, N. Igarashi, N. Matsugaki, S. Wakatsuki and I. Artsimovitch, *Nat. Struct. Mol. Biol.*, 2005, **12**, 1086–1093.
- R. E. Mitchell and R. D. Durbin, *Phytochemistry*, 1983, **22**, 1425–1428.
- R. E. Mitchell, J. M. Coddington and H. Young, *Tetrahedron Lett.*, 1989, **30**, 501–504.
- (a) T. Sammakia, T. B. Hurley, D. M. Sammond, R. S. Smith, S. B. Sobolov and T. R. Oeschger, *Tetrahedron Lett.*, 1996, **37**, 4427–4430; (b) B. R. Dent, R. H. Furneaux, G. J. Gainsford and G. P. Lynch, *Tetrahedron*, 1999, **55**, 6977–6996.
- (a) M. Ioannou, M. J. Porter and F. Saez, *Chem. Commun.*, 2002, 346–347; (b) M. Ioannou, M. J. Porter and F. Saez, *Tetrahedron*, 2005, **61**, 43–50.

- 9 H. Driguez, C. McAuliffe, R. V. Stick, D. M. Tilbrook and S. J. Williams, *Aust. J. Chem.*, 1996, **49**, 343–348.
- 10 (a) L. S. Jeong, R. F. Schinazi, J. W. Beach, H. O. Kim, S. Nampalli, K. Shanmuganathan, A. J. Alves, A. Mcmillan, C. K. Chu and R. Mathis, *J. Med. Chem.*, 1993, **36**, 181–195; (b) L. S. Jeong, R. F. Schinazi, J. W. Beach, H. O. Kim, K. Shanmuganathan, S. Nampalli, M. W. Chun, W. K. Chung, B. G. Choi and C. K. Chu, *J. Med. Chem.*, 1993, **36**, 2627–2638.
- 11 M. Sasaki, K. Tachibana and H. Nakanishi, *Tetrahedron Lett.*, 1991, **32**, 6873–6876.
- 12 L. S. Li and Y. L. Wu, *Tetrahedron Lett.*, 2002, **43**, 2427–2430.

# Chemical Biology

An exciting news supplement providing a snapshot of the latest developments in chemical biology



Free online and in print issues of selected RSC journals!\*

**Research Highlights** – newsworthy articles and significant scientific advances

**Essential Elements** – latest developments from RSC publications

**Free links** to the full research paper from every online article during month of publication

\*A separately issued print subscription is also available

RSC Publishing

[www.rsc.org/chemicalbiology](http://www.rsc.org/chemicalbiology)