

A Stereocontrolled Synthesis of D-(+)-Showdomycin

Sung Ho Kang* and Sung Bae Lee

Department of Chemistry, Korea Advanced Institute of Science and Technology, Taejeon 305-701, Korea

Total synthesis of D-(+)-showdomycin **1** is accomplished starting from the enantiomerically pure *syn*-2,5-disubstituted dihydrofuran **2**.

In recent years C-ribosides have received considerable attention, because of their metabolic stability, in the development of antiviral, anticancer or anti-AIDS drugs.¹ Classified as tetrahydrofuran derivatives, they may be synthesized efficiently by the preferential formation of a tetrahydrofuranyl ring, followed by the necessary functionalization. This synthetic approach, though rarely employed, can offer advantages owing to the relatively constrained steric environment of the ring system. With this consideration, we have developed the stereoselective synthesis of 2,5-disubstituted tetrahydrofurans.² We herein describe an enantioselective total synthesis³ of an antitumour compound D-(+)-showdomycin **1**,⁴ starting from the enantiomerically pure *syn*-2,5-disubstituted dihydrofuran **2**, $[\alpha]_{\text{D}}^{20} -11.7$ (CHCl₃, *c* 0.83).^{2b}


Osmylation of dihydrofuran **2** with a catalytic amount of osmium tetroxide in the presence of *N*-methylmorpholine *N*-oxide (NMO) at -20°C produced a quantitative 10:1 mixture of the desired α -dihydroxylated product and the isomeric β -dihydroxylated compound (Scheme 1). The dihydroxy groups of the mixture were protected as acetonides, which were readily

separated to afford the required acetonide **3**, $[\alpha]_{\text{D}}^{20} -2.2$ (CHCl₃, *c* 0.82) in 82% yield. Acetonide **3** was oxidized using bromine⁵ in the presence of potassium carbonate in aqueous THF at 0°C , and the resulting lactols **4** were converted into the regio- and stereo-isomeric mixture of lactones **5** by the portionwise addition of chromium trioxide in Celite⁶ in ethereal dichloromethane at 0°C . Since the transformation of **4** or **5** into maleic anhydride was unsuccessful under various reaction conditions including Swern oxidation,⁷ Dess–Martin periodinane,⁸ Ley's tetrapropylammonium perruthenate⁹ and so forth, compounds **5** were subjected to reaction with hydroxylamine in methanolic pyridine[†] followed by esterification with diazomethane to give methyl esters **7** via oxime carboxylic acids **6** in 86% overall yield from **3**. The oxime hydroxy groups were mesylated and subsequently eliminated using mesyl chloride in the presence of diisopropylethylamine[†] at -30 to 0°C to furnish a 3:1 mixture of nitriles **8a** and **8b** in 88% combined yield.¹⁰ For the construction of the required maleimide ring in showdomycin **1** from **8a** and **8b**, we attempted to hydrolyse their nitrile groups to amides. However, the desired hydrolysis could not be realized in our hands in spite of employing various reaction conditions such as manganese dioxide in silica,¹¹ basic hydrogen peroxide,¹² *tert*-butyl alcoholic potassium hydroxide,¹³ etc. The major problem was the migration of the double bonds in **8a** and **8b** to the side of the tetrahydrofuranyl ring under basic reaction conditions. In order to avoid the use of basic media in the formation of maleimide, we decided to form the methoxymethyl esters instead of methyl esters. Accordingly, **6** was esterified using chloromethyl methyl ether (MOMCl) in the presence of diisopropylethylamine[†] at low temperature, and the resulting reaction mixture was treated *in situ* with mesyl chloride to provide a 3:1 mixture of nitriles **9a**, $[\alpha]_{\text{D}}^{20} -14.5$ (CHCl₃, *c* 0.75) and **9b**, $[\alpha]_{\text{D}}^{20} +2.2$ (CHCl₃, *c* 0.49) in 73% overall yield from **3**. Exposure of **9a** and **9b** to a mixture of trifluoroacetic acid and trifluoroacetic anhydride at 45°C induced the cyclization of the conjugate nitrile ester functionality to maleimide,^{3e,h} and the deprotection of the resulting products was completed by the addition of water to the same pot to produce the desired D-(+)-showdomycin **1**, mp. 150 – 151°C , $[\alpha]_{\text{D}}^{20} 50.6$ (H₂O, *c* 0.455) in 72% overall yield.

Total synthesis of D-(+)-showdomycin **1** has been accomplished in 43% overall yield starting from *syn*-2,5-disubstituted dihydrofuran **2**, in which the seven-step sequence of the conversion of the furanyl group into the maleimide moiety did not require any chromatographic purification.

This work was supported by the Ministry of Science and Technology, and the Organic Chemistry Research Center sponsored by the Korea Science and Engineering Foundation.

Received, 30th January 1995; Com. 5/00553A



Scheme 1 Reagents and conditions: i, OsO₄ (cat.), NMO, aq. acetone, -20°C ; ii, *p*-TsOH (cat.), acetone, room temp.; iii, Br₂ (2 equiv.), K₂CO₃ (0.5 equiv.), THF–H₂O (5:1), 0°C ; iv, CrO₃ on Celite (3 equiv.), CH₂Cl₂–Et₂O (3:1), 0°C ; v, HONH₂·HCl, pyridine–MeOH (1:1), room temp.; vi, CH₂N₂, Et₂O, 0°C ; vii, MeSO₂Cl, Pr₂NEt, CH₂Cl₂, -20 to 0°C ; viii, MOMCl, Pr₂NEt, CH₂Cl₂, -30 to -20°C , then MeSO₂Cl, -30 to 0°C ; ix, CF₃CO₂H, (CF₃CO)₂O, 45°C , then H₂O, room temp.

Footnote

[†] The use of triethylamine induced the isomerization of the double bond to the side of the tetrahydrofuranyl ring.

References

- V. Hacksell and G. D. Daves, *Prog. Med. Chem.*, 1985, **22**, 1; M. H. St Clair, T. Spector, K. J. Weinhold, W. H. Miller, A. J. Langlais and P. A. Furman, *Antimicrob. Agents Chemother.*, 1987, **31**, 1972.
- (a) S. H. Kang and S. B. Lee, *Tetrahedron Lett.*, 1993, **34**, 1955; (b) 1993, **34**, 7579.

- 3 For the previous syntheses of D-(+)-showdomycin **1**, see: (a) L. Kalvoda, J. Farkas and F. Sorm, *Tetrahedron Lett.*, 1970, **11**, 2297; (b) G. Trummelitz and J. G. Moffatt, *J. Org. Chem.*, 1973, **38**, 1841; (c) R. Noyori, T. Sato and Y. Hayakawa, *J. Am. Chem. Soc.*, 1978, **100**, 2561; (d) J. G. Buchanan, A. R. Edgar, M. J. Power and C. T. Shanks, *J. Chem. Soc., Perkin Trans. 1*, 1979, 225; (e) T. Inoue and I. Kuwajima, *J. Chem. Soc., Chem. Commun.*, 1980, 251; (f) G. Just, T. J. Liak, M.-I. Lim, P. Potvin and Y. S. Tsantrizos, *Can. J. Chem.*, 1980, **58**, 2024; (g) A. P. Kozikowski and A. Ames, *J. Am. Chem. Soc.*, 1981, **103**, 3923; (h) M. Ohno, Y. Ito, M. Arita, T. Shibata, K. Adachi and H. Sawai, *Tetrahedron*, 1984, **40**, 145; (i) P. D. Kane and J. Mann, *J. Chem. Soc., Perkin Trans. 1*, 1984, 657; (j) N. Katagiri, T. Haneda and N. Takahashi, *Heterocycles*, 1984, **22**, 2195; (k) T. Sato, Y. Hayakawa and R. Noyori, *Bull. Chem. Soc. Jpn.*, 1984, **57**, 2515; (l) A. G. M. Barrett, H. B. Broughton, S. V. Attwood and A. A. L. Gunatilaka, *J. Org. Chem.*, 1986, **51**, 495; (m) T. Kametani, K. Kawamura and T. Honda, *J. Am. Chem. Soc.*, 1987, **109**, 3010; (n) M. S. P. Gonzalez, R. M. D. Aciego and F. J. L. Herrera, *Tetrahedron*, 1988, **44**, 3715; (o) D. H. R. Barton and M. Ramesh, *J. Am. Chem. Soc.*, 1990, **112**, 891; (p) G. Kim, S. Kang and S. N. Kim, *Tetrahedron Lett.*, 1993, **34**, 7627.
- 4 H. Nishimura, M. Mayama, Y. Komatsu, H. Kato, N. Shimaoka and Y. Tanaka, *J. Antibiot. Ser. A*, 1964, **17**, 148; R. J. Suhadolnik, *Nucleoside Antibiotics*, Wiley-Interscience, New York, 1970, p. 354; G. D. Daves, Jr. and C. C. Cheng, *Prog. Med. Chem.*, 1976, **13**, 303.
- 5 J. Jurczak and S. Pikul, *Tetrahedron Lett.*, 1985, **26**, 3039.
- 6 S. J. Flatt, G. W. J. Fleet and B. J. Taylor, *Synthesis*, 1979, 815.
- 7 A. J. Mancuso and D. Swern, *Synthesis*, 1981, 165.
- 8 D. B. Dess and J. C. Martin, *J. Am. Chem. Soc.*, 1991, **113**, 7277.
- 9 W. P. Griffith and S. V. Ley, *Aldrichimica Acta*, 1990, **23**, 13.
- 10 For the related transformation, see: J. S. Buck and W. S. Ide, *Organic Syntheses*, 1943, Coll. vol. 2, p. 622; C. O. Yang, H. Y. Wu, E. B. Fraser-Smith and K. A. M. Walker, *Tetrahedron Lett.*, 1992, **33**, 37; N. Minakawa and A. Matsuda, *Tetrahedron Lett.*, 1993, **34**, 661.
- 11 K.-T. Liu, M.-H. Shi, H.-W. Huang and C.-J. Hu, *Synthesis*, 1988, 715; P. Breuilles, R. Leclerc and D. Uguen, *Tetrahedron Lett.*, 1994, **35**, 1401.
- 12 L. F. Fieser and M. Fieser, *Reagents for Organic Synthesis*, Wiley, New York, 1967, vol. 1, p. 469.
- 13 J. H. Hall and M. Gisler, *J. Org. Chem.*, 1976, **41**, 3769.