This article was downloaded by: [Universitaetsbibliothek Giessen] On: 15 December 2014, At: 01:32 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

http://www.tandfonline.com/loi/gpss20

Synthesis and ¹⁷O Nmr

Spectroscopy of a Series of ¹⁷O Labeled Triarylphosphine OXIDES

Martin Pomerantz, Michael S. Terrazas, Yang Cheng & Xiaomin Gu

^a Department of Chemistry and Biochemistry, Box 19065, The University of Texas at Arlington, Arlington, Texas 76019-0065, USA

^b Department of Chemistry and Biochemistry, Box 19065, , The University of Texas at Arlington , Arlington, Texas 76019-0065, USA

 $^{\rm c}$ Department of Chemistry and Biochemistry, Box 19065 , The University of Texas at Arlington , Arlington , Texas 76019-0065, USA

^d Department of Chemistry and Biochemistry, Box 19065, The University of Texas at Arlington, Arlington, Texas 76019-0065, USA Published online: 06 Aug 2008.

To cite this article: Martin Pomerantz , Michael S. Terrazas , Yang Cheng & Xiaomin Gu (1996) Synthesis and ¹⁷O Nmr Spectroscopy of a Series of ¹⁷O Labeled Triarylphosphine OXIDES, Phosphorus, Sulfur, and Silicon and the Related Elements, 109:1-4, 505-508, DOI: <u>10.1080/10426509608545201</u>

To link to this article: http://dx.doi.org/10.1080/10426509608545201

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor

& Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sublicensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <u>http://</u> www.tandfonline.com/page/terms-and-conditions

SYNTHESIS AND ¹⁷O NMR SPECTROSCOPY OF A SERIES OF ¹⁷O LABELED TRIARYLPHOSPHINE OXIDES

MARTIN POMERANTZ,* MICHAEL S. TERRAZAS, YANG CHENG, AND XIAOMIN GU

Department of Chemistry and Biochemistry, Box 19065, The University of Texas at Arlington, Arlington, Texas 76019-0065, USA

Abstract. A series of nine ¹⁷O labeled triarylphosphine oxides $[(p-R-C_6H_4)_3PO]$ was synthesized, ¹⁷O NMR spectroscopic studies were carried out (toluene solvent / 95 °C and CDCl₃ / 60 °C) and the spectrum was fit with two Lorentzian peaks. The chemical shifts range from δ 51.8 to 55.7 in toluene and δ 44.8 to 48.9 in CDCl₃, while ¹J_{PO} varies from 159.6 to 168.6 Hz in toluene. The data were fit to the Taft DSP and Hammett equations and related to other NMR parameters for this system and the analogous λ^5 -phosphazenes [($p-R-C_6H_4$)₃PNPh]. Using the Taft DSP equation the ¹⁷O substituent chemical shifts gave ρ_I and ρ_R with opposite signs which is different from what is observed with the λ^5 -phosphazenes. ¹J_{PO}, on the other hand correlates best with the Hammett σ_P^+ constants. The data are consistent with a triple bond contribution to the PO bonding.

Key Words: Triarylphosphine oxides, O-17 NMR spectroscopy

INTRODUCTION

For a number of years we have been interested in the synthesis and properties of phosphorus and nitrogen compounds, in particular λ^5 -phosphazenes (R₃P=N-R'; phosphinimines) and related molecules. We have carried out considerable ³¹P, ¹⁵N and ¹³C NMR spectroscopic studies of several series of λ^5 -phosphazenes, 1-4 [1-8], phosphines 5 and phosphine oxides 6 [7,8]. Using the NMR data along with PRDDO



molecular orbital calculations, we described the electronic structure in terms of charge densities and induced dipoles and suggested that $p\pi$ - σ * bonding between phosphorus and nitrogen [using a σ * P-C(phenyl) orbital] was an important factor in the PN bonding. Systems 4 and 6 were similar to each other [7,8] in that changes in R resulted in essentially the same changes in the chemical shifts of the substituted ring carbon, but the phosphorus

506 SYNTHESIS AND ¹⁷O NMR OF TRIARYLPHOSPHINE OXIDES

chemical shift in 6 was twice as sensitive as that in 4 to changes in R [7,8]. This, we believe, is the result of the *N*-phenyl ring in 4 being able to delocalize charge. We have now turned to the synthesis of ¹⁷O labeled phosphine oxides 6, for ¹⁷O NMR spectroscopic studies, in order to attempt to better understand the bonding in the oxides and how series 6 relates to series 4. Since the parent 6, Ar = C_6H_5 , had been prepared previously with an ¹⁷O label and its ¹⁷O NMR parameters reported [9-11], we did not anticipate problems with the synthesis or NMR spectroscopy.

RESULTS AND DISCUSSION

We have synthesized the series of triarylphosphine oxides 6^{-17} O with an approximately 10% ¹⁷O label and have obtained the ¹⁷O NMR spectra, which provided the chemical shifts and ¹⁷O₋³¹P coupling constants. The synthesis is presented in Scheme 1. ¹⁷O NMR

Scheme 1



spectra are generally quite broad due to the rapid quadrupolar relaxation [12] and we have found that rather elevated temperatures (95 °C / toluene solvent) and deconvolution using two overlapping Lorentzian bands were required to resolve the spectra into two peaks. However, with CDCl₃ solvent at 60 °C several of the compounds showed a single broad peak and resolution into two peaks was not very accurate. It is well known that higher temperatures and lower viscosity solvents help in sharpening up the peaks [12]. Table 1 gives the ¹⁷O NMR spectral data for series **6**-¹⁷O.

Analysis of the data for 6-¹⁷O (toluene / 95 °C) provided the following relationships: ¹⁷O SCS = 2.93 $\sigma_I - 5.37 \sigma_R^o$ (f = 0.160) and ${}^{1}J_{17O-31P} = 3.71 \sigma_P^+ + 165.9$ (r = 0.976) [13]. Perhaps the most interesting aspect of these data is that the Taft DSP treatment of the ¹⁷O SCS data gives opposite signs for ρ_I and ρ_R^o which is in marked contrast to the situation with 4 where ρ_I and ρ_R are both negative [7]. Thus the resonance effect in both systems appear similar (similar negative ρ values, -5.37 and -4.65 respectively) while the inductive effect in the two systems is opposite (ρ values of +2.93 and -1.99 respectively). This difference is consistent with a triple bond contribution to the PO bond where electron donation / withdrawal to the oxygen can occur via both inductive and resonance effects through two different $p\pi$ - σ^* / $p\pi$ -d π orbitals [14]. That is, a substituent such as dimethylamino might donate electrons by a resonance effect onto the oxygen through a π bond ($p\pi$ - σ^* or $p\pi$ -d π) while at the same time electron density can be withdrawn from the oxygen (which now has an excess of electron density because of this resonance effect) through the perpendicular π -bond and into the sigma system of the phenyl ring(s). It should be mentioned that the ¹⁷O-³¹P coupling constant correlation with a positive slope

	_	Chemical Shift (δ)	¹⁷ O ^{_31} P Coupling Constant (Hz)	Chemical Shift (δ)
Compound	R =	Toluene	Toluene	CDCl ₃
6a- ¹⁷ O	N(CH ₃) ₂	54.65	159.6	48.9
6b- ¹⁷ O	OCH ₃	54.97	163.1	48.1
6c- ¹⁷ O	CH ₃	52.77	163.6	45.4
6d- ¹⁷ O	Н	51.90	166.6	44.8
6e- ¹⁷ O	F	55.73	165.9	48.7
6f- ¹⁷ O	Cl	53.89	167.1	46.8
6g- ¹⁷ O	COOCH ₃	51.84	167.6	46.6
6h- ¹⁷ O	CF ₃	52.52	167.1	45.4
6i- ¹⁷ O	CN	52.86	168.6	46.0

TABLE 1. ¹⁷O NMR chemical shifts and coupling constants for series 6-¹⁷O.

also stands in contrast to the negative slope of the corresponding ${}^{15}N_{-31}P$ coupling constant in 4 (${}^{1}J_{15}N_{-31}P = -2.15 \sigma_{p} + 33.4 (r = 0.927)$ [7]. This again is consistent with a different type of PO and PN bonding. The chemical shifts in toluene span the range of δ 51.8 to 55.7 and these are deshielded relative to those in CDCl₃ which are δ 44.8 to 48.9. As expected, the chemical shifts are to higher field in chloroform [11], in agreement with the observations that more polar or hydrogen bonding solvents shield the oxygen [11], presumably due to the stabilization of greater negative charge on the oxygen. The coupling constants from the CDCl₃ experiments at the lower temperature, where they could be measured, however, were much less accurate than those from the toluene experiments. The peaks were broader and so the resolution of the doublets was much poorer and with several compounds single, broad peaks were observed.

There are two reports of the ¹⁷O NMR spectral parameters of triphenylphosphine oxide, **6d**, in the literature. In the first, CDCl₃ solvent, 30 °C, δ is given as 43.3 and ¹J_{PO} as 160 ± 2.4 Hz [10] and in the other, CD₃CN solvent, 70 °C, δ is given as 47.7 and ¹J_{PO} as 153.9 Hz [9]. While the chemical shifts seem to agree reasonably well with those obtained here, the coupling constants do not. A reasonable explanation for the differences, particularly in the latter study at 70 °C [9], is that with two broad overlapping peaks, peak fitting is required. This can be seen in Figure 1 (6b-¹⁷O, R = OCH₃, toluene solvent, 95 °C) where the doublet is only partially resolved and the fitted peaks show a greater separation (larger coupling constant) than the peak maxima from the original spectrum.

Acknowledgment. We thank the Robert A. Welch Foundation (Grant Y-0684) and the National Science Foundation (AMP Program) for financial support.





REFERENCES AND NOTES

- 1. M. POMERANTZ, B. T. ZIEMNICKA, Z. M. MERCHANT, W.-N. CHOU, W. B. PERKINS, and S. BITTNER, J. Org. Chem., 50, 1757 (1985).
- M. POMERANTZ, D. S. MARYNIČK, K. RAJESHWAR, W.-N. CHOU, L. THROCKMORTON, E. W. TSAI, P. C. Y. CHEN and T. CAIN, J. Org. Chem., 51, 1223 (1986).
- 3. M. POMERANTZ, W.-N. CHOU, M. K. WITCZAK, and C. G. SMITH, J. Org. Chem., 52, 159 (1987).
- 4. M. POMERANTZ, W.-N. CHOU, M. K. WITCZAK, and C. G. SMITH, *Phosphorus and Sulfur*, **30**, 799 (1987).
- 5. W.-N. CHOU, M. POMERANTZ, and M. K. WITCZAK, J. Org. Chem., 55, 716 (1990).
- 6. M. POMERANTZ and W.-N. CHOU, *Phosphorus, Sulfur and Silicon*, 51/52, 241 (1990).
- 7. W.-N. CHOU and M. POMERANTZ, J. Org. Chem., 56, 2762 (1991), .
- 8. M. POMERANTZ, in *Phosphorus-31 NMR Spectral Properties in Compound Characterization and Structural Analysis*, edited by L. D. Quin and J. G. Verkade (VCH Publishers, New York, 1994), Chap. 10, pp 115-128.
- 9. L. D. QUIN, J. SZEWCZYK, K. LINEHAN and D. L. HARRIS, Magn. Reson. Chem., 25, 271 (1987).
- R. D. SAMMONS, P. A. FREY, K. BRUZIK, and M.-D. TSAI, J. Am. Chem. Soc., 105, 5455 (1983).
- S. A. EVANS JR., in ¹⁷O NMR Spectroscopy in Organic Chemistry, edited by D. W. Boykin (CRC Press, Boca Raton, FL, 1991), Chapter 10.
- L. G. BUTLER, in ¹⁷O NMR Spectroscopy in Organic Chemistry, edited by D. W. Boykin (CRC Press, Boca Raton, FL, 1991), Chapter 1.
- 13. SCS = Substituent Chemical Shift; f = SD / rms.
- 14. D. G. GILHEANY, Chem. Rev., 94, 1339 (1994), and references cited therein.