

Reaction Mechanism

Insertion of CS₂ into Iridium–Fluorine Bonds**


Paul Kläring and Thomas Braun*

Because of the major importance of fluorinated organic molecules in material sciences,^[1] drug development,^[2] agrochemistry,^[1b,3] and as [¹⁸F] PET tracers,^[4] it is desirable to expand the chemist's toolbox for fluorination methods and to understand their reaction mechanisms. Transition-metal-mediated fluorination methods can be considered as a smart alternative when compared to conventional fluorination methods.^[5] Catalytic procedures often allow a nucleophilic or electrophilic fluorination under mild reaction conditions and considerable progress has been made in the last decade.^[6,7] A fluorination step to form a C–F bond generally proceeds either by a transition-metal fluoro complex as an intermediate or by fluorination of a ligand. For the former pathway reductive elimination can lead to C–F bonds.^[8]

Insertion reactions into metal–fluorine bonds would also be an interesting approach for transition-metal-mediated C–F bond-formation reactions, but so far they have hardly been identified. The formation of (β-fluorovinyl)gold complexes by reaction of gold(I) fluorides with alkynes has been reported but the conversions proceed, most likely, by an intermolecular fluorination.^[9] To the best of our knowledge, the formation of the fluorodithiocarbonato complex [Pt(κ²-(S,S)-S₂CF)(PPh₃)₂][HF₂] from a reaction of [Pt(F)(PPh₃)₂][HF₂] with CS₂ represents a unique example for which the formation of a C–F bond by an intermolecular insertion of a substrate into a metal–fluorine bond was suggested.^[10,11] However, the compound was characterized by infrared spectroscopy and single-crystal structure analysis, but the identity of the bifluoride anion [F⋯F 2.84(2) Å] is ambiguous.^[12]

Herein we report on the synthesis of iridium(III) fluorodithiocarbonato complexes by insertion of CS₂ into the Ir–F bond of the iridium(III) fluorides *trans*-[Ir(Ar^F)(F)(H)(PiPr₃)₂] (**1a**: Ar^F = 4-C₅NF₄; **1b**: Ar^F = 2-C₆H₃F₂; Scheme 1).^[13] The insertion mechanism is supported by density-functional theory (DFT) calculations.

Treatment of a solution of **1a** in THF with CS₂ led to the formation of *trans*-[Ir(4-C₅NF₄)(H)(κ²-(S,S)-S₂CF)(PiPr₃)₂] (**2a**) at room temperature (Scheme 1). The ¹³C-labeled isotopologue *trans*-[Ir(4-C₅NF₄)(H)(κ²-(S,S)-S₂¹³CF)(PiPr₃)₂]



Scheme 1. Syntheses of **2a** and **2b**. THF = tetrahydrofuran.

(¹³C)-**2a**) can be obtained by a comparable procedure using ¹³CS₂. Analogously, *trans*-[Ir(2-C₆H₃F₂)(H)(κ²-(S,S)-S₂CF)(PiPr₃)₂] (**2b**) is formed by a reaction of **1b** with CS₂ in THF at room temperature.

The ³¹P{¹H} NMR spectrum of **2a** displays a doublet at δ = 9.7 ppm (²J_{FP} = 5.8 Hz) for the phosphorus atoms in a mutually *trans* arrangement. The ¹⁹F{¹H} NMR spectrum shows a broad singlet at remarkably low field at δ = 96.8 ppm and four multiplets (δ = -98.4, -99.8, -108.8, -117.2 ppm) in a 1:1:1:1 ratio. Whereas the four multiplets indicate a restricted rotation of the tetrafluoropyridyl ligand about the metal–carbon bond,^[13,14] the singlet at δ = 96.8 ppm can be assigned to the fluorine atom of the fluorodithiocarbonato ligand. For the isotopologue [¹³C]-**2a** an additional coupling constant of ¹J_{C,F} = 395 Hz is observed. The IR (ATR) spectrum of **2a** displays an absorption band at 2218 cm⁻¹ for the IrH moiety.^[13,14b,15] The complex **2b** shows comparable spectroscopic data to that of **2a**.

DFT calculations were performed to model the reaction of **1a** with CS₂ to give **2a**.^[16] The formation of **2a** is predicted to be exothermic by an energy difference of -12.0 kJ mol⁻¹. In accordance with the NMR data the computed molecular structure of **2a** exhibits a distorted octahedral arrangement as it is depicted in Figure 1. The S₂CF moiety coordinates through its two sulfur atoms, which are located in the *trans* positions to the metal-bound hydrogen atom and to the tetrafluoropyridyl ligand. Note that the fluorodithiocarbonato ligand features a surprisingly short C–F bond length of 1.320 Å. For the above-mentioned cation [Pt(κ²-(S,S)-S₂CF)(PPh₃)₂]⁺ a C–F bond length of 1.27(3) Å was determined by X-ray crystallography.^[11] Similar short C–F bonds have been found for 2-fluoroamidinium, 2-fluoroimidazolium, and 2-fluoroimidazolium cations.^[17]

Two possible reaction pathways for the formation of **2a** have been considered (Scheme 2). Pathway A features

[*] P. Kläring, Prof. Dr. T. Braun
Humboldt-Universität zu Berlin, Department of Chemistry
Brook-Taylor-Strasse 2, 12489 Berlin (Germany)
E-mail: thomas.braun@chemie.hu-berlin.de

[**] We gratefully acknowledge support from the research training group "Fluorine as a Key Element" funded by the Deutsche Forschungsgemeinschaft. We would like to thank L. Zámotná and T. Ahrens for experimental assistance and Dr. C. Müller (Free University Berlin) for helpful discussions concerning the DFT calculations.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201305106>.




Figure 1. Computed structure of **2a**. The hydrogen atoms of the isopropyl groups are omitted for clarity. Selected distances [Å] and angles [°]: Ir1–P1 2.466, Ir1–P2 2.401, Ir1–C1 2.062, Ir1–H1 1.575, Ir1–S1 2.499, Ir1–S2 2.605, S1–C24 1.677, S2–C24 1.667, C24–F5 1.320; P1–Ir1–P2 170.9, C1–Ir1–H1 86.7, C1–Ir1–S1 172.0, C1–Ir1–S2 103.2, S1–C24–S2 120.5, S1–C24–F5 119.5, S1–C24–S2 120.5.

a reductive elimination of HF and subsequent coordination of CS₂ at the metal center and fluorination of the CS₂ ligand with HF. The second pathway B involves a concerted mechanism which is characterized by an insertion of CS₂ into the Ir–F bond. The formation of the reactive 14-electron iridium(I) intermediate *trans*-{Ir(4-C₅NF₄)(PiPr₃)₂} (**A2**; pathway A) via a three-membered transition state was found to have an activation energy of +129.1 kJ mol⁻¹. The structure of transition-state **TS1** is depicted in Figure 2 and is characterized by a simultaneous elongation of the Ir–F bond (2.337 Å) and the Ir–H bond (1.650 Å). The H··F separation is significantly shorter (1.240 Å) compared to that in **1a** (2.929 Å). For an HF molecule solid-state X-ray and gas-phase neutron diffraction experiments revealed H–F separa-





Figure 2. Computed structure for the transition state (**TS1**) of the reductive elimination of HF from **1a**. The hydrogen atoms of the isopropyl groups are omitted for clarity. Selected distances [Å] and angles [°]: Ir1–C1 1.980, Ir1–H1 1.650, Ir1–F5 2.337, F5–H1 1.240; P1–Ir1–P2 167.2, C1–Ir1–H1 129.5, C1–Ir1–F5 159.9.

tions in a range from 0.92 Å to 0.97 Å, which is in accordance with a computed distance of 0.918 Å.^[18] The complex *trans*-{Ir(4-C₅NF₄)(FH)(PiPr₃)₂} (**A1**) is formed as a shallow intermediate, which then releases HF to yield *trans*-{Ir(4-C₅NF₄)(PiPr₃)₂} (**A2**). The subsequent coordination of CS₂ to the 14-electron intermediate **A2** results in the formation of *trans*-[Ir(4-C₅HF₄)(κ²-(C,S)-SCS)(PiPr₃)₂] (**3**) at –2.5 kJ mol⁻¹. Note that the formal reductive elimination of HF from *trans*-[Ir(4-C₅NF₄)(F)(H)(CO)(PiPr₃)₂] (**4**) was observed before.^[13]

The computed reaction profile for pathway B, which involves the concerted insertion of CS₂ into the metal–fluorine bond, proceeds via a four-membered transition state (**TS2**) at +71.1 kJ mol⁻¹ (Figure 3). The Ir–F separation of 2.197 Å in **TS2** shows an incipient Ir–F bond cleavage and a significant elongation when compared to the Ir–F distance

in **1a** (crystal structure: 2.039(2) Å, computed: 2.076 Å). The simultaneous formation of the C–F bond and Ir–S bond is demonstrated by a C··F distance of 1.739 Å and an Ir··S distance of 2.609 Å. Comparable four-membered transition states were also discussed for the reverse reaction—a concerted C–F bond cleavage either by β-fluoride elimination or a metathesis-like mechanism.^[19] Note that an interaction of CS₂ with the Ir center requires a vacant coordination site at the metal. The LUMO of **1a** has σ-antibonding character in the Ir–H bond, but is essentially metal-centered with one lobe



Scheme 2. Energy profile for the reaction of **1a** with CS₂. Calculated energies are given in kJ mol⁻¹ and include a zero-point correction.





Figure 3. Computed structure for the transition state (**TS2**) of the insertion of CS_2 into the Ir–F bond of **1a**. The hydrogen atoms of the isopropyl groups are omitted for clarity. Selected distances [Å] and angles [°]: Ir1–C1 2.016, Ir1–H1 1.556, Ir1–S1 2.609, Ir1–F5 2.197, S1–C24 1.634, S2–C24 1.586, C24–F5 1.739; P1–Ir1–P2 165.5, C1–Ir1–H1 84.2, C1–Ir1–S1 108.6, C1–Ir1–F5 173.6, S1–C24–S2 148.8, S1–C24–F5 101.5, S1–C24–S2 109.7.

pointing towards the vacant *trans* position of the metal-bound hydrogen atom of **1a** (see the Supporting Information). As the reaction proceeds, **TS2** links to a shallow intermediate (**B1**) at +61.8 kJ mol^{−1}, and features a monodentate fluoro-dithiocarbonato ligand. The C–F separation of 1.388 Å in **B1** reveals the C–F bond formation, whereas an Ir···F distance of 2.620 Å denotes only a weak interaction between the metal and the fluorine atom.^[20] Subsequently, a rotation about the C–S bond and coordination of the second sulfur atom to Ir results in the formation of **2a**. Overall, the reaction pathway B is computed to be kinetically favored over the alternative route including HF elimination and CS_2 coordination (pathway A). In addition, a mechanism which involves a metallophosphorane intermediate generated by initial migration of the fluorine atom to one of the phosphorous atoms was computed to be unfavorable by an energy of +272.6 kJ mol^{−1}.^[8b,d,21]

Note that it was reported before that the reaction of **1a** with CO_2 results in the formation of the hydrogencarbonato complex *trans*-[Ir(4- C_5NF_4)(H)(κ^2 -(*O,O*)- O_2COH)($\text{P}i\text{Pr}_3$)₂] (**7**) in the presence of traces of water.^[13] An insertion of CO_2 into the Ir–F bond of **1a** was not observed. We also computed the formation of the putative insertion product *trans*-[Ir(4- C_5NF_4)(H)(κ^2 -(*O,O*)- O_2COF)($\text{P}i\text{Pr}_3$)₂] by DFT calculations and found it to be endothermic by +32.9 kJ mol^{−1}. Furthermore, several attempts to optimize a transition state comparable to **TS2** only converged to the structures of **1a** and CO_2 .

Experimental studies gave further support for the improbability of a fluorination of coordinated CS_2 . The complex *trans*-[Ir(4- C_5NF_4)(η^2 - CS_2)($\text{P}i\text{Pr}_3$)₂] (**3**) was prepared on treatment of *trans*-[Ir(4- C_5NF_4)(η^2 - C_2H_4)($\text{P}i\text{Pr}_3$)₂] (**5**) with CS_2 (Scheme 3). The ³¹P{¹H} NMR spectrum of **3** shows a singlet at δ = 12.1 ppm for the phosphine ligands in a mutually *trans* position. The ¹⁹F NMR spectrum of **3** displays four multiplets at δ = −98.4, −112.5, −118.0, and −117.2 ppm in a 1:1:1:1 ratio for the fluorine atoms of the tetrafluoropyridyl ligand. The molecular structure of **3** was also confirmed by X-ray diffraction analysis at −173 °C (Figure 4).^[22] The complex **3**



Scheme 3. Synthesis of *trans*-[Ir(4- C_5NF_4)(κ^2 -(*C,S*)-SCS)($\text{P}i\text{Pr}_3$)₂] (**3**).





Figure 4. An ORTEP diagram of **3**. Ellipsoids are drawn at the 50% probability level. hydrogen atoms are omitted for clarity. Selected bond length [Å] and angles [°] with standard deviations in parentheses and computed values in italics: Ir1–P1 2.391(3) 2.449, Ir1–P2 2.375(5) 2.446, Ir1–C1 2.038(9) 2.052, Ir1–C24 1.987(9) 1.973, Ir1–S1 2.459(2) 2.477, S1–C24 1.697(12) 1.685, S2–C24 1.602(11) 1.620; P1–Ir1–P2 166.58(9) 169.6, C1–Ir1–C24 124.4(4) 129.4, C1–Ir1–S1 169.3(3) 171.6, Ir1–C24–S1 83.3(4) 84.8, Ir1–C24–S2 136.2(7) 134.3, S1–C24–S2 140.4(7) 140.8.

features an almost square-planar arrangement of the phosphine ligands (P1–Ir1–P2 166.58(9) Å), the apical carbon atom of the tetrafluoropyridyl ligand and one sulfur atom of the CS_2 ligand [C1–Ir1–S1 169.3(3)°]. The carbon atom of the CS_2 ligand is bound to the metal with an C1–Ir1–C24 angle of 124.4(4)°. The C24–S1 bond [1.697(12) Å] is found to be longer than the C24–S2 bond [1.602(11) Å], and the CS_2 moiety is bent [S1–C24–S2 140.4(7)°] in a similar fashion as reported for other compounds bearing a η^2 - CS_2 ligand at rhodium(I), nickel(0) or palladium(0).^[23]

In accordance with the computational studies the iridium(I) complex **3** does not react with $\text{Et}_3\text{N}\cdot 3\text{HF}$ (THF, 12 h, 50 °C, Scheme 3) or $\text{py}\cdot\text{HF}$ (C_6D_6 , 12 h, 50 °C). Additionally, the presence of CsF or $\text{Et}_3\text{N}\cdot 3\text{HF}$ does not have any influence on the reaction rate for the conversion of **1a** with CS_2 . However, the insertion of CS_2 into the Ir–F bond of **1a** is reversible at 50 °C. The complex **1a** was observed by NMR spectroscopy. The complex **3** was not detected, but several other compounds were, and they could not be identified further. To get more insight into that reaction, **2a** was treated with Ph_3SnCl at 50 °C. This reaction led to the generation of the chlorido complex *trans*-[Ir(4- C_5NF_4)(Cl)(H)($\text{P}i\text{Pr}_3$)₂] (**6**)^[13,24] and a colorless solid, which was identified as Ph_3SnF ^[25] by elemental analysis and IR spectroscopy (Scheme 4). CS_2 was detected in the reaction mixture by GC/MS and ¹³C NMR experiments. We assume that initially **1a** and CS_2 are formed from **2a** at elevated temperatures and a subsequent reaction of **1a** with Ph_3SnCl yields the chlorido



Scheme 4. Reactivity of **2a** towards Ph_3SnCl and CO .

complex **6** and Ph_3SnF . In addition, heating of **2a** in the presence of CO led also to the formation of CS_2 as well as the formation of *trans*- $[\text{Ir}(4\text{-C}_5\text{NF}_4)(\text{F})(\text{H})(\text{CO})(\text{P}i\text{Pr}_3)_2]$ (**4**; Scheme 4).^[13] However, we did not observe any reaction of **2a** with metal fluorides (CsF , NaF , Me_3SnF , Ph_3SnF), nor did it react with $\text{Et}_3\text{N}\cdot 3\text{HF}$ or Bu_4NF in THF at 50°C to yield fluorodithiocarbonato derivatives.

In conclusion a combined experimental and computational approach revealed that an insertion of an organic substrate into a metal–fluorine bond is feasible, indeed. The insertion of CS_2 into the metal–fluorine bond of the iridium(III) fluorido complexes **1a** and **1b** yields the fluorodithiocarbonato compounds **2a** and **2b**, respectively. Computational studies support an unprecedented metathesis-like concerted mechanism, which requires a vacant coordination site at an electron-deficient metal center. The insertion of unsaturated molecules into transition-metal–fluorine bonds to achieve a C–F bond formation has to be considered as a promising alternative to ligand fluorination.

Received: June 13, 2013

Published online: August 28, 2013

Keywords: density functional calculations · fluorine · iridium · reaction mechanism · structure elucidation

- [1] For reviews on fluorinated materials, see: a) J. Scheirs, *Modern Fluoropolymers* (Ed.: J. Scheirs), Wiley, New York, **1997**; b) K. Johns, G. Stead, *J. Fluorine Chem.* **2000**, *104*, 5–18; c) T. Hiyama, *Organofluorine Compounds. Chemistry and Applications*, Springer, Berlin, **2000**; d) B. Ameduri, B. Boutevin, *Well-Architected Fluoropolymers: Synthesis Properties and Applications*, Elsevier, Amsterdam, **2004**; e) P. Kirsch, *Modern Fluoroorganic Chemistry. Synthesis Reactivity, Applications*, Wiley-VCH, Weinheim, **2004**; f) J. A. Gladysz, *Handbook of Fluorous Chemistry*, Wiley-VCH, Weinheim, **2004**; g) *Fluorinated Materials for Energy Conversion*, (Eds.: T. Nakajima, H. Groult), Elsevier, Amsterdam, **2005**; h) K. Reichenbacher, H. I. Süß, J. Hulliger, *Chem. Soc. Rev.* **2005**, *34*, 22–30; i) R. Berger, G. Resnati, P. Metrangolo, E. Weber, J. Hulliger, *Chem. Soc. Rev.* **2011**, *40*, 3496–3508; j) P. Metrangolo, F. Meyer, T. Pilati, G. Resnati, G. Terraneo, *Angew. Chem.* **2008**, *120*, 6206–6220; *Angew. Chem. Int. Ed.* **2008**, *47*, 6114–6127.

- [2] For reviews on medicinal aspects of fluorine chemistry, see: a) C. Isanbor, D. O'Hagan, *J. Fluorine Chem.* **2006**, *127*, 303–319; b) K. L. Kirk, *J. Fluorine Chem.* **2006**, *127*, 1013–1029; c) K. Müller, C. Faeh, F. Diederich, *Science* **2007**, *317*, 1881–1886; d) J.-P. Bégué, D. Bonnet-Delpon, *Bioorganic and Medicinal Chemistry of Fluorine*, Wiley, Hoboken, **2008**; e) S. Purser, P. R. Moore, S. Swallow, V. Gouverneur, *Chem. Soc. Rev.* **2008**, *37*, 320–330; f) D. O'Hagan, *J. Fluorine Chem.* **2010**, *131*, 1071–1081.
- [3] a) *Fluorine in Agriculture, 9–11 January 1995, Conference Papers* (Ed.: R. E. Banks), Fluorine Technology Ltd., Cheshire, **1994**; b) P. Jeschke, *ChemBioChem* **2004**, *5*, 570–589.
- [4] For examples of applications of ^{18}F in PET, see: a) M. E. Phelps, *Proc. Natl. Acad. Sci. USA* **2000**, *97*, 9226–9233; b) M. J. Adam, D. S. Wilbur, *Chem. Soc. Rev.* **2005**, *34*, 153–163; c) S. M. Ametamey, M. Honer, P. A. Schubiger, *Chem. Rev.* **2008**, *108*, 1501–1516; d) P. W. Miller, N. J. Long, R. Vilar, A. D. Gee, *Angew. Chem.* **2008**, *120*, 9136–9172; *Angew. Chem. Int. Ed.* **2008**, *47*, 8998–9033; e) E. Lee, A. S. Kamlet, D. C. Powers, C. N. Neumann, G. B. Boursalian, T. Furuya, D. C. Choi, J. M. Hooker, T. Ritter, *Science* **2011**, *334*, 639–642; f) M. Tredwell, V. Gouverneur, *Angew. Chem.* **2012**, *124*, 11590–11602; *Angew. Chem. Int. Ed.* **2012**, *51*, 11426–11437; g) E. Lee, J. M. Hooker, T. Ritter, *J. Am. Chem. Soc.* **2012**, *134*, 17456–17458; h) I. S. R. Stenhagen, A. K. Kirjavainen, S. J. Forsback, C. G. Jørgensen, E. G. Robins, S. K. Luthra, O. Solin, V. Gouverneur, *Chem. Commun.* **2013**, *49*, 1386–1388.
- [5] For reviews on transition-metal-mediated fluorination, see: a) V. V. Grushin, *Chem. Eur. J.* **2002**, *8*, 1006–1014; b) C. Bobbio, V. Gouverneur, *Org. Biomol. Chem.* **2006**, *4*, 2065–2075; c) J. M. Brown, V. Gouverneur, *Angew. Chem.* **2009**, *121*, 8762–8766; *Angew. Chem. Int. Ed.* **2009**, *48*, 8610–8614; d) V. Gouverneur, *Science* **2009**, *325*, 1630–1630; e) T. Furuya, J. E. M. N. Klein, T. Ritter, *Synthesis* **2010**, 1804–1821; f) T. W. Lyons, M. S. Sanford, *Chem. Rev.* **2010**, *110*, 1147–1169; g) T. Furuya, A. S. Kamlet, T. Ritter, *Nature* **2011**, *473*, 470–477; h) C. Hollingworth, V. Gouverneur, *Chem. Commun.* **2012**, *48*, 2929–2942; i) T. Liang, C. N. Neumann, T. Ritter, *Angew. Chem.* **2013**, *125*, 8372–8423; *Angew. Chem. Int. Ed.* **2013**, *32*, 8214–8264.
- [6] a) L. Hintermann, A. Togni, *Angew. Chem.* **2000**, *112*, 4530–4533; *Angew. Chem. Int. Ed.* **2000**, *39*, 4359–4362; b) A. Togni, A. Mezzetti, P. Barthazy, C. Becker, I. Devillers, R. Frantz, L. Hintermann, M. Perseghini, M. Sanna, *Chimia* **2001**, *55*, 801–805; c) P. Barthazy, A. Togni, A. Mezzetti, *Organometallics* **2001**, *20*, 3472–3477; d) Y. Hamashima, K. Yagi, H. Takano, L. Tamáz, M. Sodeoka, *J. Am. Chem. Soc.* **2002**, *124*, 14530–14531; e) Y. Hamashima, H. Takano, D. Hotta, M. Sodeoka, *Org. Lett.* **2003**, *5*, 3225–3228; f) Y. Hamashima, T. Suzuki, H. Takano, Y. Shimura, M. Sodeoka, *J. Am. Chem. Soc.* **2005**, *127*, 10164–10165; g) K. L. Hull, W. Q. Anani, M. S. Sanford, *J. Am. Chem. Soc.* **2006**, *128*, 7134–7135; h) T. Suzuki, Y. Hamashima, M. Sodeoka, *Angew. Chem.* **2007**, *119*, 5531–5535; *Angew. Chem. Int. Ed.* **2007**, *46*, 5435–5439; i) T. Suzuki, T. Goto, Y. Hamashira, M. Sodeoka, *J. Org. Chem.* **2007**, *72*, 246–250; j) B. C. Gorske, C. T. Mbofana, S. J. Miller, *Org. Lett.* **2009**, *11*, 4318–4321; k) A. Hazari, V. Gouverneur, J. M. Brown, *Angew. Chem.* **2009**, *121*, 1322–1325; *Angew. Chem. Int. Ed.* **2009**, *48*, 1296–1299; l) W. Wang, J. Jasinski, G. B. Hammond, B. Xu, *Angew. Chem.* **2010**, *122*, 7405–7410; *Angew. Chem. Int. Ed.* **2010**, *49*, 7247–7252; m) S. Qiu, T. Xu, J. Zhou, Y. Guo, G. Liu, *J. Am. Chem. Soc.* **2010**, *132*, 2856–2857; n) P. Tang, T. Furuya, T. Ritter, *J. Am. Chem. Soc.* **2010**, *132*, 12150–12154; o) M. H. Katcher, A. G. Doyle, *J. Am. Chem. Soc.* **2010**, *132*, 17402–17404; p) C. Hollingworth, A. Hazari, M. N. Hopkinson, M. Tredwell, E. Benedetto, M. Huiban, A. D. Gee, J. M. Brown, V. Gouverneur, *Angew. Chem.* **2011**, *123*, 2661–2665; *Angew. Chem. Int. Ed.* **2011**, *50*, 2613–2617; q) M. H. Katcher, A. Sha,

- A. G. Doyle, *J. Am. Chem. Soc.* **2011**, *133*, 15902–15905; r) W. Liu, X. Huang, M.-J. Cheng, R. J. Nielsen, W. A. Goddard III, J. T. Groves, *Science* **2012**, *337*, 1322–1325; s) P. S. Fier, J. Luo, J. F. Hartwig, *J. Am. Chem. Soc.* **2013**, *135*, 2552–2559; t) M.-G. Braun, M. H. Katcher, A. G. Doyle, *Chem. Sci.* **2013**, *4*, 1216–1220.
- [7] a) A. J. Blake, R. W. Cockman, E. A. V. Ebsworth, J. H. Holloway, *J. Chem. Soc. Chem. Commun.* **1988**, 529–530; b) E. A. V. Ebsworth, N. Robertson, L. J. Yellowlees, *J. Chem. Soc. Dalton Trans.* **1993**, 1031–1037; c) S. A. Brewer, K. S. Coleman, J. Fawcett, J. H. Holloway, E. G. Hope, D. R. Russell, P. G. Watson, *J. Chem. Soc. Dalton Trans.* **1995**, 1073–1076; d) J. E. Veltheer, P. Burger, R. G. Bergman, *J. Am. Chem. Soc.* **1995**, *117*, 12478–12488; e) P. Barthazy, L. Hintermann, R. M. Stoop, M. Wörle, A. Mezzetti, A. Togni, *Helv. Chim. Acta* **1999**, *82*, 2448–2453; f) P. Barthazy, R. M. Stoop, M. Wörle, A. Togni, A. Mezzetti, *Organometallics* **2000**, *19*, 2844–2852; g) L. Hintermann, F. Läng, P. Maire, A. Togni, *Eur. J. Inorg. Chem.* **2006**, 1396–1412; h) T. Furuya, H. M. Kaiser, T. Ritter, *Angew. Chem.* **2008**, *120*, 6082–6085; *Angew. Chem. Int. Ed.* **2008**, *47*, 5993–5996; i) T. Furuya, T. Ritter, *J. Am. Chem. Soc.* **2008**, *130*, 10060–10061; j) N. D. Ball, M. S. Sanford, *J. Am. Chem. Soc.* **2009**, *131*, 3796–3797; k) T. Furuya, A. E. Strom, T. Ritter, *J. Am. Chem. Soc.* **2009**, *131*, 1662–1663; l) T. Furuya, T. Ritter, *Org. Lett.* **2009**, *11*, 2860–2863; m) P. Tang, T. Ritter, *Tetrahedron* **2011**, *67*, 4449–4454; n) D. Breyer, T. Braun, P. Kläring, *Organometallics* **2012**, *31*, 1417–1424.
- [8] a) W. J. Marshall, V. V. Grushin, *Organometallics* **2003**, *22*, 555–562; b) V. V. Grushin, W. J. Marshall, *J. Am. Chem. Soc.* **2004**, *126*, 3068–3069; c) W. J. Marshall, V. V. Grushin, *Organometallics* **2004**, *23*, 3343–3347; d) S. A. Macgregor, D. C. Roe, W. J. Marshall, K. M. Bloch, V. I. Bakmutov, V. V. Grushin, *J. Am. Chem. Soc.* **2005**, *127*, 15304–15321; e) V. V. Grushin, W. J. Marshall, *Organometallics* **2007**, *26*, 4997–5002; f) D. V. Yandulov, N. G. Tran, *J. Am. Chem. Soc.* **2007**, *129*, 1342–1358; g) D. A. Watson, M. Su, G. Teverovkiy, Y. Zhang, J. Garzía-Fortanet, T. Kinzel, S. L. Buchwald, *Science* **2009**, *325*, 1661–1664; h) T. Furuya, D. Benitez, E. Tkatchouk, A. E. Strom, P. Tang, W. A. Goddard III, T. Ritter, *J. Am. Chem. Soc.* **2010**, *132*, 3793–3807.
- [9] J. A. Akana, K. X. Bhattacharyya, P. Müller, J. P. Sadighi, *J. Am. Chem. Soc.* **2007**, *129*, 7736–7737.
- [10] We denote the S_2CF^- anion as fluorodithiocarbonate anion and the corresponding ligand fluorodithiocarbonato ligand in accordance to: a) *Nomenclature of Inorganic Chemistry, IUPAC Recommendations 1990* (Ed.: G. J. Leigh), Blackwell Scientific Publications, Oxford, **1990**, pp. 122–142; b) X. Zhang, U. Gross, K. Seppelt, *Angew. Chem.* **1995**, *107*, 2019–2021; *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 1858–1860; c) *Nomenclature of Inorganic Chemistry, IUPAC Recommendations 2005* (Eds.: N. G. Connelly, T. Damhus, R. M. Hartshorn, A. T. Hutton), RSC Publishing, Cambridge, **2005**, pp. 124–141.
- [11] J. A. Evans, M. J. Hacker, R. D. W. Kemmitt, D. R. Russell, J. Stocks, *J. Chem. Soc. Chem. Commun.* **1972**, 72–73.
- [12] For F···F separations in bifluoride ions, see: a) D. Lentz, K. Seppelt, U. Rüdiger, *J. Fluorine Chem.* **1997**, *84*, 103–106; b) S. Y. Troyanov, I. V. Morozov, E. Kemnitz, *Z. Anorg. Allg. Chem.* **2005**, *631*, 1651–1654, and references herein; c) T. Braun, A. Steffen, V. Schorlemer, B. Neumann, H.-G. Stammer, *Dalton Trans.* **2005**, 3331–3335; d) P. Tang, W. Wang, T. Ritter, *J. Am. Chem. Soc.* **2011**, *133*, 11482–11484.
- [13] P. Kläring, A.-K. Jungton, T. Braun, C. Müller, *Eur. J. Inorg. Chem.* **2012**, 1430–1436.
- [14] a) M. Ahijado, T. Braun, D. Noveski, N. Kocher, B. Neumann, D. Stalke, H.-G. Stammer, *Angew. Chem.* **2005**, *117*, 7107–7111; *Angew. Chem. Int. Ed.* **2005**, *44*, 6947–6951; b) M. Ahijado Salomon, T. Braun, I. Krossing, *Dalton Trans.* **2008**, 5197–5206; c) M. Ahijado Salomon, A.-K. Jungton, T. Braun, *Dalton Trans.* **2009**, 7669–7677; d) G. Meier, T. Braun, *Angew. Chem.* **2012**, *124*, 12732–12737; *Angew. Chem. Int. Ed.* **2012**, *51*, 12564–12569.
- [15] P. Kläring, S. Pahl, A. Penner, T. Braun, *Dalton Trans.* **2011**, 40, 6785–6791.
- [16] See the Supporting Information for computational details and a full reference for the Gaussian 09 programme package: Gaussian09, Revision A.02, M. J. Frisch et al., Gaussian Inc., Wallingford CT, **2009**.
- [17] a) N. Kuhn, H. Bohnen, J. Fahl, D. Blaser, R. Boese, *Chem. Ber.* **1996**, *129*, 1579–1585; b) T. Böttcher, B. S. Bassil, G.-V. Rösenthaller, *Inorg. Chem.* **2012**, *51*, 763–765.
- [18] For experimentally determined H–F separations in solid, liquid, and gaseous hydrogen fluoride, see: a) M. Atoji, W. Lipscomp, *Acta Crystallogr.* **1954**, *7*, 173–175; b) M. Deraman, J. C. Core, J. G. Powles, J. H. Holloway, P. Chieux, *Mol. Phys.* **1985**, *55*, 1351–1367; c) T. Pfeleiderer, I. Waldner, H. Bertagnolli, K. Tödheide, H. E. Fischer, *J. Chem. Phys.* **2000**, *113*, 3690–3696; d) S. E. McLain, C. J. Benmore, J. E. Siewenie, J. Urquidi, J. F. C. Turner, *Angew. Chem.* **2004**, *116*, 1986–1989; *Angew. Chem. Int. Ed.* **2004**, *43*, 1952–1955; e) M. Kreitmeir, G. Heusel, H. Bertagnolli, K. Tödheide, C. J. Mundy, *J. Chem. Phys.* **2005**, *122*, 154511–8.
- [19] a) L. A. Watson, D. V. Yandulov, K. G. Caulton, *J. Am. Chem. Soc.* **2001**, *123*, 603–611; b) S. P. Reade, M. F. Mahon, M. K. Whittlesey, *J. Am. Chem. Soc.* **2009**, *131*, 1847–1861; c) J. A. Panetier, S. A. Macgregor, M. K. Whittlesey, *Angew. Chem.* **2011**, *123*, 2835–2838; *Angew. Chem. Int. Ed.* **2011**, *50*, 2783–2786; d) A. Nova, R. Mas-Ballesté, A. Lledós, *Organometallics* **2012**, *31*, 1245–1256, and references herein; e) J.-P. Werner, P. Burger, *J. Phys. Chem. A* **2011**, *115*, 13885–13895; f) D. Lentz, T. Braun, M. F. Kuehnell, *Angew. Chem.* **2013**, *125*, 5328–5332; *Angew. Chem. Int. Ed.* **2013**, *52*, 3328–3348; g) M. F. Kühnel, P. Holstein, M. Kliche, J. Krüger, S. Matthies, D. Nitsch, J. Schutt, M. Sparenberg, D. Lentz, *Chem. Eur. J.* **2012**, *18*, 10701–10714; h) M. F. Kühnel, D. Lentz, *Angew. Chem.* **2010**, *122*, 2995–2998; *Angew. Chem. Int. Ed.* **2010**, *49*, 2933–2936; i) R. P. Hughes, *J. Fluorine Chem.* **2010**, *131*, 1059–1070; j) J. Yuan, R. P. Hughes, A. L. Rheingold, *Organometallics* **2006**, *25*, 2908–2910.
- [20] a) H. Plenio, *Chem. Rev.* **1997**, *97*, 3363–3384, and references herein; b) K. Stanek, B. Czarniecki, R. Aardoom, H. Rügger, A. Togni, *Organometallics* **2010**, *29*, 2540–2546.
- [21] a) S. Erhardt, S. A. Macgregor, *J. Am. Chem. Soc.* **2008**, *130*, 15490–15498; b) A. Nova, S. Erhardt, N. A. Jasim, R. N. Perutz, S. A. Macgregor, J. E. McGrady, A. C. Whitwood, *J. Am. Chem. Soc.* **2008**, *130*, 15499–15511; c) J. Goodman, S. A. Macgregor, *Coord. Chem. Rev.* **2010**, *254*, 1295–1306; d) A. Nova, M. Reinhold, R. N. Perutz, S. A. Macgregor, J. E. McGrady, *Organometallics* **2010**, *29*, 1824–1831; e) V. V. Grushin, *Acc. Chem. Res.* **2010**, *43*, 160–171; f) M. Teltewskoi, J. A. Panetier, S. A. Macgregor, T. Braun, *Angew. Chem.* **2010**, *122*, 4039–4043; *Angew. Chem. Int. Ed.* **2010**, *49*, 3947–3951; g) E. Clot, O. Eisenstein, N. Jasim, S. A. Macgregor, J. E. McGrady, R. N. Perutz, *Acc. Chem. Res.* **2011**, *44*, 333–348; h) E. Q. Jiao, F. T. Xia, H. Zhu, *Comput. Theor. Chem.* **2011**, *965*, 92–100; i) A. L. Raza, J. Panetier, M. Teltewskoi, S. A. Macgregor, T. Braun, *Organometallics* **2013**, *32*, 3795–3807.
- [22] See the Supporting Information for supplementary crystallographic data of **3**. CCDC 941549 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [23] a) T. Kashiwagi, N. Yasuoka, T. Ueki, N. Kasai, M. Kakudo, S. Takahashi, N. Nagihara, *Bull. Chem. Soc. Jpn.* **1968**, *41*, 296–303; b) C. Bianchini, D. Masi, C. Mealli, A. Meli, M. Sabat, *Organometallics* **1985**, *4*, 1014–1019; c) E. Lindner, B. Keppeler,

- H. A. Mayer, K. Gierling, R. Fawzi, M. Steimann, *J. Organomet. Chem.* **1996**, 526, 175–183; d) M. Doux, N. Mezailles, L. Ricard, P. Le Floch, *Organometallics* **2003**, 22, 4624–4626; e) J. S. Anderson, V. M. Iluc, G. L. Hillhouse, *Inorg. Chem.* **2010**, 49, 10203–10207.
- [24] A.-K. Jungton, P. Kläring, T. Braun, A. Eißler, *Z. Anorg. Allg. Chem.* **2012**, 638, 505–511.
- [25] a) B. W. A. Sharp, J. M. Winfield, *J. Chem. Soc.* **1965**, 2278–2279; b) R. C. Poller, *Spectrochim. Acta* **1966**, 22, 935–939; c) A. C. Sau, L. A. Carpino, R. R. Holmes, *J. Organomet. Chem.* **1980**, 197, 181–197.
-