The present results put this idea on very solid ground, demonstrating that alcohols or water can serve as general acids/bases for catalysis of O-O bond cleavage. Therefore, we have provided viable mechanisms for cytochrome P-450 catalysis.

$$RS - H\overline{m} + O_{2} = RSH\overline{m}O_{2} \frac{e^{-}}{[H^{+}]} RS - H\overline{m}OOH \qquad (38)$$

$$RSH\overline{m}OOH + H_{2}O - \left[\begin{array}{c} HOH \\ RSH\overline{m}OOH \end{array} \right] - \left[\begin{array}{c} HOH \\ RSH\overline{m}OOH \end{array} \right]$$

$$RSH\overline{m}OOH + OH^{-} + H_{2}O \qquad (39)$$

$$RSH\overline{m}OOH + H_{3}O^{+} - \left[\begin{array}{c} H_{3}O^{+} \\ RSH\overline{m}OOH \end{array} \right] - \left[\begin{array}{c} H_{3}O^{+} \\ RSH\overline{m}OOH \end{array} \right] - \left[\begin{array}{c} H_{3}O^{+} \\ RSH\overline{m}OOH \end{array} \right]$$

RSHmO + 2H₂O (40)

An attractive alternative process would use the thiolate as a reductant.

 $RSH\bar{m}O_2 \approx RSH\bar{m}OOH \xrightarrow{H_1O^+} RS^{+}Hm^+=O + 2H_2O$ (41)

Conclusions. The effect of concentration and acidities of protic solvents on the rate of reactions of hydroperoxides with iron(III) porphyrins and the solvent isotope effects on these rates have been investigated and shown to be almost identical with those effects on the reaction of hydroperoxides with R₂S as well as with reactions of peracids with hemins. These and previously reported data allow two conclusions to be drawn. First, the reactions of hydrogen peroxide, hydroperoxides, and peracids with iron(III) porphyrins in protic solvents all proceed by heterolytic cleavage whether buffer catalyzed or solvent catalyzed. In the absence of efficient solvent buffer catalysis, the complex HmOOR is likely to cleave in a homolytic fashion.^{14b,37} Second, alcohol and water catalyses of this reaction have been demonstrated, providing a rational mechanism for water-catalyzed breaking of the O-O bond in the cytochrome P-450 catalytic cycle.

Acknowledgment. Support for this by a grant from the National Science Foundation (CHE 87-21364) is gratefully acknowledged.

Molybdenum-Mediated Imido Group Transfer: Stoichiometric and Catalytic Reactions and Structures

Edgar W. Harlan and R. H. Holm*

Contribution from the Department of Chemistry, Harvard University, Cambridge, Massachusetts 02138. Received April 14, 1989

Abstract: The isoelectronic relationship $O^{2-} = RN^{2-}$ implies imido group transfer chemistry related to oxygen atom transfer reactivity, which is extensive for certain elements. This matter has been subjected to its first systematic examination, using the N,N-diethyldithiocarbamate (Et₂dtc) complexes of molybdenum and the N-tosylimido (NTs) group. The imido group transfer reactions to Mo, $MoO(Et_2dtc)_2 + XNR$ and $Mo(CO)_2(Et_2dtc)_2 + XNR$, afford $MoO(NTs)(Et_2dtc)_2$ and $Mo(NTs)_2(Et_2dtc)_2$, respectively, in high yield with the imido group transfer reagents $XNR = C_3H_3NNTs$, PhMeSNTs, Ph₃SbNTs, PhINTs, and TsN₃. MoO(NTs)(Et₂dtc)₂·¹/₂PhMe crystallizes in triclinic space group $P\overline{1}$ with a = 9.212 (2) Å, b = 12.080 (4) Å, c = 14.310 (4) Å, $\alpha = 115.02$ (2)°, $\beta = 95.51$ (2)°, $\gamma = 100.34$ (2)°, and Z = 2; the structure was refined to R = 12.0804.8% Mo(NTs)₂(Et₂dtc)₂ was obtained in monoclinic space group $P2_1/n$ with a = 10.632 (2) Å, b = 16.307 (3) Å, c = 19.023(5) Å, $\beta = 101.29$ (2)°, and Z = 4; the structure was refined to R = 4.0%. The two compounds have a distorted octahedral structure with the multiply bonded groups O and NTs in cis positions. Comparison of these structures with that of MoO₂(Et₂dtc)₂ reveals that the NTs group is a nearly isostructural replacement for oxo. The imido group transfer reactions from Mo, $MoO(NTs)(Et_2dtc)_2 + R_3P$, proceed but with the more basic phosphines favoring imido (R_3PNTs) rather than oxo (R_3PO) transfer. Imido transfers to and from Mo involve the purple intermediates $Mo_2O_2(NTs)(Et_2dtc)_4$ and $MoO(NTs)_2(Et_2dtc)_4$, respectively, which are in presumed equilibrium with Mo(IV,VI) complexes. The complex Mo(NTs)(Et₂dtc)₂, resulting from imido group transfer, was too unstable to be isolated. Group transfers to and from Mo were coupled to form the catalytic reactions XNTs + $R_3P \rightarrow X + R_3PNTs$, which did not proceed at a measurable rate in the absence of Mo(NTs)₂(Et₂dtc)₂. The system based on PhMeSNTs + Ph₂MeP showed 28 turnovers in 7.5 h in chloroform at 35 °C. The reaction PhMeSNTs + 2PhSH \rightarrow PhSMe + PhSSPh + TsNH₂, in which benzenethiol acts as an electron donor, was also catalyzed by Mo-(NTs)₂(Et₂dtc)₂. All known imidometal groups are tabulated, and oxo and imido group transfer reactions of Mo are schematically summarized.

A rapidly expanding research area is that including the preparations, structures, and reactions of compounds containing the metal-ligand multiple bond groups 1-8. This set encompasses



selenido ligands that interact as four- or six-electron donors with

metals in their high oxidation states to afford groups whose bond

lengths are indicative of multiple bonding. Properties of compounds containing groups 1 and 2,^{1,2} 3,³ 4,^{1,4,5} 6,^{1,6,7} and 7⁸ have been summarized. The newest group is 5, of which there are only several examples.9 Of these groups, by far the greatest amount

- (2) Schrock, R. R. Acc. Chem. Res. 1979, 12, 98.
- Dehnicke, K.; Strähle, J. Angew. Chem., Int. Ed. Engl. 1981, 20, 413.
 Nugent, W. A.; Haymore, B. L. Coord. Chem. Rev. 1980, 31, 123.
- (4) Nugeni, W. A.; Haymore, B. E. Cona, Chem. Rev. Loo, 11, 125.
 (5) Chisholm, M. H.; Rothwell, I. P. In Comprehensive Coordination Chemistry; Pergamon Press: New York, 1987; Vol. 2, Part 13.4.
- (6) Holm, R. H. Chem. Rev. 1987, 87, 1401.
 (7) Griffith, W. P. Coord. Chem. Rev. 1970, 5, 459.

(8) Müller, A.; Diemann, E. In Comprehensive Coordination Chemistry; Pergamon Press: New York, 1987; Vol. 2, Parts 16.1 and 16.3.

⁽¹⁾ Nugent, W. A.; Mayer, J. M. Metal-Ligand Multiple Bonds; Wiley-Interscience: New York, 1988.

IMIDOMETAL AND OXO/IMIDOMETAL FUNCTIONAL GROUPS



Figure 1. Imido and oxoimido functional groups established in isolated compounds or inferred in transient species (Mn, Fe) from reactivity properties. Two of the Os entries are molecules rather than groups but are included to show the existence of stable species with Os=O and Os=NR groups in the same molecule.

of structural and reactivity information is available for compounds containing one or more of the M=O bonds, 6. A reaction type of oxometal compounds that is of particular current interest is oxygen atom transfer. The most numerous and best understood examples of this reaction type occur with molybdenum complexes, 6.10 the two most pervasive processes being the primary oxo transfer reaction 1⁶, and the μ -oxo dimerization reaction 2. In the first, compounds XO/X function as oxo donors/acceptors.

$$Mo^{V_1}O_2L_n + X \rightleftharpoons Mo^{V_2}OL_n + XO$$
 (1)

$$Mo^{V_1}O_2L_n + Mo^{V_2}O_L_n = Mo^{V_2}O_3L_{2n}$$
(2)

After oxo, the imido ligand is probably the most common in the set 1-8. There is an increasing number of essentially isostructural molecules that differ only in the substitution of group 6 by the M=NR group 2. Molecules of this type that are pertinent to the present work include MoO₂(Et₂dtc)₂/Mo(NPh)₂- $(Et_2dtc)_2$.¹¹⁻¹³ Similarly, the μ -O bridge may be replaced by μ -NR, as in the pair [Fe(salen)]₂O/[Fe(salen)]₂(Ntol).^{14,15} The known examples of metal-imido or mixed metal-oxo/imido functional groups¹⁶ are shown in Figure 1. As would be expected, their occurrence usually follows those oxidation state and delectron configuration parameters that stabilize the M=O group, 6. These are an effective metal charge sufficient to reduce the basicity of the imido ligand such that it can survive electrophilic attack under the usual conditions of preparation and purification, and the configurations $d^{\leq 4}$, which in the expected orbital energy order for a M=NR unit, would maintain a bond order of at least 2.

A comparison of Figure 1 with an analogous presentation of oxometal functional groups⁶ reveals that there are over a dozen more cases of the latter than the former. For example, there are

(9) (a) Hitchcock, P. B.; Lappert, M. F.; Leung, W.-P. J. Chem. Soc., Chem. Commun. 1987, 1282. (b) For the current state of terminal phosphilo/phosphinidene complexes, cf.: Cowley, A. R.; Barron, A. R. Acc. Chem. Res. 1988, 21, 81. (10) Holm, R. H. Coord. Chem. Rev. 1990, in press.

(11) Abbreviations: Cp, cyclopentadienyl; Et₂dtc, N,N-diethyldithio-carbamate(1-); MeCp, methylcyclopentadienyl; py = pyridine; salen, N,N'bis(salicylidenamino)ethane(2-); tol, p-tolyl, tosyl, p-toluenesulfonyl; TPP, tetraphenylporphyrinate(2-).

(12) Berg, J. M.; Hodgson, K. O. Inorg. Chem. 1980, 19, 2180.
 (13) Haymore, B. L.; Maatta, E. A.; Wentworth, R. A. D. J. Am. Chem.

Soc. 1979, 101, 2063.

(14) Davies, J. E.; Gatehouse, B. M. Acta Crystallogr. 1973, B29, 1934. 15) Nichols, P. J.; Fallon, G. D.; Murray, K. S.; West, B. O. Inorg. Chem. 1988, 27, 2795.

no characterized imido complexes of Ti, Tc, and Ru, but there are three types of oxotechnetium groups and four types of oxoruthenium groups. Further, in comparison with the very large number of metal-mediated oxo transfer reactions, such as reaction 1, the number of imido group transfer reactions is minuscule. Such reactions have been limited to, primarily, the oxidation of metal complexes by organic azides and the oxidation of organic substrates by transient high-valent manganese and iron imidoporphyrin species. In addition, there is evidence for catalytic imido group transfer.¹⁷⁻²² Overall, however, the relatively small number of M=NR groups and the narrow range of known imido group transfer reactivity have not made as apparent as might be desired the utility of imidometal species in stoichiometric and catalytic transformations.

We have begun an investigation of imido group transfer chemistry whose ultimate purpose is to develop and systematize group transfer reactivity. As molybdenum has the best understood oxo transfer chemistry, this metal was selected for the initial studies. Further, imidomolybdenum groups are well established for $M_{2}(VI)$,^{13,23–25} $M_{0}(V)$,^{26,27} and $M_{0}(IV)$.²⁸ Reported herein are an examination of the reactivity of certain substrate molecules X=NR in group transfer, complete characterization of new Mo(VI) imido complexes, and certain reactivity aspects of these molecules.

Experimental Section

Preparation of Compounds.¹¹ The compounds S-methyl-S-phenyl-Ntosylsulfilimine,²⁹ S,S-dimethyl-N-tosylsulfilimine,³⁰ S,S-diphenyl-N-tosylsulfilimine,²⁹ N-1-pyridinium sulfonamidate,³¹ triphenylphosphine N-tosylimine,³² triphenylarsine N-tosylimine,³³ triphenylstibine N-tosylimine,³⁴ cyclohexano-N-tosylaziridine,³⁵ tosyl azide,³⁶ (N-tosylimino)phenyliodinane,³⁷ triethylphosphine N-tosylimine,³⁸ Mo(CO)₃(MeCN)₃,³⁹

- (17) Sharpless, K. B.; Chong, A. O.; Oshima, K. J. Org. Chem. 1976, 41, 177.
- (18) Breslow, R.; Gellman, S. H. J. Chem. Soc., Chem. Commun. 1982, 1400.
- (19) Breslow, R.; Gellman, S. H. J. Am. Chem. Soc. 1983, 105, 6728. (20) Barton, D. H. R.; Hay-Motherwell, R. S.; Motherwell, W. B. J. Chem. Soc., Perkin Trans. 1983, 445.
- (21) Mansuy, D.; Mahy, J.-P.; Dureault, A.; Bedi, G.; Battioni, P. J. Chem.
- Soc., Chem. Commun. 1984, 1161.
 (22) Svastis, E. W.; Dawson, J. H.; Breslow, R.; Gellman, S. H. J. Am. Chem. Soc. 1985, 107, 6427.

(23) Maatta, E. A.; Haymore, B. L.; Wentworth, R. A. D. Inorg. Chem. 1980, 19, 1055.

(24) Bishop, M. W., Chatt, J.; Dilworth, J. R.; Neaves, B. D.; Dahlstrom,

 P.; Hyde, J.; Zubieta, J. J. Organomet. Chem. 1981, 213, 109.
 (25) (a) Dehnicke, K.; Wieher, U.; Fenske, D. Z. Anorg. Allg. Chem.
 1979, 456, 71. (b) Bishop, M. W.; Chatt, J.; Dilworth, J. R.; Hursthouse, M. B.; Jayaweera, S. A. A.; Quick, A. J. Chem. Soc., Dalton Trans. 1979, 914. (c) Chatt, J.; Choukroun, R.; Dilworth, J. R.; Hyde, J.; Vella, P.; Zubieta, J. Transition Met. Chem. 1979, 4, 59. (d) Nugent, W. A.; Harlow, R. L. J. Am. Chem. Soc. 1980, 102, 1759. (e) Chisholm, M. H.; Folting, K.; Huffman, J. C.; Raterman, A. L. Inorg. Chem. 1982, 21, 978. (f) Chou, C. Y.; Huff-

man, J. C.; Maatta, E. A. J. Chem. Soc., Chem. Commun. 1984, 1184.
 (26) Wall, K. L.; Folting, K.; Huffman, J. C.; Wentworth, R. A. D. Inorg.

Chem. 1983, 22, 2366. (27) (a) Chou, C. Y.; Huffman, J. C.; Maatta, E. A. Inorg. Chem. 1986, (a) Chou, C. 1., Huffman, J. C., Waatta, L. A. Inforg. Chem. 1980, 25, 822.
(b) Noble, M. E.; Huffman, J. C.; Wentworth, R. A. D. Inorg. Chem. 1982, 21, 2101.
(c) Edelblut, A. W.; Folting, K.; Huffman, J. C.; Wentworth, R. A. D. J. Am. Chem. Soc. 1981, 103, 1927.
(d) Noble, M. E.; Huffman, J. C.; Wentworth, R. A. D. Inorg. Chem. 1983, 22, 1756.
(e) Glowiak, T.; Sabat, M. J. Cryst. Mol. Struct. 1975, 5, 247.
(28) (a) Dilworth, J. R.; Dahlstrom, P. L.; Hyde, J. R.; Zubieta, J. Inorg. Chim. 464, 1983, 21, 21.

Chim. Acta 1983, 71, 21. (b) Chou, C. Y.; Devore, D. D.; Huckett, S. C.; Maatta, E. A.; Huffman, J. C.; Takusagawa, F. Polyhedron 1986, 5, 301. (c) Bernal, I.; Draux, M.; Brunner, H.; Hoffmann, B.; Wachter, J. Organo-metallics 1986, 5, 655.

(29) Tsujihara, K.; Furukawa, N.; Oae, K.; Oae, S. Bull. Chem. Soc. Jpn. 1969, 42, 2631.

(30) King, C. J. Org. Chem. 1960, 25, 352.

(31) Balasubramian, A.; McIntosh, J. M.; Snieckus, V. J. Org. Chem. 1970, 35, 433.

(32) Franz, J. E.; Osuch, C. Tetrahedron Lett. 1963, 841.

 (33) Wittig, G.; Hellwinkel, D. Chem. Ber. 1964, 97, 769.
 (34) Pinchuk, A. M.; Kuplennik, Z. I.; Belaya, Z. N. J. Gen. Chem. USSR 1976, 46, 2155.

(35) Barton, D. H. R.; Britten-Kelly, M. R.; Ferreira, D. J. Chem. Soc., Perkin Trans. 1 1978, 1682.

(36) Doering, W. von E.; DePuy, C. J. Am. Chem. Soc. 1953, 79, 5955.
(37) Yamada, Y.; Yamamoto, T.; Okawara, M. Chem. Lett. 1975, 361.
(38) Mann, F. G.; Chaplin, E. J. J. Chem. Soc. 1937, 527.

⁽¹⁶⁾ This tabulation is confined to M=NR groups (R = H, alkyl, aryl), whose imido (vs nitrene) character is recognizable by short bond distances corresponding to multiple bonds. It does not include the numerous cases of metal-diazenido, -hydrazido, and related interactions, in which metal and ligand oxidation states may be more ambiguous. For an extensive examination of such compounds, cf.: Johnson, B. F. G.; Haymore, B. L.; Dilworth, J. R. In Comprehensive Coordination Chemistry; Pergamon Press: New York, 1987; Vol. 2, Part 13.3.

and $MoO(Et_2dtc)_2^{40}$ were prepared by published methods. In the preparations below, all manipulations were performed under a purified dinitrogen atmosphere, and all solvents were rigorously dried and degassed prior to use.

 $M_0(CO)_2(Et_2dtc)_2$. The following procedure is more convenient than that reported earlier⁴¹ because it does not require the preparation of $M_0(CO)_4Cl_2$. To a solution of 2.00 g (6.60 mmol) of $M_0(CO)_3(MeCN)_3$ in 30 mL of methanol was added 2.00 g of tetraethylthiuram disulfide (Aldrich). Upon being stirred at room temperature, the mixture became yellow, and after about 15 min, an orange solid appeared. The product was collected by filtration, washed with methanol, and dried in vacuo to afford 2.43 g (82%) of a dark red-purple solid. The compound was identified by its IR spectrum.⁴¹

MoO(NTs) (Et₂dtc)₂·0.5C₆H₃Me. A. With S-Methyl-S-phenyl-Ntosylsulfilimine. A solution of 1.00 g (2.50 mmol) of MoO(Et₂dtc)₂ in 25 mL of dichloromethane was treated with a solution of 0.80 g (2.7 mmol) of the sulfilimine in 10 mL of dichloromethane. The solution rapidly became dark purple and over the course of 2 h changed to dark red-brown. The solution was layered with 25 mL of toluene and was maintained overnight at -20 °C. The solid that separated was collected by filtration, washed with hexane, and dried in vacuo to give 1.20 g (85%) of red-orange crystalline product. IR (Nujol): ν_{MOO} 876 cm⁻¹. ¹H NMR (CDCl₃): δ 1.27 (t, 12), 2.33 (s, 1.5), 2.38 (s, 3), 3.73 (q, 8), 7.1-7.2 (m, 2.5), 7.25 (d, 2), 7.88 (d, 2). UV $\lambda_{max} (\epsilon_M)$ (CH₂Cl₂): 375 (5200), 500 (sh) nm. Anal. Calcd for C_{20.5}H₃₁N₃O₃S₅Mo: C, 39.47; H, 5.01; N, 6.74; S, 25.70; Mo, 15.38. Found: C, 39.25; H, 4.95; N, 6.74; S, 26.95; Mo, 15.59.

B. With Tosyl Azide. The previous procedure on the same scale but with 1.1 equiv of tosyl azide in place of the sulfilimine afforded 1.32 g (94%) of product.

C. With N-1-Pyridinium Sulfonamidate. A mixture of 0.400 g (0.0098 mmol) of MoO(Et₂dtc)₂ and 0.260 g (0.011 mmol) of the amidate was suspended in 20 mL of toluene. The stirred mixture rapidly turned deep purple and then more slowly assumed a yellow-orange color. The yellow-orange solid was collected and recrystallized from dichloromethane/toluene to yield 0.450 g (74%) of product.

D. With Triphenylstibine N-Tosylimine. The reaction and recrystallization were performed as in preparation C but with 1.1 equiv of the stibinimine in place of the pyridinium sulfonamidate and gave 0.380 g (62%) of product.

Mo(NTs)₂(Et₂dtc)₂. To a solution of 2.00 g (4.50 mmol) of Mo-(CO)₂(Et₂dtc)₂ in 30 mL of dichloromethane was added 2.90 g (9.90 mmol) of S-methyl-S-phenyl-N-tosylsulfilimine. Gas evolution was accompanied by a gradual color change from red-purple to red-brown. The solution was stirred for 10 h, and then 30 mL of hexane was added. After storage overnight at -20 °C, the solid was collected by filtration, washed with hexane, and dried in vacuo to afford 2.40 g (73%) of red microcrystalline product. ¹H NMR (CDCl₃): δ 1.24 (t, 12), 2.39 (s, 6), 3.69 (q, 8), 7.20 (d, 4), 7.75 (d, 4). UV λ_{max} (ϵ_M) (CH₂Cl₂): 332 (12800), 400 (sh). Anal. Calcd for C₂₄H₃₄N₄O₄S₆Mo: C, 39.44; H, 4.69; N, 7.67; S, 26.32; Mo, 13.13. Found: C, 39.20; H, 4.59; N, 7.60; S, 26.32; Mo, 13.90.

Similar reactions with other tosylimido donors gave the following isolated yields: N-1-pyridinium sulfonamidate, 60%; triphenylstibine N-tosylimine, 77%; tosyl azide, 75%.

Methyldiphenylphosphine N-Tosylimine. A toluene solution of 0.50 g (2.5 mmol) of Ph₂MeP was warmed to 50 °C and treated dropwise with 0.50 g (2.5 mmol) of tosyl azide. After N₂ evolution ceased, the solution volume was reduced to one-half in vacuo, and the solution was maintained at -20 °C for 4 h. The product was collected by filtration and dried in vacuo to give 0.84 g (91%) of white crystalline solid: mp 83-85 °C. ¹H NMR (CDCl₃): δ 2.29 (d, 3), 2.30 (s, 3), 7.03 (d, 2), 7.38-7.48 (m, 4), 7.50-7.70 (m, 8).

Collection and Reduction of X-ray Data. Diffraction-quality crystals of MoO(NTs)(Et₂dtc)₂·¹/₂PhMe (A, dark orange) and Mo(NTs)₂-(Et₂dtc)₂ (B, red) were obtained by slow evaporation of concentrated dichloromethane/toluene solutions (4:1 v/v). The crystals were sealed in glass capillaries under a dinitrogen atmosphere. Data were collected on a Nicolet P3F diffractometer. Unit cells were determined by using 25 machine-centered reflections over $20^{\circ} \le 2\theta \le 25^{\circ}$. Three check reflections registered every 123 reflections showed no significant decay over the course of the data collections. Data were processed with xTAPE of the SHELXTL program package (Nicolet XRD Corporation, Madison, WI 53711), and empirical absorption corrections were applied with the

Table I. Crystallographic Data^a

	MoO(NTs)(Et ₂ dtc) ₂ ·0.5C ₆ H ₅ Me	Mo(NTs) ₂ (Et ₂ dtc) ₂
formula	C20 5H31 MoN3O3S5	C24H34M0N4O4S6
formula wt	623.71	730.84
a, Å	9.212 (2)	10.632 (2)
b, Å	12.080 (4)	16.307 (3)
c, Å	14.310 (4)	19.023 (5)
α , deg	115.02 (2)	
β , deg	95.51 (2)	101.29 (2)
γ , deg	100.34 (2)	
V, Å ³	1393.0 (7)	3234.2 (1.1)
Z	2	4
space group	PĪ	$P2_1/n$
Project (Pobed)	1.49 (1.51)	1.50 (1.49)
μ , cm ⁻¹	8.4	8.0
$R(F_{a}), \%$	4.79	3.98
$R_{\rm w}(F_{\rm o}^{2}), \%$	4.71	4.81

 ${}^{a}T = 298 \text{ K}, \lambda = 0.71069 \text{ Å} (Mo \text{ K}\alpha).$

Table II. Atom Positional Parameters ($\times 10^4$) for MoO(NTs)(Et₂dtc)₂·0.5C₇H₈

atom	x/a	y/b	z/c
Mo(1)	3326.5 (8)	1961.6 (5)	1468.3 (4)
S(41)	2342 (3)	1428 (2)	2431 (1)
C(41)	971 (9)	2001 (7)	2110 (5)
S(42)	1140 (3)	2634 (2)	1425 (1)
N(41)	-64 (8)	1928 (7)	2368 (5)
C(42)	-271 (14)	1223 (9)	2890 (7)
C(44)	-1409 (19)	2304 (13)	1980 (11)
C(45)	-1199 (16)	2920 (11)	2315 (9)
C(43)	-12 (18)	1557 (12)	3568 (10)
N(21)	4078 (8)	2930 (5)	1711 (4)
S(21)	4510 (3)	3877 (2)	1742 (2)
O(21)	5756 (8)	3918 (6)	1612 (6)
O(22)	4263 (10)	4187 (5)	2405 (4)
C(21)	3498 (10)	4398 (6)	1040 (5)
C(26)	2464 (11)	4827 (7)	1171 (5)
C(22)	3789 (11)	4397 (7)	369 (6)
C(24)	2019 (11)	5292 (6)	-50 (6)
C(23)	3039 (13)	4851 (8)	-173 (5)
C(25)	1718 (11)	5277 (7)	620 (6)
C(27)	1246 (16)	5792 (8)	-653 (7)
S(32)	3143 (3)	2315 (2)	201 (1)
C(31)	2132 (9)	1495 (6)	-39 (5)
S(31)	1968 (3)	898 (2)	680 (1)
N(31)	1543 (8)	1345 (5)	-694 (4)
C(32)	618 (10)	658 (7)	-866 (5)
C(34)	1788 (13)	1845 (9)	-1293 (5)
C(35)	919 (18)	2537 (11)	-1437 (8)
C(33)	1247 (12)	-123 (8)	-1020 (7)
N(11)	4583 (7)	1228 (5)	1752 (4)
S(11)	5236 (3)	397 (2)	2118 (2)
O(11)	6542 (7)	404 (5)	2049 (5)
O(12)	4934 (9)	333 (5)	2819 (4)
C(11)	4479 (10)	-381 (6)	1583 (6)
C(12)	3476 (12)	-801 (7)	1786 (6)
C(16)	4797 (14)	-588 (8)	933 (7)
C(14)	3121 (18)	-1586 (9)	763 (10)
C(15)	4092 (20)	-1206(11)	501 (8)
C(13)	2846 (15)	-1402(9)	1372 (10)
C(17)	2449 (29)	-2278 (13)	282 (15)

program PSICOR. Space groups were uniquely determined by axial symmetries from axial photographs and, for B, by systematic absences. They were confirmed by successful solutions and refinements of the structures. Crystallographic data are presented in Table I.

Structure Solutions and Refinements. Atom scattering factors were taken from the tabulation of Cromer and Waber.⁴² For A, the Mo atom was located by Patterson methods. The position of the Mo atom in B was determined by direct methods (MULTAN). All other non-hydrogen atoms were found by successive Fourier and difference Fourier maps and were refined with the program CRYSTALS. The asymmetric unit of A consists of one molecule of $MOO(NTs)(Et_2dtc)_2$ and one-half of a toluene molecule of solvation. The latter resides on an inversion center and is

 ⁽³⁹⁾ Tate, D. P.; Knipple, W. R.; Augl, J. M. Inorg. Chem. 1962, 1, 433.
 (40) Chen, G. J.-J.; McDonald, J. W.; Newton, W. E. Inorg. Chem. 1976, 15, 2612.

⁽⁴¹⁾ Colton, R.; Scollary, G. R.; Tompkins, I. B. Aust. J. Chem. 1968, 21, 15.

⁽⁴²⁾ Cromer, D. T.; Waber, J. T. International Tables for X-Ray Crystallography; Kynoch Press: Birmingham, England, 1974.

Table III. Atom Positional Parameters (×10⁴) for $Mo(NTs)_2(Et_2dtc)_2$

atom	x/a	y/b	z/c
Mo(1)	2263.2 (9)	3995.1 (8)	2422.7 (7)
S(41)	4489 (3)	5442 (2)	3756 (2)
C(41)	5582 (9)	4368 (9)	3247 (7)
S(42)	4632 (3)	3025 (2)	2160 (2)
N(41)	6991 (8)	4596 (7)	3690 (5)
C(42)	7864 (10)	3630 (10)	3326 (7)
C(44)	7795 (11)	5803 (10)	4575 (7)
C(45)	8607 (13)	6698 (11)	4181 (8)
C(43)	7720 (12)	2856 (12)	3938 (10)
N(11)	1402 (7)	2726 (7)	2691 (5)
S(11)	759 (3)	1250 (2)	2389 (2)
O(12)	-504 (7)	765 (6)	1541 (5)
O(11)	521 (8)	1174 (6)	3334 (5)
C(11)	2202 (10)	488 (8)	1948 (7)
C(13)	4422 (12)	-180 (9)	2319 (8)
C(12)	3260 (11)	451 (10)	2683 (8)
C(14)	4442 (11)	-740 (9)	1258 (8)
C(16)	2232 (10)	-59 (9)	865 (7)
C(15)	3369 (11)	-658 (10)	541 (8)
C(17)	5679 (13)	-1419 (13)	880 (11)
O (1)	1314 (7)	5129 (6)	2963 (5)
S(32)	1023 (3)	2645 (2)	602 (2)
C(31)	2007 (10)	3690 (9)	217 (8)
S(31)	3244 (3)	4952 (3)	1253 (2)
N(31)	1841 (9)	3501 (8)	-778 (6)
C(34)	2751 (12)	4463 (12)	-1043 (9)
C(35)	2016 (15)	5493 (13)	-959 (11)
C(32)	801 (13)	2343 (12)	-1648 (9)
C(33)	-576 (17)	2545 (14)	-1970 (13)

disordered. The toluene methyl group was fixed with half-occupancy. Isotropic refinements converged at R values of 7.1% (A) and 7.8% (B). Because of the large thermal parameters, certain ethyl carbon atoms of B were refined isotropically; all other atoms in both crystals were refined anisotropically. In the final stages of refinement, hydrogen atoms were placed at 0.95 Å from, and assigned thermal parameters 1.2 times those of, the bonded carbon atoms. Final R factors are included in Table I, and positional parameters are collected in Tables II and III.4

Other Physical Measurements. All measurements were made under anaerobic conditions. Absorption spectra were determined on a Perkin-Elmer Lambda 4C spectrophotometer, and infrared spectra were measured on a Perkin-Elmer 599B spectrophotometer. ¹H NMR spectra were obtained on a Brüker AM 250 or 300 spectrometer and were referenced to Me₄Si. ³¹P NMR spectra were measured at 121.4 MHz on a Brüker WM 300 spectrometer and are referenced to external 85% H₃PO₄.

Results and Discussion

Classes of Imido Group Transfer Reactions and Imido Group Donors. Two classes of these reactions can be recognized. The first class of imido group transfer processes, described generally by reaction 3, is an oxo/imido substitution process without change

$$M^{z}O_{a}L_{n} + XNR \rightarrow M^{z}O_{a-1}(NR)L_{n} + XO$$
 (3)

XNR = ArNCO,⁴⁶ ArNSO,⁴⁷ Ph_3PNR ,^{23,48} $(Me_3Si)_2NR^{49}$

$$M^{z}(NR)_{a}L_{n} + XNR \rightleftharpoons M^{z+2}(NR)_{a+1}L_{n} + X \qquad (4)$$

$$M^{z}O_{a}L_{n} + XO \rightleftharpoons M^{z+2}O_{a+1}L_{n} + X$$
 (5)

in the oxidation state of the metal.⁴⁴ Principal imido group donors in this reaction include aryl isocyanates, sulfinylamines, and phosphinimines, and bis(silyl)amines. Selected works using these reagents are cited; other synthetic routes by substitution reactions are summarized elsewhere.^{4,5} The second class of imido group transfer processes is represented by reaction 4, which involves oxidation or reduction of the metal. It is strictly analogous to the primary oxo transfer reaction $5,^6$ of which there are many examples, and with M = Mo, it is the subject of this investigation. Compounds 9-17 have been identified as potential or known donors of the imido group in reaction 4. These include the pyridinium 1-amidates (9), sulfilimines (10), sulfoximes (11), sul-



fonediimines (12),⁴⁵ phosphinimines (13), arsinimines (14), and stibinimines (15), all of which to our knowledge have never been used for this purpose. (N-Tosylimino)phenyliodinane (16) has been employed in the generation of transient manganese and iron imido species whose existence has been inferred from their reactivities,17-22 which find analogy with the behavior of the corresponding oxo complexes. Aryl azides (17) have been the most frequently utilized reagent in reaction 4.15,23,25f,50

In this work, we have investigated the stoichiometric and catalytic imido group transfer reactions of molybdenum, utilizing dithiocarbamate complexes as convenient vehicles in reactions with a variety of imido compounds. Of the members of the preceding set, the reactivities of all but 12, with the invariant substituent N-tosyl,¹¹ have been examined. The related oxo chemistry of these complexes, in the form of reactions 1 and 2, has been worked out in considerable detail.⁶ These reactions proceed readily with a variety of oxo donors and acceptors, and Mo(VI,V,IV) complexes have been isolated. Somewhat similar behavior might be expected in imido systems. At the outset, we note the earlier contributions of Wentworth, Maatta, and their co-workers to the chemistry of imidomolybdenum compounds.^{13,23,25f,26,27a-d,28b,50-52}

Imido Transfer to Mo(II,IV). N-Tosyl derivatives of all imido reagents except 12 have been reported. We have employed most of these compounds in this work. Reactions 6-10 with 1 equiv

	pyNTs	+ ру	(6)
	PhMeS=NTs	+ PhSMe	(7)
$MoO(Et_2dtc)_2 +$		\rightarrow MoO(NTs)(Et ₂ dtc) ₂	
	Ph ₃ Sb=NTs	+ Ph ₃ Sb	(8)
	PhI=NTs	+ PhI	(9)
	TsN ₃	$+ N_2$	(10)

of imido donor proceed smoothly at room temperature, resulting in facile imido group transfer via a deep purple intermediate (vide infra). On a preparative scale in dichloromethane, each of these reactions proceeds in high yield. When examined in situ by ¹H

⁽⁴³⁾ See the paragraph at the end of this article concerning supplementary material available.

⁽⁴⁴⁾ We adopt here a nomenclature analogous to that used for oxo com-plexes and their atom transfer reactions.⁶ When used to designate a ligand, the term "imido" refers to the formal anion RN2-, inasmuch as we know of no persuasive reason why in the molecules of interest here the ligands should be regarded otherwise. Reactions of type 3 in which the metal changes oxidation state by two units result, in effect, in the transfer of a nitrene (RN)group. Some may prefer the description "nitrene" instead of "imido" transfer reactions. For simplicity of nomenclature, we retain the latter usage.

⁽⁴⁵⁾ For a summary of the chemistry of compounds 10-12, cf.: Oae, S.;

^{(46) (}a) Bevore, D. D.; Lichtenhan, J. D.; Takusagawa, F.; Maatta, E. J.
Am. Chem. Soc. 1987, 109, 7408. (b) Nielson, A. J.; Waters, J. M.; Bradley, D. C. Polyhedron 1985, 2, 285. (c) Bradley, D. C.; Hursthouse, M. B.; Abdul, Malik K. M. Nielson, A. J. L. Chem. Soc. 1987, 109, 7408. (c) Bradley, D. C.; Hursthouse, M. B.; Abdul, Malik K. M. Nielson, A. J. Shot, P. L. Chem. Soc. 1993. D. C. Polynearon 1965, 2, 263. (c) Brauley, D. C.; Hurstnouse, M. B.; Abdul Malik, K. M.; Nielson, A. J.; Short, R. L. J. Chem. Soc., Dalton Trans. 1983, 2621. (d) Young, C. G.; Broomhead, J. A.; Boreham, C. J. J. Chem. Soc., Dalton Trans. 1983, 2135. (e) Chen, G. J.-J.; McDonald, J. W.; Newton, W. E. Inorg. Chim. Acta 1980, 41, 49.

 ⁽⁴⁷⁾ Cenini, S.; Pizzotti, M. Inorg. Chim. Acta 1980, 42, 65.
 (48) (a) Maatta, E.; Kim, C. Inorg. Chem. 1989, 28, 624. (b) Chong, A.
 O.; Oshima, K.; Sharpless, K. B. J. Am. Chem. Soc. 1977, 99, 3420. (c) Chatt, J.; Dilworth, J. R. J. Chem. Soc., Chem. Commun. 1972, 549.

⁽⁴⁹⁾ Slawisch, T. Z. Anorg. Allg. Chem. 1970, 374, 291; Naturwissen-schaften 1969, 56, 369. Related reactions employ Me₃SiNHR.⁵

^{(50) (}a) Maatta, E. A.; Wentworth, R. A. D. Inorg. Chem. 1979, 18, 2409. (b) Edelblut, A. W.; Wentworth, R. A. D. Inorg. Chem. 1979, 1110. (c)
 Edelblut, A. W.; Wentworth, R. A. D. Inorg. Chem. 1980, 19, 1110. (c)
 Edelblut, A. W.; Wentworth, R. A. D. Inorg. Chem. 1980, 19, 2006.
 (51) (a) Edelblut, A. W.; Haymore, B. L.; Wentworth, R. A. D. J. Am.
 Chem. Soc. 1978, 100, 2250. (b) Noble, M. E.; Folting, K.; Huffman, J. C.;

Wentworth, R. A. D. Inorg. Chem. 1983, 22, 3671.
 (52) (a) Devore, D. D.; Maatta, E. A. Inorg. Chem. 1985, 24, 2846. (b)

Devore, D. D.; Maatta, E. A.; Takusagawa, F. Inorg. Chim. Acta 1986, 112, 87



Figure 2. Absorption spectra of $MoO_2(Et_2dtc)_2$, $MoO(NTs)(Et_2dtc)_2$, and $Mo(NTs)_2(Et_2dtc)_2$ in dichloromethane solutions; absorption maxima are indicated.

NMR, the reactions were found to be quantitative. No other Mo-containing product was detected. Use of toluene as a cosolvent in these reactions resulted in the isolation of the oxoimido-Mo(VI) complex as the toluene hemisolvate. In refluxing 1,2-dichloroethane, 2-3 equiv of Ph₃PNTs, Ph₃AsNTs, Me₂SNTs, Ph₂SNTs, and PhMeS(O)NTs did not react with MoO(Et₂dtc)₂. Parallel behavior is found with Ph₃PO and several sulfones, which have not been shown to be oxo donors to any Mo(IV) complex. It is also reported that the foregoing two sulfilimines do not react with *tert*-phosphines in DMF at 120 °C.⁵³

The behavior of the imido reactants in reactions 6–8 parallels the behavior of their oxo analogues, which afford high-yield conversions to $MoO_2(R_2dtc)_2$.^{6.54,55} However, Ph₃AsO, unlike Ph₃AsNTs, does oxidize $MoO(Et_2dtc)_2$.⁵⁵ Also, while cyclohexano-*N*-tosylaziridine reacts with the Mo(IV) complex, no cyclohexene was observed, in contrast to the reaction with the epoxide,⁵⁶ and the ¹H NMR spectrum revealed multiple Mocontaining products. Two previously established mechanistic features of oxo and imido group transfer are expected to apply to the reaction systems examined here: reactions that are firstorder in Mo(IV) and XNR,⁵⁴ and retention of the α -N atom of TsN₃ in the NTs group.^{50a}

The bis(imido)molybdenum(VI) complex was accessible in good yield by four-electron oxidation of $Mo(CO)_2(Et_2dtc)_2$ with 2 equiv of the same imido donors as in reactions 6–10. The compound is analogous to $Mo(Ntol)_2(Et_2dtc)_2$, which has been obtained by a similar reaction using *p*-tolyl azide.¹³ Use of 1 equiv of imido donor led to formation of $Mo(NTs)_2(Et_2dtc)_2$ in reduced yield, indicating that reaction 12 is competitive in rate or much faster than reaction 11. Spectra of $MoO_2(Et_2dtc)_2$ (yellow), $MoO-Mo(CO)_2(Et_2dtc)_2 + XNTS \rightarrow$

 $Mo(NTs)(Et_2dtc)_2 + X + 2CO$ (11)

 $Mo(NTs)(Et_2dtc)_2 + XNTs \rightarrow Mo(NTs)_2(Et_2dtc)_2 + X$ (12)

 $(NTs)(Et_2dtc)_2$ (orange), and Mo $(NTs)_2(Et_2dtc)_2$ (red-orange) are compared in Figure 2 for the purpose of identification. Introduction of the tosylimido group causes a red shift of the absorption profile. Imido group transfer reactions to Mo are summarized in Figure 3.

Structures of Molybdenum(VI) Tosylimido Complexes. The structures of $MoO(NTs)(Et_2dtc)_2$,¹/₂PhMe and $Mo(NTs)_2$ -($Et_2dtc)_2$ are shown in Figures 4 and 5, respectively, and selected interatomic distances and angles are contained in Table IV. Both molecules display severely distorted octahedral coordination. The multiply bonded ligands are mutually cis, as expected for a d⁰ complex.⁵⁷ The distortions arise from the large N-Mo-O/N and



Figure 3. Summary of imido group transfer reactions to and from Mo in Et_2dtc complexes.

MoO(NTs)(Et₂dtc)₂ ·1/2 toluene



Figure 4. Structure of $MoO(NTs)(Et_2dtc)_2$ as its toluene hemisolvate, showing 50% probability ellipsoids and the atom numbering scheme.



Figure 5. Structure of $Mo(NTs)_2(Et_2dtc)_2$, showing 50% probability ellipsoids and the atom numbering scheme.

small (mean) dithiocarbamate bite angles, which are 104.8 (3)° and 69° in $MoO(NTs)(Et_2dtc)_2$ and 103.9 (4)° and 72° in

⁽⁵³⁾ Aida, T.; Furukawa, N.; Oae, S. Chem. Lett. 1973, 805.

⁽⁵⁴⁾ Reynolds, M. S.; Berg, J. M.; Holm, R. H. Inorg. Chem. 1984, 23, 3057.

⁽⁵⁵⁾ Lu, X.; Sun, J.; Tao, X. Synthesis 1982, 185.

 ⁽⁵⁶⁾ Moloy, K. G. Inorg. Chem. 1988, 27, 677.
 (57) Tatsumi, K.; Hoffmann, R. Inorg. Chem. 1980, 19, 2656.

Table IV. Selected Interatomic Distances (Å) and Angles (deg)

	MoO(NTs)(Et ₂ dtc) ₂ .	
	0.5C7H8	$Mo(NTs)_2(Et_2dtc)_2$
Mo-O(1)/N(21)	1.703 (6)	1.790 (8)
Mo-N(11)	1.801 (7)	1.797 (7)
Mo-S(31)	2.551 (3)	2.550 (3)
Mo-S(32)	2.432 (2)	2.452 (3)
Mo-S(41)	2.457 (2)	2.444 (3)
Mo-S(42)	2.642 (2)	2.559 (3)
N(11) - S(11)	1.624 (8)	1.617 (8)
N(21) - S(21)		1.611 (8)
C(31) - S(31)	1.72 (1)	1.72 (1)
C(31)-S(32)	1.73 (1)	1.72 (1)
C(41)-S(41)	1.743 (9)	1.74 (1)
C(41) - S(42)	1.706 (9)	1.70 (1)
O(1)/N(21)-Mo-N(11)	104.8 (3)	104.0 (4)
$M_0 - N(11) - S(11)$	153.5 (4)	156.8 (5)
$M_0 - N(21) - S(21)$		165.1 (5)
S(31)-Mo-S(32)	70.16 (8)	69.82 (9)
S(41)-Mo-S(42)	68.74 (8)	69.85 (9)

 $Mo(NTs)_2(Et_2dtc)_2$. In the latter complex, the tosylimido groups are oriented so as to minimize interactions between them, and the two Mo=N distances differ by 0.047 Å.

Comparisons with dimensional parameters of MoO₂(Et₂dtc)₂ and certain molybdenum(VI,V) imido complexes, summarized in Table V, provide several interesting results. MoO(NTs)- $(Et_2dtc)_2$ is only the second Mo complex containing both an oxo and an imido ligand whose structure has been determined. The first is MoO(NH)Cl₂(EtPh₂PO)₂;^{25c} parameters of the MoO(NH) group are comparable with those of $MoO(NTs)(Et_2dtc)_2$. The Mo=O distance of the latter is indistinguishable from that in $MoO_2(Et_2dtc)_2$. Its Mo=N distance is the same as those in $Mo(NTs)_2(Et_2dtc)_2$ and is among the longest known, but the entire range covers only 0.1 Å. The O/N-Mo-O/N angles in these three compounds are essentially identical. In Mo(NTs)₂(Et₂dtc)₂, the two Mo-N-S angles differ by only 8°. The virtually equal values of the smaller angle and that in MoO(NTs)(Et₂dtc)₂ indicate that both angles are not strongly affected by steric effects. Also, these angles are closely comparable to that in seven-coordinate [Mo- $(NSO_2Ph)(Et_2dtc)_3]^+$. The situation is very different from that of Mo(NPh)₂(Et₂dtc)₂, where the two Mo-N-C angles differ by 30°. Further, the correlation in this molecule of the shorter Mo=N distance with the larger angle does not hold in Mo- $(NTs)_2(Et_2dtc)_2$, where the two Mo=N distances are indistinguishable from each other and from the distance associated with the smaller angle in $Mo(Ntol)_2(Et_2dtc)_2$. The trans influence on Mo-S bond lengths by the tosylimido group in MoO(NTs)- $(Et_2dtc)_2$ and Mo(NTs)₂ $(Et_2dtc)_2$ is appreciable and nearly equal in the two compounds (ca. 0.1 Å) and is half as large as the effect of an oxo ligand in the former compound. Examination of all comparative structural data leads to the principal conclusion that the structural features of the Mo=O and M=NTs groups are independently manifested in MoO(NTs)(Et₂dtc)₂. The Mo-N-S bond angles of 154-165° suggest that the tosylimido group is acting as an electron donor intermediate between the four-electron (ca. 120°) and six-electron (ca. 180°) limits.

Imido Transfer from Mo(VI) and Formation of Mo(V) Dimers. Having established imido group transfer to Mo(IV) with certain reagents (9, 10, and 15-17), the reverse process has been examined using *tert*-phosphines as substrates. Reactions 13-17 are pertinent

	$Mo(NTs)(Et_2dtc)_2 + R_3PO$ (1)	3)
MoO(NTs)(Et ₂ dtc) ₂ +	R ₃ P	
	$MoO(Et_2dtc)_2 + R_3PNTs $ (1	4)
$MoO(NTs)(Et_2dtc)_2 +$	$Mo(NTs)(Et_2dtc)_2 \longrightarrow Mo_2O(NTs)_2(Et_2dtc)_2$	c)₄ I5)
MoO(NTs)(Etadtc)a +	$MoO(Et_2dtc)_2 \longrightarrow Mo_2O_2(NTs)(Et_2dtc)_4 (1)$	6)
MoO ₂ (Et ₂ dtc) ₂ +	+ MoO(Et ₂ dtc) ₂ \longrightarrow Mo ₂ O ₃ (Et ₂ dtc) ₄ (1	17)

here and were carried out in chloroform or dichloromethane solutions, in which the results were the same. Phosphorus reaction products with the indicated chemical shifts were identified by ³¹P NMR.

Reaction 13 with excess Ph₃P led to an intermediate deep violet solution (λ_{max} 538 nm), which slowly decayed to a red-brown solution with an essentially featureless visible spectrum. One equivalent of Ph₃PO (+25.0 ppm) was formed. The immediate product of this reaction is interpreted to be Mo(NTs)(Et₂dtc)₂, which is captured in reaction 15 to afford a binuclear purple chromophore. The latter is identified as the indicated μ -oxo Mo(V) complex on the basis of its spectral similarity to Mo₂O-(Ntol)₂(Et₂dtc)₄ (λ_{max} 533 nm in toluene), which has been isolated and characterized by Devore and Maatta.^{52a} Attempts to isolate Mo₂O(NTs)₂(Et₂dtc)₄ by similar means led to a red-brown solution. Its ¹H NMR spectrum revealed an envelope of dithiocarbamate ethyl signals corresponding to several metal-containing products that were not identified.

Treatment of MoO(NTs)(Et₂dtc)₂ with excess Ph₂MeP resulted in the formation of Ph₂MePO (+30.0 ppm) and Ph₂MePNTs (+14.6 ppm) in a ca. 2:1 ratio; i.e., reactions 13 and 14 both proceed. A deep violet intermediate is also observed and is due to combined products of reactions 15 and 16. When MoO-(NTs)(Et₂dtc)₂ is reacted with stoichiometric Et₃P, only reaction 14 occurs to a detectable extent, affording Et₃PNTs (+37.8 ppm) and MoO(Et₂dtc)₂, which was detected spectrophotometrically (λ_{max} 385, 508 nm). Here again a violet intermediate is generated (λ_{max} 514 nm). Reaction 16 was confirmed by combining equimolar quantities of the authentic Mo(IV) and Mo(VI) complexes, which produced the identical chromophore. This is the same reaction that produces the violet intermediate in imido group transfer to Mo(IV).

Reactions of $Mo(NTs)_2(Et_2dtc)_2$ with phosphines were not encouraging. With 30 mM complex and 5 equiv of Ph_3P , the formation of Ph_3PNTs (+10.5 ppm) is less than 5% complete after 24 h. With 1 equiv of Ph_2MeP , the reaction affords quantitative formation of $Ph_2MePNTs$ in 5 h. The ¹H NMR spectrum indicated the presence of several Mo-containing products and no evidence, from a visible color or absorption band, was found for a μ -NTs product. A reaction system containing 5 equiv of Ph_2MeP resulted in bleaching.

The formation of a bridged intermediate in reactions 13 and 14 is entirely analogous to the intervention of purple Mo_2O_3 -(Et₂dtc)₄ (λ_{max} 512 nm in 1,2-dichloroethane⁵⁴) in oxo transfer

Table V.	Comparison	of Selected Bond	Lengths (A	 and 	Angles	(deg) f	for Molybdenum	Oxo	/Imido	Complexes
----------	------------	------------------	------------	-------------------------	--------	---------	----------------	-----	--------	-----------

				Mo—S ^a				
complex	Мо—О	Mo=N	cis O/N	trans N	trans O	O/N—Mo—O/N	Mo-N-C/S	ref
$MoO_2(Et_2dtc)_2$	1.703 (2)		2.450 (1)		2.639 (1)	105.8 (1)		12
$MoO(NTs)(Et_2dtc)_2$	1.703 (6)	1.801 (7)	2.44 (1)	2.551 (3)	2.642 (2)	104.8 (3)	153.5 (4)	с
$Mo(NPh)_2(Et_2dtc)_2$		1.754 (4)	2.457 (6)	2.602 (2)		103.5 (2)	169.4 (4)	13
		1.789 (4)		2.755 (2)			139.4 (4)	
$Mo(NTs)_2(Et_2dtc)_2$		1.790 (8)	2.448 (6)	2.559 (3)		104.0 (4)	165.1 (5)	с
		1.797 (7)		2.550 (3)			156.8 (5)	
$Mo(NPh)Cl_2(Et_2dtc)_2$		1.734 (4)	2.508 (6)				166.8 (3)	23
$[Mo(NSO_2Ph)(Me_2dtc)_3]^+$		1.70 (2)	2.49 (2) ⁶				161 (1)	24
$[Mo(NCPh_3)(Me_2dtc)_3]^+$		1.731 (2)	2.498 (9)	2.606 (2)			175.1 (5)	24
$Mo(Ntol)(S_2P(OEt)_2)_3$		1.732 (4)	2.49 (3)	2.645 (2)			168.4 (4)	27b

^a Distances cis or trans to the indicated atoms; mean values given where appropriate. ^b Irregular structure; mean of all values. ^c This work.



Figure 6. Time course (to 7.5 h and 29 turnovers) of the Mo-catalyzed tosylimido group transfer between PhMeSNTs and Ph₂MeP in chloroform solution at 35 °C as observed by ³¹P NMR spectroscopy. Initial concentrations were [PhMeSNTs] = 700 mM, [Ph₂MeP] = 350 mM, and [Mo(NTs)₂(Et₂dtc)₂] = 9.0 mM. Chemical shifts and signal assignments are indicated.

reactions.⁶ This compound exists in equilibrium 17,⁵⁸ whose reversibility may be expected to extend to reactions 15 and 16. While an imido-bridged product may not be formed in these reactions, there are precedents for the unsupported μ -NR group, among them [Fe(salen)]₂(Ntol).^{15,59} It should be noted that Mo₂O₂(NTs)(Et₂dtc)₄ could dissociate into MoO₂(Et₂dtc)₂ and $Mo(NTs)(Et_2dtc)_2$ as well as the species in reaction 16. In this event, Mo(NTs)₂(Et₂dtc)₂ and R₃PO (with all phosphines) would be expected upon imido transfer to and from Mo, respectively. These species were not observed. In the only previous example of imido group transfer to a phosphine, Cr(Ntol)(TPP) and Ph₃P yielded Cr(TPP) and Ph₃PNtol.⁶⁰ While one of the immediate products of reaction 13 is Mo(NTs)(Et₂dtc)₂, this compound could not be isolated because of its instability, despite a number of attempts to do so. Indeed, Devore and Maatta^{52a} were unable to isolate pure Mo(Ntol)(Et₂dtc)₂ from the reaction of MoO- $(Ntol)(Et_2dtc)_2$ and excess Ph₃P. The compound was ultimately trapped and crystallographically characterized as the dimethylacetylene dicarboxylate adduct.⁵² Our inability to isolate $Mo_2O(NTs)_2(Et_2dtc)_4$ is a likely consequence of the instability of Mo(NTs)(Et₂dtc)₂, whose decomposition would remove the binuclear complex from the system.

Catalytic Imido Transfer. Successful imido group transfer to and from Mo suggests the coupling of reactions 18 and 19 to form a catalytic cycle with net reaction 20. In the catalytic system,

$$Mo(NTs)_2(Et_2dtc)_2 + R_3P \rightarrow Mo(NTs)(Et_2dtc)_2 + R_3PNTs$$
(18)

 $Mo(NTs)(Et_2dtc)_2 + XNTs \rightarrow Mo(NTs)_2(Et_2dtc)_2 + X$ (19)

$$XNTs + R_3P \rightarrow X + R_3PNTs$$
 (20)

reaction 19, rendered sufficiently rapid by a large excess of substrate, should arrest decomposition of the unstable imidomolybdenum(IV) complex. This expectation has been realized.

The system in Figure 6, initially containing the mole ratio PhMeSNTs:Ph₂MeP:Mo(NTs)₂(Et₂dtc)₂ = 78:39:1, was maintained at 35 °C and was monitored periodically by ³¹P NMR. After 7.5 h, the reaction was 72% complete, representing 28 turnovers of sulfilimine substrate relative to the Mo complex. No other signals were observed in the range -40 to +50 ppm. The final solution was examined by ¹H NMR and found to contain

equimolar amounts of PhSMe (δ 2.46) and Ph₂MePNTs (δ 2.29, d), with the indicated methyl shifts. Other than unreacted PhMeSNTs (δ 2.81) and Ph₂MeP (δ 1.59, d), no additional species were observed. The catalytic reaction could have been continued for longer periods inasmuch as there was no sign of degradation at 7.5 h. A similar system containing Ph₃P evidenced reaction but at a very much slower rate. The system in Figure 6 without the Mo complex showed no reaction after 24 h. Earlier, a number of phosphines, including Ph₃P, had been demonstrated to reduce quantitatively arylalkyl-N-tosylsulfilimines at 120 °C in DMF solutions.⁶¹

A chloroform solution containing initially 450 mM Ph₃SbNTs, 400 mM Ph₂MeP, and 9.0 mM Mo(NTs)₂(Et₂dtc)₂ was maintained at 25 °C and monitored by ³¹P NMR. After 6 h, formation of Ph₂MePNTs was 79% complete, equivalent to 35 turnovers. Again, no degradation of the system was observed. A control reaction in the absence of the Mo complex revealed a slower reaction, with 22% conversion of phosphine to phosphinimine at 24 h. It is clear that the Mo complex catalyzes the reaction.

Benzenethiol was examined as an alternative electron donor in the catalytic sulfilimine reduction, reaction 21. A system initially containing 150 mM PhMeSNTs, 200 mM PhSH, and

$$PhMeS = NTs + 2PhSH \rightarrow PhSMe + PhSSPh + TsNH_2$$
(21)

10 mM Mo(NTs)₂(Et₂dtc)₂ in chloroform at room temperature was monitored by ¹H NMR. After 56 h, all the thiol had been consumed, as evidenced by the disappearance of the SH resonance at δ 3.40, a 67% conversion to the sulfilimine to the sulfide had occurred, and equal quantities of the sulfide and sulfonamide had been produced. The reaction proceeds because Mo(NTs)₂(Et₂dtc)₂ is reduced to Mo(NTs)(Et₂dtc)₂ by the thiol.⁶² A chloroform solution of the sulfilimine and PhSH, both at 200 mM, showed no reaction after 3 days at 25 °C.⁶³ Earlier, we have demonstrated the Mo-catalyzed reduction of sulfoxides using benzenethiol as an electron source.⁶⁴

Lastly, MoO(Ntol)(Et_2dtc)₂ has been shown to catalyze the oxidation of Ph₃P by Me₂SO, presumably by the intermediacy of Mo(Ntol)(Et_2dtc)₂.^{28b,52a} Similar reactivity of MoO(NTs)-(Et_2dtc)₂ was tested in a system initially containing 150 mM Me₂SO, 100 mM Ph₃P, and 10 mM complex in chloroform. After 24 h at room temperature, all Ph₃P had been converted to Ph₃PO, indicating the ability of Mo(NTs)(Et_2dtc)₂ to react with oxo as well as imido donors.

Summary. The following are the principal findings and conclusions from this investigation.

1. N-Tosylimido group transfer reactions to Mo(II) (reactions 11 and 12) and oxomolybdenum(IV) in high yield using reagents 9, 10, and 15-17 have been developed.

2. Oxygen atom and N-tosylimido group transfer reactions from (oxoimido)molybdenum(VI) to *tert*-phosphines proceed, but with the more basic phosphines⁶⁵ favoring imido rather than oxo transfer (reactions 13 and 14).

3. The comparative structures of the set $MoXY(Et_2dtc)_2$ with X = Y = O, X = Y = NTs, and X = O, Y = NTs show that the metric features of the MoXY unit are almost independent of X and Y; tosylimido is a nearly isostructural replacement for oxo.

4. Reactions 1 and 2 involve the purple intermediates with probable bridge units $Mo_2O(NTs)(\mu-O)$ and $Mo_2(NTs)_2(\mu-O)$, respectively, analogous to the structurally established $Mo_2O_2(\mu-O)$ bridge in $Mo_2O_3(Pr_2dtc)_4^{66}$ and that present in isolated Mo_2O_3

⁽⁵⁸⁾ Matsuda, T.; Tanaka, K.; Tanaka, T. Inorg. Chem. 1979, 18, 454.

⁽⁵⁹⁾ For an example of an imido/oxo-bridged Mo structure, cf. ref 51a. (60) Elliott, R. L.; Nichols, P. J.; West, B. O. Polyhedron **1987**, 6, 2191.

⁽⁶¹⁾ Aida, T.; Furukawa, N.; Oae, S. C. J. Chem. Soc., Perkin Trans. 2 1976, 1438.

⁽⁶²⁾ The analogous reduction of MoO₂(Et₂dtc)₂ to MoO(Et₂dtc)₂ by PhSH has been demonstrated: McDonald, J. W.; Corbin, J. L.; Newton, W. E. Inorg. Chem. **1976**, 15, 2056.

⁽⁶³⁾ In a different, uncatalyzed transformation, RR'SNTs with ArS⁻ yields RSSAr and TsNH₂: Oae, S.; Aida, T.; Nakajima, M.; Furukawa, N. *Tetrahedron* **1974**, *30*, 947.

⁽⁶⁴⁾ Caradonna, J. P.; Harlan, E. W.; Holm, R. H. J. Am. Chem. Soc. 1986, 108, 7856.

⁽⁶⁵⁾ Bush, R. C.; Angelici, R. J. Inorg. Chem. 1988, 27, 681.

OXO TRANSFER REACTIONS OF MOLYBDENUM



IMIDO GROUP TRANSFER REACTIONS OF MOLYBDENUM



Figure 7. Generalized representations of the established oxo and imido group transfer reactions of Mo. Double-headed curved arrows indicate that transformations between species connected with joined straight line segments are bidirectional (but not with the same reagent); single-headed arrows indicate a unidirectional transformation. Allowance is made for the possible existence of bridged units Mo₂(NR)₃ and bridging groups μ -NR, although neither has yet been proven.

(Ntol)₂(Et₂dtc)₄.^{52a} The Mo(IV) complex Mo(NTs)(Et₂dtc)₂, produced in reactions 11 and 13 and implicated in equilibrium 15, proved too unstable for isolation in the pure form.

5. The catalytic reactions 20 with XNR = PhMeSNTs or Ph_3SbNTs and $R_3P = Ph_2MeP$ and reaction 21 in which a thiol serves as an electron donor-all initiated by the catalyst Mo-(NTs)₂(Et₂dtc)₂-were demonstrated. Systems are robust and could have been operated for much longer times than those used here

All known oxygen atom and imido group transfer reactions of Mo are summarized in Figure 7 as transformations between oxidation states and oxo/imido Mo groups. Here XO/X and XNR/X are oxo and imido donors/acceptors, single- and double-headed curved arrows represent uni- and bidirectional transformations between species connected by joined straight-line segments, and double arrows link species in known or potential equilibrium. The schemes are generalized; not all transformations are spontaneous with a given Mo complex or donor/acceptor. Oxo transfer reactions are presented and discussed at length elsewhere.^{6,10} Imido group transfer reactions are drawn from this work and previously cited references. Schemes such as those in Figure 7 provide a unified presentation of all known transformations by the processes of atom or group transfer. They represent a desirable

(66) Ricard, L.; Estienne, J.; Karagiannidis, P.; Toledano, P.; Fisher, J.; Mitschler, A.; Weiss, R. J. Coord. Chem. 1973, 3, 277.

goal in the evolvement of the atom or group transfer chemistry of a given element, for they summarize known reactions and imply possible reactions of other compounds of the element.

The imido group transfer reactions to and from Mo demonstrated here show more clearly than heretofore that a significant body of such reactions is possible for Mo and, by implication, for other elements. Further, catalytic imido group transfer can be improved and expanded using the results here as a base. Earlier, catalytic amidation of cyclohexane and alkenes, primarily using manganese and iron porphyrins, had been achieved but with limited turnovers.¹⁷⁻²¹ The active intermediate imidometal species have not been isolated or otherwise detected.⁶⁷ Lastly, inspection of Figure 1 reveals untouched areas for investigation. For example, given the extensive oxo chemistry of Ru,⁶ imido-Ru complexes are especially conspicuous by their absence. The most recent addition to the table of imidometal groups is the highly reactive Zr=NR^{68,69} and Ir=NR.⁷⁰ One example of each has been structurally characterized.^{68,70} In an inverted contrast, the Zr=O and Ir=O groups remain elusive. Metal-imido multiple bonds have very recently been extended to the actinides, in compounds such as $(MeCp)_3UNPh^{71}$ and Cp_3MNPPh_3 (M = Th, U).⁷²

Other metal-ligand multiply bonded units collected at the outset remain attractive subjects for investigations of atom and group transfer reactions. For example, the first instance of nitrogen atom transfer, from a Mn^{V} =N porphyrin to metal(II,III) porphyrins, has been reported.⁷³ There is also at least one example of what might be termed an alkylidene group transfer reaction. Treatment of Cp₂TaMe(Me₃P) with the ylide Et₃PCHMe results in the transfer of a carbene equivalent to the metal, yielding Cp₂TaMe(CHMe).⁷⁴ The availability of sulfonium⁷⁵ and iodonium⁷⁶ ylides suggests extensions of this reaction type.

Acknowledgment. This research was supported by NSF Grant CHE 85-21365. X-ray equipment was obtained through NIH Grant 1 S10 RR-02247. We thank Dr. B. S. Snyder for experimental assistance and useful discussions.

Supplementary Material Available: Tables of crystallographic data, positional and thermal parameters, and calculated hydrogen atom positions for MoO(NTs)(Et₂dtc)₂ and Mo(NTs)₂(Et₂dtc)₂, and details of the data collection (13 pages); table of calculated and observed structure factors (38 pages). Ordering information is given on any current masthead page.

(71) Brennan, J. G.; Andersen, R. A. J. Am. Chem. Soc. 1985, 107, 514. (72) Cramer, R. E.; Edelmann, F.; Mori, A. L.; Roth, S.; Gilje, J. W.; Tatsumi, K.; Nakamura, A. Organometallics 1988, 7, 841. (73) (a) Woo, L. K.; Goll, J. G. J. Am. Chem. Soc. 1989, 111, 3755. (b)

York, 1975.

(76) (a) Schank, K.; Lick, C. Synthesis 1983, 392. (b) Mansuy, D.; Battioni, J. P.; Akhrem, I.; Dupre, D.; Fischer, J.; Weiss, R. J. Am. Chem. Soc. 1984, 106, 6112. (c) Hadjiarapoglou, L.; Spyroudis, S.; Varvoglis, A. J. Am. Chem. Soc. 1985, 107, 7178.

⁽⁶⁷⁾ For recent work related to this point, cf.: Mahy, J.-P.; Battioni, P.; Bedi, G.; Mansuy, D.; Fischer, J.; Weiss, R.; Morgenstern-Badarau, I. J. Am. Chem. Soc. 1988, 27, 353.

⁽⁶⁸⁾ Walsh, P. J.; Hollander, F. J.; Bergman, R. G. J. Am. Chem. Soc. 1988, 110, 8729.

⁽⁶⁹⁾ Cummins, C. C.; Baxter, S. M.; Wolczanski, P. T. J. Am. Chem. Soc. 1988, 110, 8731.

⁽⁷⁰⁾ Glueck, D. S.; Hollander, F. J.; Bergman, R. G. J. Am. Chem. Soc. 1989, 111, 2719.